

## CORRECTION



# Correction: Mortality attributable to sugar sweetened beverages consumption in Mexico: an update

Ariela Braverman-Bronstein, Dalia Camacho-García-Formentí, Rodrigo Zepeda-Tello, Frederick Cudhea, Gitanjali M. Singh, Dariush Mozaffarian and Tonatiuh Barrientos-Gutierrez

© The Author(s), under exclusive licence to Springer Nature Limited 2024

*International Journal of Obesity* (2024) 48:747; <https://doi.org/10.1038/s41366-024-01498-3>

Correction to: *International Journal of Obesity* <https://doi.org/10.1038/s41366-019-0506-x>, published online 10 December 2019

In the section “SSBs intake”, the following sentence:

“We defined SSBs-servings (8 oz) according to the following categories: sugar-sweetened carbonated beverages, industrialized juices including added sugar, and industrialized “aguas frescas” including added sugar. These include all beverages subject to the SSBs tax implemented in Mexico in 2014 [16], but exclude homemade and restaurant-made preparations which represent up to 10.5% of SSBs in Mexico [11]”

has been corrected as follows to include all beverages and not just industrialized preparations:

“We defined SSBs-servings (8 oz) according to the following categories: sugar-sweetened carbonated beverages; industrialized juices and “aguas frescas” with added sugar; prepared juices and “aguas frescas” with added sugars made at home or in a food establishment.”

The original article has been corrected.