

SCIENTIFIC SERIALS

THE *American Naturalist* for February, among others, contains an article by Dr. Gill on "The Limits of the Class of Fishes," in which he endeavours to modify their generally accepted classification by dividing them up into two classes and three sub-classes, of equal significance with the reptiles and birds. The names he proposes are (1) Pisces; (2) Marsipobranchii; and (3) Leptocardii, which sufficiently indicate the genera he includes in each class. Such an amount of division we think excessive, and it would undoubtedly necessitate the removal of the crocodiles from the reptilia, among other changes. Mr. A. S. Packard gives an account of one of the beaks of a cuttle-fish, probably *Architeuthis dux*, which is four and a half inches long; he also describes other colossal specimens. There is a paper by Prof. Jordan on the colours of vegetation, one by Dr. Abbott on the habits of certain crawfish, and another by Dr. Foster on the pottery of the mound-builders, which is fully illustrated.

THE Munich *Zeitschrift für Biologie*, Bd. 8, Heft 4, contains the following papers of purely medical interest: on the occurrence of enteric fever in the Bavarian army, by Dr. Port, with charts of the mortality in the different barracks and of the amount of subsoil water; on the present state of the cholera problem, by Prof. von Pettenkofer; and on the processes of decomposition which result from venesection, by Dr. J. Bauer.

Schriften der Naturforschenden Gesellschaft in Danzig, New Series, vol. 3, Part I. The first paper in this publication of the Danzig Society is a contribution to primitive German history by Dr. Lissaicer of Danzig, being a very careful and elaborate monograph on some skulls found at Meisterswalde and Krissau, a short distance from Danzig. The paper is accompanied by some capitally executed photographs of the skulls. The next paper is also a contribution to the history of the early inhabitants of Pomerania, being a description by Herr Kasiski of the numerous and varied contents of some of the ancient graves which abound in the district around the village of Persanzig, on the river Persante, a short distance west of Neustettin. The district abounds with material for the archaeologist. The paper is accompanied with numerous illustrations of the contents of the graves. The next paper is a long one by Dr. C. J. H. Lampe, of Danzig, on the Movement of Water in pipes, accompanied by some calculations as to the pressure and speed of the water in the pipes by which Danzig is now supplied with water from a considerable distance. This paper is also illustrated, as is also the last one, which is the fifth part of A. Menge's Catalogue of Prussian Spiders.

Der Zoologische Garten (Frankfurt a. M.), January 1873, contains an excellent article, with maps in illustration, of the geographical distribution of the Birds of Paradise, with which are included *Epimachus* and *Ptiloris*. There is also an article by Dr. H. Dörner on the tongue of the Ka-ka Parrot (*Nestor meridionalis*), in which he shows clearly that in structure it presents none of the characters of the *Trichoglossinae*, and in other points his results quite agree with those read before the Zoological Society of London in June last, although he, following Dr. Finsch, does not feel disposed to remove this parrot from among those with trichoglossal tongues, because of a supposed similarity in their beaks, which we find it difficult to appreciate, the Ka-ka's being black and ribbed, whilst that of *Lorius* is smooth and with an orange tint. There is not the least doubt that, now it has been doubly demonstrated that their tongues are not similarly constructed, there is not any good reason for associating the Nestors with the Lories.

SOCIETIES AND ACADEMIES

LONDON

Royal Society, March 6.—"On the Vapour-density of potassium."—Preliminary notice. By James Dewar and William Dittmar.

The results of their observations conclusively show that the density of potassium-vapour, as produced in the process described, cannot exceed 45 times that of hydrogen, and that therefore the molecule of potassium consists of two atoms (K_2).

"On New Sources of Ethyl- and Methyl-Aniline." By John Spiller, F.C.S.

"On a new genus of Amphipod Crustaceans. By Rudolph von Willemoes-Suhm, Ph.D., Naturalist to the *Challenger* exploring expedition.

In lat. $35^{\circ} 47'$, long. $8^{\circ} 23'$, off Cape St. Vincent, the trawl was sent down to a depth of 1090 fathoms on the 28th of January and brought up among other very interesting things a large, transparent Amphipod with enormous faceted eyes. The animal evidently hitherto unknown, will be the type of a new genus, having the following characters:—

THAUMOPS, nov. gen.

Caput oblongum, inflatum, oculis maximis superiorem capitis partem tegentibus. Segmenta thoracica 6, abdominalia 5. Antennarum in feminis par unum, maxillarum par unum, pedum paria duo minima maxillarum locum tenentia. Mandibulæ nullæ. Pedes thoracici 5, abdominalia 3 in quoque latere. Appendices caudales 4. Gangliorum pectoralium paria 5, abdominalium 3.

T. pellucida, n. sp.

Corpus longitudine 14 mm., latitudine 21 mm., pellucidum.

It could not be made out whether *T. pellucida* inhabits the deep sea, or whether it is, like *Phronima*, a pelagic animal, having been caught by the trawl only as the latter came up from the depth.

Geological Society, February 26.—Prof. Ramsay, F.R.S., vice-president, in the chair.—The following communications were read:—"On the Jurassic Rocks of Skye and Raasay," by Dr. James Bryce. In this paper the author described numerous sections of Jurassic rocks exposed chiefly in the sea-cliffs of Skye and Raasay, indicating the presence in those islands of a complete series of beds ascending from the Lower Lias to the middle of the Middle Oolite. He noticed the occurrence in these sections of fossils belonging to the zones of *Ammonites angulatus* and *A. Buchlandi* in the Lower Lias, to the zones of *A. Jamesoni*, *A. capricornus*, *A. margaritatus*, and *A. spinatus* in the Middle Lias, of Upper Lias fossils, including *Anmonites communis*, *falcifer*, *heterophyllus*, and *bifrons* and of others indicating beds belonging to the Inferior Oolite and Cornbrash, and to the Oxford Clay. The Loch Staffin beds were described as an estuarine series, nearly approaching the Oxford Clay in geological age, and including a bed almost entirely made up of shells of *Ostrea hebridica*. The whole series of Jurassic rocks in these islands reposes on the Torridon sandstone of Cambrian age; and the author discussed the question whether or not the intervening beds have ever existed in this locality, and came to the conclusion that they probably existed, and have been swept away by denudation. He remarked further upon the resemblance in lithological characters of the beds described with the corresponding deposits elsewhere in Britain. The traprocks intruded between the Jurassic deposits he regarded as of post-oolitic date.—"Observations on the more remarkable Boulders of the North-West of England and the Welsh Borders," by Mr. D. Mackintosh. In this paper the author described the situation and indicated the probable origin of many of the more striking known boulders in Westmoreland, Cumberland, Lancashire, Cheshire, and on the borders of Wales. The northern boulders seem to have originated chiefly from Westdale Crag, Criffel, Ennerdale, and Eskdale; those of Cheshire chiefly from the Lake District and South of Scotland; and many of those on the Welsh borders from the mountains of Wales. Many of the boulders noticed by the author exhibit glacial striae. The author also especially referred to the occurrence of boulders at high levels.

Linnæan Society, March 6.—Mr. Bentham made some observations on the homology of the perigynium or utricle of the female flowers of *Carex* and *Uncina*, with a view to calling to the disputed points in question the attention of botanists used to microscopical investigation, who may have the opportunity of examining living specimens in the earliest stages of flowering. Two principal explanations of the homology of the perigynium of *Carex* have been given. Brown, relying upon its being composed of two squamæ, considered that it represents a perianth, and Payer and Schleiden have adopted the same view, after an examination of its appearance at a very early stage. Kunth, on the contrary, believed it to be formed of a single scale, and to be an ordinary glume subtending the female flower on a secondary axis, of which the seta of many species of *Carex*, and of