

of the combination of "the pleasure of intense stimulus" and the "pleasure of conflict" in our enjoyment of a tragedy, are models of delicate æsthetic analysis. The author's attitude towards the various current theories of "play" is eminently judicious. As he well points out, both the "surplus activity" theory and the "recreation" theory are one-sided, the former doing less than justice to the pastimes of adults, the latter to those of children. His own view that play must be regarded by the biologist primarily as the great educator and perfecter of imperfect instincts has been most nearly approached by Prof. Baldwin. Prof. Groos's treatment of the sociological aspects of "play," both as the child's earliest form of experimentation and as the earliest school of obedience to authority, should prove useful to students of ethics as well as to professed sociologists. The admirable literary style of the book, no less than the interest of its contents, should recommend it to all persons of general culture who care for anthropological studies.

A. E. T.

*Physique et Chimie Viticoles.* By A. de Saporta. Pp. iv + 300. (Paris: G. Carré and C. Naud, 1899)

IN the preface to this book, contributed by M. P. P. Dehérain, the immense importance of the vine culture to France is pointed out, the wine from the department of Hérault alone having in 1897 a value of 212,000,000 francs. The questions of suitability of soil, of manures, of the remedies against the many diseases of the vine, of fermentation, and preservation of wine all depend largely upon simple chemical and physical considerations; hence arises the necessity for such a work as the present, dealing with the physics and chemistry of vine culture and wine production. Of the eight chapters composing the book, the first two are preliminary, giving a very brief outline of the atomic theory and the measuring instruments used in the laboratory. The third chapter deals with the soil, especial attention being directed to the use of various insecticides, and to the causes of vine disease generally residing in the soil. In the third chapter, on account of the importance of the estimation of calcium carbonate in the soil, numerous calcimeters are described, some of considerable and apparently unnecessary complexity, as, for example, the self-registering calcimeter of Houdaille. The description of the properties of manures is lucid, and their analysis is treated in a simple manner. Chapter vi., dealing with the remedies for vine diseases, is, on account of the evident practical knowledge of the author, the most valuable portion of the book. The number of remedies that have been invoked to combat mildew, black rot, chlorosis, phylloxera, and other vine diseases, is so great as to render their classification and intelligent use difficult. Especial attention is here directed to the use of carbon bisulphide, ferrous sulphate, sulphur, copper sulphate and acetate, and mercury salts, the last-named being emphatically condemned in spite of their undoubted efficacy in combating fungoid diseases. The concluding chapters deal briefly with the fermentation of the grape, analysis of the wine, and the diseases to which it is liable. The book will be of great practical service to vine growers.

*Cours Élémentaire de Zoologie.* Par Remy Perrier. Pp. 734. 697 illustrations. (Paris: Masson et Cie., 1899.)

THIS work contains a great deal in brief that is to be found in its predecessor, the author's "Éléments d'Anatomie Comparée," published in 1893. In some respects it may be said to be a "Grundriss" to that volume, but, in contradistinction to it, the Vertebrata are here treated on a greater equality with the Invertebrata, and the order of presentation is more rational and in accordance with precedent. For example, the Chætopod

Worms are dealt with before the Arthropods, the inversion of this order being a notorious feature of the "Éléments." Chapter i. is devoted to broad principles and definitions, Chapter ii. to the elements of histology, and Chapter iii. to the classification of the metazoa—177 pp. in all. Tables of affinity and structural relationship are here and there given, and the 565 remaining pages of the work are devoted to a systematic consideration of each of the greater groups of animal forms in an ascending order, the Echinoderms, Rotifers, Polyzoa, and Brachiopods being taken after the Cœlenterates and before the Leeches and Worms. Some of the groups receive but scanty treatment, meagre and wholly insufficient, and throughout the work the author has conspicuously neglected the rendering clear the extremes of modification of the great groups, which we consider should be an indispensable feature of an elementary text-book on organic forms that shall do justice to our present knowledge. In dealing with such an assemblage as the Tunicata, where octoradiata, valved, stalked, and many other well-known forms occur, a great opportunity has in this way been lost, and the same may be said of the author's treatment of the Bryozoa.

The illustrations are for the most part good and clear; some of the new ones are admirable, and we congratulate the author upon such as his aortic arch series (p. 602), which are the most accurate and up to date of any text-book set yet published. They are sure to be popular and reproduced *ad nauseam*. But why that old nightmare the Cuvier's "Chimæra" (Fig. 589), a badly drawn Chimæra with a Callorhynchus tail! Surely the time has come when this and other persistent atrocities of our text-books, which have so long offended, should be condemned.

A really sound elementary treatise on zoology has long been a desideratum, and the present work is the outcome of a commendable attempt to supply the need. Though desperately thin in parts it is up to date in its leading themes, well arranged, and written in a good easy style, and it may be safely recommended as trustworthy so far as it goes.

#### LETTERS TO THE EDITOR.

[The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, or to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

#### Thermometric Scales for Meteorological Use.

IN the course of some recent work on the meteorology of Ben Nevis, which involved extensive extracting and computing work, I have again had forcibly impressed on me the great advantage which Fahrenheit's thermometer has over that of Celsius for meteorological use, especially in temperate regions.

In chemistry and physics the range of temperature covered is so great that Celsius' scale, which is now universally used, adequately meets every case. The size of the degree and the change of sign at the melting point of ice do not cause any inconvenience in the laboratory. It is otherwise in the meteorological observatory. There the range of temperature dealt with is very restricted, and the Celsius degree is too large, while the change of sign in the middle of the working part of the scale is simply intolerable. The latter peculiarity is the fruitful introducer of error into both the observations and the reductions, and besides it greatly increases the fatigue of both classes of work.

In view of the agitation to abolish the use of Fahrenheit's scale, and to replace it universally by that of Celsius, it may not be inopportune to direct attention to some of the advantages in securing accuracy and in relieving labour which Fahrenheit's scale offers over that of Celsius when used for meteorological purposes.

In tropical countries it matters little whether one scale or the other is used, except that the size of Fahrenheit's degree is much