imported timber, and the evils, including physical degeneration of the race, and coal fogs in the big cities, which have been shown elsewhere to result from England's neglect of its forests. The reference to De Wet in Prof. Perry's communication is unfortunate: a small quick-moving army would probably have caught him. And surely cheap coal and luxury is not the *summum bonum*. Rather let us have hamlets of strong forest workers than the luxurious town dwellers of to-day with their decayed muscle and cheap mechanical power! Compare a European engine-driver with the runner castes of India and The engine-driver shows us perhaps fine inherited Japan. muscle, but going to decay for want of use ; the Eastern runners show the development of muscle by both use and inheritance. Which would have the best chance of catching De Wet a hundred years hence?

As far back as 1882 the discovery was made by Sir D. Brandis and myself that Eucalypts planted on tropical mountains will produce wood fuel at the rate of 20 tons (dry weight at 60 lbs. per cubic foot) per acre per year in perpetuity. The eucalypt plantation reproduces itself when cut, without further expense, and its dry timber, heavier than coal (which, as met with commercially, weighs 50 lbs. to 52 lbs. the cubic foot) has an equal or a higher thermal power, bulk for bulk, than coal. We obtained this result as the maximum yield of *Eucalyptus globulus* on the Nilgiris, Southern India. No doubt there are other instances where higher yields are produced now, and no doubt also when the coal supply is exhausted, selection and experiment will produce a forest vegetation that will produce more than 20 tons per acre per year. The sugar beet and all the fruits and vegetables of civilisation show how the vegetable kingdom can be moulded to suit man's wants. If a chance tree on a chance mountain in a chance soil can produce the equivalent of 20 tons of coal per acre per year, it seems not unreasonable to suppose that by selection we can produce, say, double this, or 40 tons. To produce this in perpetuity we should probably have to find a tree with the moderate soil requirements of the Conifers. A powerful sun, a heavy rainfall, and a very rapid forced growth would be the essentials of such a production of wood fuel.

Looking at a rainfall map of the world, one sees that these conditions are fulfilled over about 8000 million acres of its surface (which is between one-fourth and one-fifth of the total land surface of 35,200 million acres). I take latitudes below 40° and rainfalls above 40 inches. One-half of this area under forest might thus yield the equivalent of 161,000 million tons of coal yearly. This is more than 288 times the world's present consumption of coal, assuming that coal and eucalypt timber are of approximately equal heating power. On the basis of the actual forest yields of to-day we have half this, or 80,500 million tons. In Germany, one-fourth of the total area is under forest, and this is held on the highest authority to be the suitable proportion for a thickly-peopled civilised country such as Germany. The forest should properly occupy a higher proportion in countries where large areas are pestilential and unsuited for human habitation. Putting this, however, aside, and taking the German standard of one-fourth forest, then on the basis of to-day's maximum yields we should obtain a yearly output of 40,250 million tons. And if to convert the maximum forest yield to an average forest yield we again divide by two, we obtain 20,175 million tons. Lower than this I do not think we can reasonably go for the class of forest under consideration. It is a little more than thirty times the world's present con-sumption of coal. The world's yearly output of coal recently was 663 million tons, says Prof. Perry. Thus we see that the yield of firewood from the world's transient provide for the provide the provide the set of the set to be a set of the transient for the provide the set of the set of

tropical and extra-tropical forests, whenever they are fully stocked and scientifically worked, will yield the equivalent of from thirty times to 122 times the present consumption of coal, or even up to 243 times the present consumption of coal if we succeed by cultivation in doubling present timber yield figures.

It may be objected that my figures are far in excess of those representing the yield of European forests and that they require confirmation. No doubt they are far in excess of European figures; but so also is the intensity of the vegetative process in these latitudes, and so also is the stature of the Sequoias of California, and the Eucalypts of Australia and South Africa above the stature of the biggest spruces and silver-firs of Europe. The Nilgiri figures I have quoted above were formally recorded in two official reports printed and published by the

NO. 1706, VOL. 66]

Madras Government in 1882.¹ They have since been confirmed by the measurements of forest officers who have subsequently had charge of the Nilgiri plantations. Similar figures have been obtained by myself and other forest officers in South Africa. They have been exceeded in several plantations in Matal, while at Johannesburg they have not been confined to Eucalypts, but have been obtained from Acacia decurrens, or

black wattle, as well as from some other trees. Therefore, "when our coal supply is exhausted, when all the races of the world have fought for the waterfalls and places of high tide," there will still remain that which Englishmen of all the civilised races of the world do most neglect-the forest.

D. E. HUTCHINS. Grootvadersbosch, Swellendam, Cape Colony, May 14.

Cold Weather in South Africa.

WE have been getting exceptional weather here of late. General French was actually snowed up at Middelburg. A good general idea of the circumstances will be obtained from the telegrams abridged below from the Diamond Fields Advertiser of June 14.

Middelburg (Cape), June 11.-For the first time for sixteen years the town is covered to a great depth with snow. Kingwilliamstown, June 11.—A fierce thunderstorm occurred last night, accompanied by heavy rain. Port Elizabeth, June 11.— The train service between Graaff-Reinet and Rosmead is to-day stopped temporarily owing to heavy snowstorms-an unusual experience for South Africa. Cradock, June 12 .- An exceptionally heavy fall of snow occurred in the Midlands on Tuesday night and yesterday. Queenstown, June 12 .- The rainfall reported during the first five months of the year is the lowest recorded for the same period for the past thirty years. The drought has, however, been broken. Rain started on June 10, and during the night there was a heavy fall of snow. Kokstad, June 12.—There was a heavy snowstorm last night, accompanied by a heavy gale. The snow is several inches deep in the streets. Bloemfontein, June 12.—The weather is unprecedentedly cold. The hills round Thaba Nchu are covered with snow. Last night snow fell in Bloemfontein.

At Kimberley it has been intensely cold, with a low barometer, wind, rain and sleet, and afterwards heavy frost. With the one exception of July 12, 1886 (when Kimberley is said to have been under snow for the whole day), the maximum shade temperature registered is the lowest on record. For the eight days ending Sunday, June 15, the temperatures have been :-

	Observatory Screen.		Stevenson Screen.	
	Max.	Min.	Max.	Min.
June 8	72° 0	34°°0	73°.7	33°.0
», 9 …	59 0	39 O	59 8	38.1
,, 10	40 2	30 0	45,9	37 0
,, II	45 4	30-9	44 9	30 1
,, 12	48 3	31 0	40, 8	31.2
,, 13	52 1	25 2	53 [°] .7	24°'2
,, 14	57° '0	26° .9	58° 0	26° • 0
,, 15	62° 0	29° 0	63°.2	28° '0

The maximum temperatures registered on June 10 and 11 are the lowest on record for any June. The maximum registered on July 12, 1886, was 35° 8. There was also a maximum temperature of 45° in July 1891. Both, however, were obtained under a Glaisher screen and are probably a little too low. Minimum temperatures lower than 25° have been registered perhaps three times; the lowest known is probably 20° in July 1888. All these previous instances have been quite transitory, the temperatures in each case being much higher both the day before and the day after. There seems to be no record of a cold spell having the duration of the one in question. J. R. SUTTON. Kenilworth, Kimberley, June 16.

¹ "Suggestions regarding Forest Administration in the Madras Presi-dency," by D. Brandis, C.I.E., Inspector-General to the Government of India (Madras, 1882). "Report on Measurements of the Growth of Australian Trees on the Nilgiris," by D. E. Hutchins, Dep. Cons. Forests, Mysore (Government Press, Madras, 1883).