part of the annual amount is sometimes made up in the course of a few days. At Okombahe, for instance, two-thirds of the annual amount of $11\frac{1}{2}$ inches fell in the course of four days (January 27-30), and of this amount 5 inches fell on January 28. Generally speaking, little or no rain falls between May and September, inclusive.

Report of the Liverpool Observatory, 1905.—This observatory, maintained at Bidston by the Mersey Docks and Harbour Board, is one of the oldest and best equipped in the United Kingdom, and it transmits daily telegraphic reports to the Meteorological Office. Under the head of automatic instruments are included anemometers of the forms designed by Dines, Osler, and Robinson. We give the comparative maximum records of these during two of the heaviest gales of the year:—

	Dines, miles	Osler, lb. on sq. ft.	Robinson, miles per hour	Direction
1905, Jan. 8 ...	73.3 ...	32 ...	55 ...	S.W.
„ Nov. 26 ...	70.1 ...	40.5 ...	08 ...	W.
„ „ 27 ...	77.0 ...	33.2 ...	61 ...	W.

The gusts recorded by the Dines and Osler anemometers were not exactly at the same time; the figures seem to show that the force of the gusts differs considerably at different points of the same locality. With respect to observations with Milne's seismograph, Mr. Plummer makes the interesting remark that during the time of the recent Antarctic expedition many earthquakes recorded by the exploring party were registered on the instrument at Bidston, although intermediate stations did not in all instances record the tremor. The average number of astronomical observations made with the transit instrument has been practically maintained during the year.

South African Meteorology.—Under the title "South Africa as seen by a Meteorologist," Dr. H. R. Mill gave a lecture before the Royal Meteorological Society on March 21, and an abstract has now been published. The address contains much instructive matter, but Dr. Mill's meteorological notes naturally refer chiefly to rainfall. Table Bay was reached on August 15, 1905, the minimum temperature at Cape Town being only 38° . The most unusual part of the meteorological equipment at the Royal Observatory was the size of the standard rain-gauge, having a diameter of about $11\frac{1}{4}$ inches; at other stations in the colony the size is 8 inches. The usual exposure of the gauges in South Africa is 4 feet high, a fact, as Dr. Mill observes, that must be borne in mind when comparing readings with gauges in this country, where they are usually placed at a height of 1 foot. Meteorology in Cape Colony suffers, the author states, by the excessively small annual grant available, and the opinion is expressed that the good work done at Kenilworth (Kimberley) makes it desirable that the institution should be placed on a permanent footing. The Transvaal Government spends a comparatively large sum on meteorological observations; at Johannesburg the observatory is admirably fitted up, and the site offers peculiar advantages for anemometer work. The rain gauges here and in the Orange River Colony are 5 inches in diameter. The site of the observatory at Grey Town is not a very good one, and is shortly to be changed. Meteorology at Bulawayo is under the charge of Father Goetz, to whose work we recently referred; he has constructed an ingenious electrical recording rain gauge which is apparently very efficient. On the homeward journey a visit was made to the Portuguese station at Beira; with regard to this observatory, Dr. Mill remarks that it is "an imposing structure and the rain gauge is of heroic dimensions." This interesting paper is embellished by many photographic illustrations.

Report of the Observatory Department of the National Physical Laboratory for the Year 1905.—As this branch is in many respects complete in itself, and its work appeals to a different class from that interested in the other departments, the director thinks it desirable to issue the report separately. The magnetographs have been in constant operation throughout the year, and the curves have again been free from any very large disturbances; the most interesting movements were those of November 12 and 15. On the latter date an auroral display was generally observed (NATURE, November 30, 1905, p. 101). The mean declination during the year was $16^{\circ} 32' 9''$ W., mean

inclination $67^{\circ} 3' 8''$ N. Owing to the disturbance of the vertical force produced by electric trams, it was found impossible to tabulate the curves for this element satisfactorily. The meteorological traces and tabulations have been, as usual, sent to the Meteorological Office for publication, therefore only the results are given as an appendix to the report in question. The maximum shade temperature was $81^{\circ} 5'$, on July 8, and the minimum $22^{\circ} 4'$, on November 22 ($23^{\circ} 4'$ on January 19). The rainfall was 22.61 inches, and the number of rain days 154, including five days on which snow was recorded. The number of instruments verified (exclusive of watches and chronometers) amounted to 26,658, being a considerable increase as compared with the previous year; about 60 per cent. of these instruments were clinical thermometers.

The Warm Air Current at the Height of 10-12 Kilometres.—In the *Meteorologische Zeitschrift* for June, Dr. R. Nimführ discusses the question of one of the most interesting results of the international balloon ascents, viz. the "inversion of temperature" at an altitude of 8-13 km. pointed out by M. L. Teisserenc de Bort (*Comptes rendus*, April 28, 1902), and by Dr. Assmann at an altitude of 10-15 km. (*Sitzb. Akad.*, Berlin, May 1, 1902). Dr. Nimführ states that the French experiments were made with paper balloons, that in about half the ascents the maximum height of the balloon was at the critical altitude of 11-12 km., and that consequently the instruments were affected by solar radiation owing to decrease of ventilation; also that Dr. Assmann's experiments were probably similarly affected, although to a less extent, as he used closed rubber balloons. Dr. Nimführ thinks that the lifting power of the balloons was decreased in the higher regions owing to the rubber becoming porous by expansion; further, that the bimetallic thermometer used in some ascents is subject to a fundamental error, now under further investigation, which possibly affects some of the results obtained. We offer no comments on the questions raised; they will no doubt receive full consideration by those engaged in this important branch of meteorological inquiry.

Rainfall in the Philippines.—The Bulletin for December last, issued by the Weather Bureau of Manila, under the direction of the Rev. Father Algué, contains a table of the monthly and annual distribution of rainfall in 1905 at fifty-three stations scattered over the different islands of the archipelago. It is shown that the stations may be arranged in three groups:—(1) where the fall is uniform in the various months; (2) where the rainfall is scarce from December to March inclusive; (3) stations with abundant rains from June to October, and little in the rest of the year. At some stations the amounts are large, e.g. Baguio, an elevated plateau, more than 165 inches, while eighteen of the other stations have falls varying from 79 inches to 115 inches. From the results of the last five years' observations at Baguio we note that the mean annual temperature there is $65^{\circ} 3'$; the lowest monthly mean is $61^{\circ} 3'$, in February, and the highest $69^{\circ} 1'$, in April and May. The absolute maximum was $84^{\circ} 7'$, in April, and the minimum $42^{\circ} 8'$, in February. Rain falls on an average on 171 days, mostly between May and October.

DISEASES OF SHEEP.

PROBABLY few persons, unless they have had reason to study the matter, have any idea of the immense economic importance of the diseases of animals. As a matter of fact, our flocks and herds are every year stricken down to an extent representing a value of hundreds of thousands of pounds!

In 1901 a committee, consisting of Prof. Hamilton, Mr. J. McCall, and Mr. E. G. Wheeler, with Mr. R. B. Greig as secretary, was appointed by the Board of Agriculture to investigate and report on the diseases of sheep known as louping-ill and braxy, and the findings of this committee have lately been published in a voluminous and interesting report.

Louping-ill is a disease which shows itself in the form of nervous spasms of the limbs and neck, or rigidity, followed by more or less complete paralysis; sometimes, however, there is a general dazed condition with speedy

collapse. It does not seem to be known in foreign countries, but in the British Isles is met with particularly on the west coasts of Ireland and Scotland, in Cumberland and Westmorland, and in small, scattered areas throughout the country. Sheep of all ages may be attacked, and the mortality may amount to 20 per cent. or even more.

Braxy is often a rapidly fatal disease. The animal goes off its feed, is restless, the belly swells, it falls on its side, becomes semi-comatose, and death soon ensues, the carcase having a characteristic odour. The disease prevails in several countries of northern Europe, and in the British Isles on the west coasts of Ireland and Scotland, central Wales, Westmorland and Northumberland, Cornwall, Wilts, and Gloucester. Sheep under one year are the chief sufferers.

A remarkable feature of both these (and certain other) diseases of the sheep is their seasonal prevalence; thus louping-ill and braxy are not met with during July and August, and the former is most prevalent from April to June, the latter from November to February.

In the case of louping-ill, for a long time the specific cause remained a mystery, carcase after carcase examined

ing the animals with cultures during the insusceptible period was adopted, and proved a decided success on the large scale. Thus, with louping-ill, 1340 sheep were treated in this manner, and a single doubtful death from the disease occurred; with braxy, 1545 sheep were treated, and there were nine possible (three being doubtful) deaths from braxy among them.

A remarkable discovery was made with regard to the seasonal susceptibility and immunity. It was found that during the period of immunity the blood of the sheep proved highly bactericidal towards the louping-ill and braxy bacilli, while during the susceptible period the bacilli were not only not destroyed by, but grew well in, the sheep's blood.

As already indicated, the diseases are mostly communicated by the fouling of the pastures by the dejecta. It has been held by some that the sheep-tick plays a part in their transmission, but experiments showed that this could only be to a very insignificant extent.

The report, which is illustrated with a number of figures and maps, is highly suggestive in many directions; the researches made promise to throw new light on the path-

FIG. 1.—Bacillus of Louping-ill in peritoneal liquid of sheep, showing the rods, some without spores, others with spores in their interior. $\times 1000$ diameters.

showing no lesions, and inoculations of the blood, &c., failed to convey the disease from one sheep to another. At last, examination of the fluid in the peritoneal cavity revealed the presence of a large sporing anærobic bacillus (Fig. 1), which, on inoculation into healthy sheep, conveyed the disease again and again. It was for a long time an enigma how this bacillus reached the peritoneal cavity, the blood and tissues being free from it. Eventually, however, in a diseased lamb the intestine was found to be swarming with the bacillus, and a long series of experiments proved that the organisms or their spores are taken in with the food, and if at the susceptible period of the year induce the disease in a large proportion of cases. The organism, being passed with the dejecta, fouls the pasture, and so the disease is propagated. Precisely the same holds good for braxy, which, however, is caused by an organism different from the louping-ill bacillus, the braxy organism being also an anærobic sporing bacillus, but being much smaller and more delicate than the louping-ill bacillus (Fig. 2).

Attempts to immunise by means of injections of attenuated organisms or by chemical products of the organisms proved not only failures, but dangerous on account of the mortality. Taking into account the fact that the organisms are intestinal, the happy idea of drench-

FIG. 2.—Bacillus of Braxy, peritoneal liquid of sheep, showing the comparatively delicate rods, some sporing, others not; those sporing have an oval or lanceolate form. $\times 1000$ diameters.

ology of many of the contagious and infectious maladies of man and the lower animals, and we congratulate Prof. Hamilton and his collaborators on the valuable work they have done.

R. T. HEWLETT.

THE NEW MUSPRATT LABORATORY OF PHYSICAL AND ELECTROCHEMISTRY AT THE UNIVERSITY OF LIVERPOOL.

THE laboratory of physical and electrochemistry, which the University of Liverpool owes to the munificent generosity of Mr. E. K. Muspratt, president of the council of the University, was formally opened by Sir William Ramsay, K.C.B., F.R.S., on Saturday, October 13. The distinguished company which assembled in Liverpool for the occasion included, amongst others, the following well-known men of science from abroad:—Prof. Ostwald (Leipzig), Prof. R. Abegg (Breslau), Prof. Ernst Cohen (Utrecht), Prof. H. Goldschmidt (Christiania), Prof. Lash Miller (Toronto), and Prof. Macallum (Toronto).

On Saturday forenoon the guests inspected the new laboratory privately, and were afterwards entertained to lunch by Sir John Brunner, Bart., M.P., at the University Club, many other prominent men of science and letters in