had to devote himself to military duties, Dr. Arthur Smith was appointed acting-professor of anatomy in the University of Sydney, and Colonel Wilson most magnanimously handed over to him this most important human relic for investigation and description. With the valuable assistance of Mr. D. M. S. Watson he has succeeded in removing the matrix and fully exposing the extraordinary features of Australia's earliest known human inhabitant, perhaps a member of the earliest family to ferry across Wallace's line and tramp dogs through Papua to Queensland.

The skull has been purchased by the Hon. Joynton Smith, M.L.C., and presented by him to the University of Sydney. The formal presentation was made the occasion of a public meeting in the University on June 18, presided over by the chancellor, when Acting-Prof. Arthur Smith and Prof. David delivered addresses on the significance of the skull and its bearing upon the problems of man's antiquity in Australia.

Prof. David explained that the skull (of the Pleistocene period) was found after a flood thirty-one years ago in the bed of a creek near Talgai station, on the Darling Downs, by a stockman. From its highly mineralised state and mode of occurrence in a region where other fossil remains had also been found, the skull was of sufficiently striking geological antiquity to be grouped with the last ice epoch of the northern hemisphere. At the close of the last glacial period, gigantic animals became extinct, and probably the big animals known to have existed in Australia ages ago likewise died at the same time as those of the northern hemisphere. The discovery of the skull might serve to explain how it was that Australia, with its marsupial fauna, had an invader in the form of the dingo, which was a sort of Asiatic jackal, and "had no more right here than the Germans had in Belgium." There was the possibility of its having been brought here by early man. However, the skull was worth its weight in gold. The University of Sydney and men of science throughout the world were under a deep debt of gratitude to Mr. Smith for purchasing the skull, which was undoubtedly a priceless specimen.

Acting-Prof. Smith stated that the condition of the teeth was such as to show that the skull was a youth's, while at the same time it had many of the characteristics of an adult skull. The canine tooth of the Talgai fossil was the largest human tooth so far discovered. The extremely primitive characteristics were such as to warrant one claiming for the skull considerable importance in anthropology, and worthy of being placed with such specimens as the prehistoric Heidelberg jaw and the Piltdown skull. There was, he concluded, certain to be a great controversy concerning the latest discovery when the results of the investigation were made known.

THE RAINFALL OF JAVA.

LYING as it does midway between India and Australia, Java forms a climatological link between the two great British Dominions, and we recognise the appropriateness of the compliment implied in the choice of English as the second language in which this important outcome of Dutch enterprise is published. Tables are given of the mean monthly rainfall, number of rain days, and mean daily maximum rainfall (for each month) at 1061 stations in Java from observations taken between

1879 and 1911. Dr. van Bemmelen discusses the trustworthiness of the data, which are far from uniform. Obviously fallacious records have been eliminated, and there is reason to believe that the data here given compare well with those available for other tropical countries. Observations were made to the nearest millimetre, so that the critical rainfall which determines whether or not a day is to be counted as a rain day is 0.5 mm., or 0.02 in., i.e. four times the amount taken as determining a rain day in this country. As an example of the variability of monthly rainfall in Java, the case is cited of one station where in September, 1902, the rainfall was 12 mm., and in September, 1904, 896 mm. The greatest average annual rainfall recorded at any station in Java is 8305 mm., and the least 882 mm., but for both stations quoted the average was only for the short period of seven years and experience shows that runs of seven years may occur with every year above or every year below the long-period average. The wettest year ever recorded at any station was 10,112 mm. at Sirah Kentjong, Kediri, in 1909, and the greatest fall in twenty-four hours was 511 mm. at Besokor, Semarang, on January 31, 1901. Perhaps I may be pardoned if I remind those who keep metric and vulgar statistics in separate compartments of their brains that these figures are equivalent to 398.1 and 20.1 in. respectively.

The most important part of Dr. van Bemmelen's work is his series of maps of average rainfall. That he was fully alive to the difficulties in his way is made plain by his quotation of the reasons which deterred him in 1908 from making the attempt. To our mind the greatest difficulty, and one that has not been overcome, is the use of averages of varying lengths without reduction to a common period. We agree with the author that such reduction was difficult, if not impossible, in the circumstances, and that any sort of isohyetal map is better than none, and for many purposes, both scientific and economic, the maps he has produced will prove most useful. He gives a large-scale (1 : 1,000,000) map showing the position of the stations, and a series of smaller maps (scale 1 : 1,500,000) for annual rainfall, and the mean monthly rainfall of four selected seasons. These are:—(1) the mean of the three wettest months, December, January, February, when the west monsoon is blowing; (2) the mean for the two months March and April, showing the main transition; and (3) the mean for the two months May and June, when the transition is almost complete, to (4) the period of the east monsoon, represented by the mean of the three months July, August, September—the driest part of the year; finally, (5) the mean of October and November, showing the transition to the wet season again. These maps are admirably executed and most suggestive; but we are of opinion that both the climatic transitions and the geographical distribution would be more clearly suggested by maps based on the homogeneous average of, say, the last ten years, than on the data actually used. We are in hearty agreement with Dr. van Bemmelen's method of drawing isohyetal lines from such data as are available by what he aptly terms "a continual compromise between deduction and induction," the deduction being based on the recognised relationship between configuration, wind-direction, and rainfall, the induction being from the data which have been actually observed. Even where the data are as numerous, long-continued, and reasonably accurate as they are for the British Isles, the most probable distribution of rainfall can only be represented cartographically by such a "continual compromise."

HUGH ROBERT MILL.

¹ "Uitkomsten der Regenwaarnemingen op Java." (Results of Rainfall Observations in Java.) By Dr. W. van Bemmelen. Pp. xxiii+173. Regenatlas (Rainfall-Atlas), vii maps. (Batavia: Javasche Boekhandel Drukkerij, 1914 and 1915.)