Oceanography and the Sea-Fisheries.*

By WILLIAM A. HERDMAN, C.B.E., D.Sc., Sc.D., LL.D., F.R.S.,

Professor of Oceanography in the University of Liverpool, President.

IT has been customary, when occasion required, for the president to offer a brief tribute to the memory of distinguished members of the Association lost to science during the preceding year. These, for the most part, have been men of advanced years and high reputation who had completed their life-work and served well in their day the Association and the sciences which it represents. Such are our late general treasurer, Prof. Perry, and our past-president, Sir Norman Lockyer, of whom the retiring president has just spoken. We have this year no other such losses to record; but it seems fitting on the present occasion to pause for a moment and devote a grateful thought to that glorious band of fine young men of high promise in science who, in the years since our Australian meeting in 1914, gave, it may be, in brief days and months of sacrifice, greater service to humanity and the advance of civilisation than would have been possible in years of normal time and work. A few names stand out already known and highly honoured—Moseley, Jenkinson, Geoffrey Smith, Keith Lucas, Gregory, and more recently Leonard Doncaster—all grievous losses; but there are also others, younger members of our Association, who had not yet had opportunity for showing accomplished work, but who equally gave up all for a great ideal. I prefer to offer a collective rather than an individual tribute. Other young men of science will arise and carry on their work, but the gap in our ranks remains. Let their successors remember that it serves as a reminder of a great example and of high endeavour worthy of our gratitude and of permanent record in the annals of science.

At the last Cardiff meeting of the British Association in 1891 you had as your president the eminent astronomer Sir William Huggins, who discoursed upon the then recent discoveries of the spectroscope in relation to the chemical nature, density, temperature, pressure, and even the motions of the stars. From the sky to the sea is a long drop, but the sciences of both have this in common: that they deal with fundamental principles and with vast numbers. More than three hundred years ago Spenser in the "Faerie Queene" compared "the seas abundant progeny" with "the starres' on hy," and recent investigations show that a litre of sea-water may contain more than a hundred times as many living organisms as there are stars visible to the eye on a clear night.

During the past quarter of a century great advances have been made in the science of the sea, and the aspects and prospects of sea-fisheries research have undergone changes which encourage the hope that a combination of the work now carried on by hydrographers and biologists in most civilised countries on fundamental problems of the ocean may result in a more rational exploitation and administration of the fishing industries.

And yet even at your former Cardiff meeting thirty years ago there were at least three papers of oceanographic interest—one by Prof. Osborne Reynolds on the action of waves and currents, another by Dr. H. R. Mill on seasonal variation in the temperature of lochs and estuaries, and the third by our honorary local secretary for the present meeting, Dr. Evans Hovle, on a deep-sea tow-net capable of being opened and closed under water by the electric current.

It was a notable meeting in several other respects,

* Presidential address delivered at the Cardiff meeting of the British Association on August 24.

of which I shall merely mention two. In Section A Sir Oliver Lodge gave the historic address in which he expounded the urgent need, in the interests of both science and the industries, of a national institution for the promotion of physical research on a large scale. Lodge's pregnant idea put forward at this Cardiff meeting, supported and still further elaborated by Sir Douglas Galton as president of the Association at Ipswich, has since borne notable fruit in the establishment and rapid development of the National Physical Laboratory. The other outstanding event of that meeting is that you then appointed a committee of eminent geologists and naturalists to consider a project for boring through a coral reef, and that led during following years to the successive expeditions to the atoll of Funafuti, in the Central Pacific, the results of which, reported upon eventually by the Royal Society, were of great interest alike to geologists, biologists, and oceanographers.

Dr. Huggins, on taking the chair in 1891, remarked that it was more than thirty years since the Association had honoured astronomy in the selection of its president. It might be said that the case of oceanography is harder, as the Association has never had an oceanographer as president; and the Association might well reply, "Because until very recent years there has been no oceanographer to have." If astronomy is the oldest of the sciences, oceanography is probably the youngest. Depending as it does upon the methods and results of other sciences, it was not until our knowledge of physics, chemistry, and biology was relatively far advanced that it became possible to apply that knowledge to the investigation and explanation of the phenomena of the ocean. No one man has done more to apply such knowledge derived from various other subjects and to organise the results as a definite branch of science than the late Sir John Murray, who may therefore be regarded as the founder of modern oceanography.

It is to me a matter of regret that Sir John Murray was never president of the British Association. I am revealing no secret when I tell you that he might have been. On more than one occasion he was invited by the council to accept nomination, and he declined for reasons that were good and commanded our respect. He felt that the necessary duties of this post would interfere with what he regarded as his primary life-work—oceanographical explorations already planned, and the last of which he actually carried out in the North Atlantic in 1912, when above seventy years of age, in the Norwegian steamer *Michael Sars* along with his friend Dr. Johan Hjort.

Anyone considering the subject-matter of this new science must be struck by its wide range, overlapping as it does the borderlands of several other sciences and making use of their methods and facts in the solution of its problems. It is not only world-wide in its scope, but it also extends beyond our globe, and includes astronomical data in their relation to tidal and certain other oceanographical phenomena. No man in his work, or even thought, can attempt to cover the whole ground, although Sir John Murray, in his remarkably comprehensive "Summary" volumes of the *Challenger* Expedition and other writings, went far towards doing so. He, in his combination of physicist, chemist, geologist, and biologist, was the nearest approach we have had to an all-round oceanographer. The International Research Council probably acted wisely at the recent Brussels Confer-

ence in recommending the institution of two International Sections in our subject, one of physical and the other of biological oceanography, although the two overlap and are so interdependent that no investigator on one side can afford to neglect the other.¹

On the present occasion I must restrict myself almost wholly to the latter division of the subject, and be content, after brief reference to the founders and pioneers of our science, to outline a few of those investigations and problems which have appeared to me to be of fundamental importance, of economic value, or of general interest.

Although the name "oceanography" was only given to this branch of science by Sir John Murray in 1880, and although, according to that veteran oceanographer Mr. J. Y. Buchanan, the last surviving member of the civilian staff of the *Challenger*, the science of oceanography was born at sea on February 15, 1873,² when at the first official dredging station of the expedition, to the westward of Teneriffe, at 1525 fathoms, everything that came up in the dredge was new, and led to fundamental discoveries as to the deposits forming on the floor of the ocean, still it may be claimed that the foundations of the science were laid by various explorers of the ocean at much earlier dates. Aristotle, who took all knowledge for his province, was an early oceanographer on the shores of Asia Minor. When Pytheas passed between the Pillars of Hercules into the unknown Atlantic and penetrated to British seas in the fourth century B.C., and brought back reports of Ultima Thule and of a sea to the north thick and sluggish like a jellyfish, he may have been recording an early planktonic observation. But passing over all such and many other early records of phenomena of the sea, we come to surer ground in claiming as founders of oceanography Count Marsili, an early investigator of the Mediterranean, and that truly scientific navigator Capt. James Cook, who sailed to the South Pacific on a Transit of Venus expedition in 1769, with Sir Joseph Banks as naturalist, and by afterwards circumnavigating the South Sea about latitude 60° finally disproved the existence of a great southern continent; and Sir James Clark Ross, who, with Sir Joseph Hooker as naturalist, first dredged the Antarctic in 1840.

The use of the naturalist's dredge (introduced by O. F. Müller, the Dane, in 1799) for exploring the sea-bottom was brought into prominence almost simultaneously in several countries of North-West Europe—by Henri Milne-Edwards in France in 1830, by Michael Sars in Norway in 1835, and by our own Edward Forbes about 1832.

The last-mentioned genial and many-sided genius was a notable figure in several sections of the British Association from about 1836 onwards, and may fairly be claimed as a pioneer of oceanography. In 1839 he and his friend the anatomist, John Goodsir, were dredging in the Shetland seas, with results which Forbes made known to the meeting of the British Association at Birmingham that summer, with such

¹ The following classification of the primary divisions of the subject may possibly be found acceptable:—

² Others might put the date later. Significant publications are Sir John Murray's Summary Volumes of the *Challenger* (1885), the inauguration of the Musée Océanographique at Monaco in 1910, the foundation of the Institut Océanographique at Paris in 1906 (see the Prince of Monaco's letter to the Minister of Public Instruction), and Sir John Murray's little book "The Ocean" (1913), where the superiority of the term "oceanography" to "thalassography" (used by Alexander Agassiz) is discussed.

good effect that a "Dredging Committee"³ of the Association was formed to continue the good work. Valuable reports on the discoveries of that committee appear in our volumes at intervals during the following twenty-five years.

It has happened over and over again in history that the British Association, by means of one of its research committees, has led the way in some important new research or development of science, and shown the Government or an industry what wants doing and how it can be done. We may fairly claim that the British Association has inspired and fostered that exploration of British seas which through marine biological investigations and deep-sea expeditions has led on to modern oceanography. Edward Forbes and the British Association Dredging Committee, Wyville Thomson, Carpenter, Gwyn Jeffreys, Norman, and other naturalists of the pre-*Challenger* days—all these men in the quarter-century from 1840 onwards worked under research committees of the British Association, bringing their results before successive meetings; and some of our older volumes enshrine classic reports on dredging by Forbes, McAndrew, Norman, Brady, Alder, and other notable naturalists of that day. These local researches paved the way for the *Challenger* and other national deep-sea expeditions. Here, as in other cases, it required private enterprise to precede and stimulate Government action.

It is probable that Forbes and his fellow-workers on this "Dredging Committee" in their marine explorations did not fully realise that they were opening up a most comprehensive and important department of knowledge. But it is also true that in all his expeditions—in the British seas from the Channel Islands to the Shetlands, in Norway, and in the Mediterranean as far as the Aegean Sea—his broad outlook on the problems of Nature was that of the modern oceanographer, and he was the spiritual ancestor of men like Sir Wyville Thomson, of the *Challenger* Expedition, and Sir John Murray, whose accidental death a few years ago, whilst still in the midst of active work, was a grievous loss to this new and rapidly advancing science of the sea.

Forbes in these marine investigations worked at border-line problems, dealing, for example, with the relations of geology to zoology and the effect of the past history of the land and sea upon the distribution of plants and animals at the present day, and in these respects he was an early oceanographer. For the essence of that new subject is that it also investigates border-line problems, and is based upon and makes use of all the older fundamental sciences—physics, chemistry, and biology—and shows, for example, how variations in the great ocean-currents may account for the movements and abundance of the migratory fishes, and how periodic changes in the physico-chemical characters of the sea, such as variations in the hydrogen-ion and hydroxyl-ion concentration, are correlated with the distribution at the different seasons of the all-important microscopic organisms that render our oceanic waters as prolific a source of food as the pastures of the land.

Another pioneer of the nineteenth century who, I sometimes think, has not yet received sufficient credit for his foresight and initiative is Sir Wyville Thomson, whose name ought to go down through the ages as the leader of the scientific staff on the famous *Challenger* Deep-Sea Exploring Expedition. It is due chiefly to him and to his friend, Dr. W. B.

³ "For researches with the dredge, with a view to the investigation of the marine zoology of Great Britain, the illustration of the geographical distribution of marine animals, and the more accurate determination of the fossils of the Pleistocene period: under the superintendence of Mr. Gray, Mr. Forbes, Mr. Goodsir, Mr. Patterson, Mr. Thompson of Belfast, Mr. Ball of Dublin, Dr. George Johnston, Mr. Smith of Jordan Hill, and Mr. A. Strickland, 60/." Report for 1839, p. xxvi.

Carpenter, that the British Government, through the influence of the Royal Society, was induced to place at the disposal of a committee of scientific experts, first the small surveying steamer *Lightning* in 1868, and then the more efficient steamer *Porcupine* in the two succeeding years, for the purpose of exploring the deep water of the Atlantic from the Faroes in the north to Gibraltar and beyond in the south, in the course of which expeditions they got successful hauls from the then unprecedented depth of 2435 fathoms, nearly three statute miles.

It will be remembered that Edward Forbes, from his observations in the Mediterranean (an abnormal sea in some respects), regarded depths of more than 300 fathoms as an azoic zone. It was the work of Wyville Thomson and his colleagues, Carpenter and Gwyn Jeffreys, on these successive dredging expeditions to prove conclusively what was beginning to be suspected by naturalists, that there is no azoic zone in the sea, but that abundant life belonging to many groups of animals extends down to the greatest depths of from four to five thousand fathoms—nearly six statute miles from the surface.

These pioneering expeditions in the *Lightning* and *Porcupine*—the results of which are not even yet fully made known to science—were epoch-making, inasmuch as they not only opened up this new region to the systematic marine biologist, but also gave glimpses of world-wide problems in connection with the physics, the chemistry, and the biology of the sea which are only now being adequately investigated by the modern oceanographer. These results, which aroused intense interest amongst the leading scientific men of the time, were so rapidly surpassed and overshadowed by the still greater achievements of the *Challenger* and other national exploring expeditions that followed in the seventies and eighties of last century, that there is some danger of their real importance being lost sight of; but it ought never to be forgotten that they first demonstrated the abundance of life of a varied nature in depths formerly supposed to be azoic, and, moreover, that some of the new deep-sea animals obtained were related to extinct forms belonging to the Jurassic, Cretaceous, and Tertiary periods.

It is interesting to recall that our Association played its part in promoting the movement that led to the *Challenger* Expedition. Our general committee at the Edinburgh meeting of 1871 recommended that the president and council be authorised to co-operate with the Royal Society in promoting "a circumnavigation expedition, specially fitted out to carry the physical and biological exploration of the deep sea into all the great oceanic areas"; and our council later appointed a committee consisting of Dr. Carpenter, Prof. Huxley, and others to co-operate with the Royal Society in carrying out these objects.

It has been said that the *Challenger* Expedition will rank in history with the voyages of Vasco da Gama, Columbus, Magellan, and Cook. Like these, it added new regions of the globe to our knowledge, and the wide expanses thus opened up for the first time, the floors of the oceans, though less accessible, are vaster than the discoveries of any previous exploration. Has not the time come for a new *Challenger* expedition?

Sir Wyville Thomson, although leader of the expedition, did not live to see the completed results, and Sir John Murray will be remembered in the history of science as the *Challenger* naturalist who brought to a successful issue the investigation of the enormous collections and the publication of the scientific results of that memorable voyage; these two Scots share the honour of having guided the destinies of what is still the greatest oceanographic exploration of all time.

In addition to taking his part in the general work of the expedition, Murray devoted special attention to three subjects of primary importance in the science of the sea, viz.: (1) The plankton or floating life of the oceans, (2) the deposits forming on the sea-bottoms, and (3) the origin and mode of formation of coral-reefs and islands. It was characteristic of his broad and synthetic outlook on Nature that, in place of working at the specicography and anatomy of some group of organisms, however novel, interesting, and attractive to the naturalist the deep-sea organisms might seem to be, he took up wide-reaching general problems with economic and geological as well as biological applications.

Each of the three main lines of investigation—deposits, plankton, and coral-reefs—which Murray undertook on board the *Challenger* has been most fruitful of results both in his own hands and in those of others. His plankton work has led on to those modern planktonic researches which are closely bound up with the scientific investigation of our sea-fisheries.

His work on the deposits accumulating on the floor of the ocean resulted, after years of study in the laboratory as well as in the field, in collaboration with the Abbé Renard, of the Brussels Museum, afterwards professor at Ghent, in the production of the monumental "Deep-Sea Deposits" volume, one of the *Challenger* reports, which first revealed to the scientific world the detailed nature and distribution of the varied submarine deposits of the globe and their relation to the rocks forming the crust of the earth.

These studies led, moreover, to one of the romances of science which deeply influenced Murray's future life and work. In accumulating material from all parts of the world and all deep-sea exploring expeditions for comparison with the *Challenger* series, some ten years later, Murray found that a sample of rock from Christmas Island, in the Indian Ocean, which had been sent to him by Comdr. (now Admiral) Aldrich, of H.M.S. *Egeria*, was composed of a valuable phosphatic material. This discovery in Murray's hands gave rise to a profitable commercial undertaking, and he was able to show that some years ago the British Treasury had already received in royalties and taxes from the island considerably more than the total cost of the *Challenger* Expedition.

That first British circumnavigating expedition on the *Challenger* was followed by other national expeditions (the American *Tuscarora* and *Albatross*, the French *Travailleur*, the German *Gauss*, *National*, and *Valdivia*, the Italian *Vettor Pisani*, the Dutch *Siboga*, the Danish *Thor*, and others) and by almost equally celebrated and important work by unofficial oceanographers such as Alexander Agassiz, Sir John Murray with Dr. Hjort in the *Michael Sars*, and the Prince of Monaco in his magnificent ocean-going yacht, and by much other good work by many investigators in smaller and humbler vessels. One of these supplementary expeditions I must refer to briefly because of its connection with sea-fisheries. The *Triton*, under Tizard and Murray in 1882, while exploring the cold and warm areas of the Faroe Channel separated by the Wyville Thomson ridge, incidentally discovered the famous Dubh-Artach fishing-grounds, which have been worked by British trawlers ever since.

Notwithstanding all this activity during the last forty years since oceanography became a science, much has still to be investigated in all seas in all branches of the subject. On pursuing any line of investigation one very soon comes up against a wall of the unknown or a maze of controversy. Peculiar difficulties surround the subject. The matters investigated are often remote and almost inaccessible. Un-

known factors may enter into every problem. The samples required may be at the other end of a rope or a wire eight to ten miles long, and the oceanographer may have to grope for them literally in the dark and under other difficult conditions which make it uncertain whether his samples when obtained are adequate and representative, and whether they have undergone any change since leaving their natural environment. It is not surprising, then, that in the progress of knowledge mistakes have been made and corrected, and that views have been held on what seemed good scientific grounds which later on were proved to be erroneous. For example, Edward Forbes, in his division of life in the sea into zones, on what then seemed to be sufficiently good observations in the Ægean, but which we now know to be exceptional, placed the limit of life at 300 fathoms, while Wyville Thomson and his fellow-workers on the *Porcupine* and the *Challenger* showed that there is no azoic zone even in the great abysses.

Or, again, take the celebrated myth of Bathybius. In the sixties of last century samples of Atlantic mud, taken when surveying the bottom for the first telegraph cables and preserved in alcohol, were found when examined by Huxley, Haeckel, and others to contain what seemed to be an exceedingly primitive protoplasmic organism, which was supposed on good evidence to be extended widely over the floor of the ocean. The discovery of this Bathybius was said to solve the problem of how the deep-sea animals were nourished in the absence of seaweeds. Here was a widespread protoplasmic meadow upon which other organisms could graze. Belief in Bathybius seemed to be confirmed and established by Wyville Thomson's results in the *Porcupine* Expedition of 1869, but was exploded by the naturalists on the *Challenger* some five years later. Buchanan in his recently published "Accounts Rendered" tells us how he and his colleague Murray were keenly on the lookout for hours at a time on all possible occasions for traces of this organism, and how they finally proved, in the spring of 1875 on the voyage between Hong-Kong and Yokohama, that the all-pervading substance like coagulated mucus was an amorphous precipitate of sulphate of lime thrown down from the sea-water in the mud on the addition of a certain proportion of alcohol. He wrote to this effect from Japan to Prof. Crum Brown, and it is in evidence that after receiving this letter Crum Brown interested his friends in Edinburgh by showing them how to make Bathybius in the chemical laboratory. Huxley at the Sheffield meeting of the British Association in 1879 handsomely admitted that he had been mistaken, and it is said that he characterised Bathybius as "not having fulfilled the promise of its youth." Will any of our present oceanographic beliefs share the fate of Bathybius in the future? Some may, but even if they do they may well have been useful steps in the progress of science. Although, like Bathybius, they may not have fulfilled the promise of their youth, yet we may add they will not have lived in the minds of man in vain.

Many of the phenomena we encounter in oceanographic investigations are so complex, are or may be affected by so many diverse factors, that it is difficult, if indeed possible, to be sure that we are unravelling them aright and see the real causes of what we observe.

Some few things we know approximately, nothing completely. We know that the greatest depths of the ocean, about six miles, are a little greater than the highest mountains on land, and Sir John Murray has calculated that if all the land were washed down into the sea the whole globe would be covered by an ocean

averaging about two miles in depth.⁴ We know the distribution of temperatures and salinities over a great part of the surface and a good deal of the bottom of the oceans, and some of the more important oceanic currents have been charted and their periodic variations, such as those of the Gulf Stream, are being studied. We know a good deal about the organisms floating or swimming in the surface waters (the epiplankton), and also those brought up by our dredges and trawls from the bottom in many parts of the world, although every expedition still makes large additions to knowledge. The region that is least known to us, both in its physical conditions and in its inhabitants, is the vast zone of intermediate waters lying between the upper few hundred fathoms and the bottom. That is the region that Alexander Agassiz, from his observations with closing tow-nets on the *Blake* Expedition, supposed to be destitute of life, or at least, as modified by his later observations on the *Albatross*, to be relatively destitute compared with the surface and the bottom, in opposition to the contention of Murray and other oceanographers that an abundant meso-plankton was present, and that certain groups of animals, such as the Challengerida and some kinds of Medusæ, were characteristic of these deeper zones. I believe that, as sometimes happens in scientific controversies, both sides were right up to a point, and both could support their views upon observations from particular regions of the ocean in certain circumstances.

But much still remains unknown or only imperfectly known even in matters that have long been studied and where practical applications of great value are obtained—such as the investigation and prediction of tidal phenomena. We are now told that theories require reinvestigation, and that published tables are not sufficiently accurate. To take another practical application of oceanographic work, the ultimate causes of variations in the abundance, in the sizes, in the movements, and in the qualities of the fishes of our coastal industries are still to seek, and, notwithstanding volumes of investigation and a still greater volume of discussion, no man who knows anything of the matter is satisfied with our present knowledge of even the best-known and economically most important of our fishes such as the herring, the cod, the plaice, and the salmon.

Take the case of our common fresh-water eel as an example of how little we know and at the same time of how much has been discovered. All the eels of our streams and lakes of North-West Europe live and feed and grow under our eyes without reproducing their kind; no spawning eel has ever been seen. After living for years in immaturity, at last near the end of their lives the large male and female yellow eels undergo a change in appearance and in nature. They acquire a silvery colour and their eyes enlarge, and in this bridal attire they commence the long journey which ends in maturity, reproduction, and death. From all the fresh waters they migrate in the autumn to the coast, from the inshore seas to the open ocean and still westward and south to the mid-Atlantic, and we know not how much further, for the exact locality and manner of spawning have still to be discovered. The youngest known stages of the *Leptocephalus*, the larval stage of eels, have been found by the Dane, Dr. Johannes Schmidt, to the west of the Azores, where the water is more than 2000 fathoms in depth. These were about one-third of an inch in length, and were probably not long hatched. I cannot now refer to all the able inves-

⁴ It was possibly in such a former world-wide ocean of ionised water that, according to the recent speculations of A. H. Church ("Thalassiophyta," 1919), the first living organisms were evolved, to become later the floating unicellular plants of the primitive plankton.

tigators—Grassi, Hjort, and others—who have discovered and traced the stages of growth of the Leptocephalus and its metamorphosis into the “elvers” or young eels which are carried by the North Atlantic drift back to the coasts of Europe and ascend our rivers in spring in countless myriads; but no man has been more indefatigable and successful in the quest than Dr. Schmidt, who in the various expeditions of the Danish investigation steamer *Thor* from 1904 onwards found successively younger and younger stages, and is during the present summer engaged in a traverse of the Atlantic to the West Indies in the hope of finding the missing link in the chain, the actual spawning fresh-water eel in the intermediate waters somewhere above the abysses of the open ocean.⁵

Again, take the case of an interesting oceanographic observation which, if established, may be found to explain the variations in time and amount of important fisheries. Otto Pettersson in 1910 discovered by his observations in the Gullmar Fjord the presence of periodic submarine waves of deeper saltier water in the Kattegat and the fjords of the west coast of Sweden, which draw in with them from the Jutland banks vast shoals of the herrings which congregate there in autumn. The deeper layer consists of “bank-water” of salinity 32 to 34 per thousand, and as this rolls in along the bottom as a series of huge undulations it forces out the overlying fresher water, and so the herrings living in the “bankwater” outside are sucked into the Kattegat and neighbouring fjords and give rise to important local fisheries. Pettersson connects the crests of the submarine waves with the phases of the moon. Two great waves of saltier water which reached up to the surface took place in November, 1910, one near the time of full moon and the other about new moon, and the latter was at the time when the shoals of herring appeared inshore and provided a profitable fishery. The coincidence of the oceanic phenomena with the lunar phases is not, however, very exact, and doubts have been expressed as to the connection; yet, if established, and even if found to be due, not to the moon, but to prevalent winds or the influence of ocean currents, this would be a case of the migration of fishes depending upon mechanical causes, while in other cases it is known that migrations are due to spawning needs or for the purpose of feeding, as in the case of the cod and the herring in the west and north of Norway and in the Barents Sea.

Then, turning to a very fundamental matter of purely scientific investigation, we do not know with any certainty what causes the great and all-important seasonal variations in the plankton (or floating minute life of the sea) as seen, for example, in our own home seas, where there is a sudden awakening of microscopic plant-life, the Diatoms, in early spring when the water is at its coldest. In the course of a few days the upper layers of the sea may become so filled with organisms that a small silk net towed for a few minutes may capture hundreds of millions of individuals. And these myriads of microscopic forms, after persisting for a few weeks, may disappear as suddenly as they came, to be followed by swarms of Copepoda and many other kinds of minute animals, and these again may give place in the autumn to a second maximum of Diatoms or of the closely related Peridiniales. Of course, there are theories as to all these more or less periodic changes in the plankton, such as Liebig’s “law of the minimum,” which limits the production of an organism by the amount of

that necessity of existence which is present in least quantity, it may be nitrogen or silicon or phosphorus. According to Raben, it is the accumulation of silicic acid in the sea-water that determines the great increase of Diatoms in spring and again in autumn. Some writers have considered these variations in the plankton to be caused largely by changes in temperature supplemented, according to Ostwald, by the resulting changes in the viscosity of the water; but Murray and others are more probably correct in attributing the spring development of phyto-plankton to the increasing power of the sunlight and its value in photosynthesis.

Let us take next the fact—if it be a fact—that the genial, warm waters of the tropics support a less abundant plankton than the cold polar seas. The statement has been made and supported by some investigators and disputed by others, both on a certain amount of evidence. This is possibly a case like some other scientific controversies where both sides are partly in the right or right under certain conditions. At any rate, there are marked exceptions to the generalisation. The German Plankton Expedition in 1889 showed in its results that much larger hauls of plankton per unit-volume of water were obtained in the temperate North and South Atlantic than in the tropics between, and that the warm Sargasso Sea had a remarkably scanty microflora. Other investigators have since reported more or less similar results. Lohmann found the Mediterranean plankton to be less abundant than that of the Baltic; gatherings brought back from tropical seas are frequently very scanty, and enormous hauls, on the other hand, have been recorded from Arctic and Antarctic seas. There is no doubt about the large gatherings obtained in northern waters. I have myself in a few minutes’ haul of a small horizontal net in the north of Norway collected a mass of the large Copepod, *Calanus finmarchicus*, sufficient to be cooked and eaten like potted shrimps by half a dozen of the yacht’s company, and I have obtained similar large hauls in the cold Labrador current near Newfoundland. On the other hand, Kofoid and Alexander Agassiz have recorded large hauls of plankton in the Humboldt current off the west coast of America, and during the *Challenger* Expedition some of the largest quantities of plankton were found in the equatorial Pacific. Moreover, it is common knowledge that on occasions vast swarms of some planktonic organism may be seen in tropical waters. The yellow alga, *Trichodesmium*, which is said to have given its name to the Red Sea, and has been familiarly known as “sea-sawdust” since the days of Cook’s first voyage,⁶ may cover the entire surface over considerable areas of the Indian and South Atlantic Oceans; and some pelagic animals, such as Salpæ, Medusæ, and Ctenophores, are also commonly present in abundance in the tropics. Then, again, American biologists’ have pointed out that the warm waters of the West Indies and Florida may be noted for the richness of their floating life for periods of years, while at other times the pelagic organisms become rare and the region is almost a desert sea.

It is probable, on the whole, that the distribution and variations of oceanic currents have more than latitude or temperature alone to do with any observed scantiness of tropical plankton. These mighty rivers of the ocean in places teem with animal- and plant-life, and may sweep abundance of food from one region to another in the open sea.

But even if it be a fact that there is this alleged deficiency in tropical plankton, there is by no means

⁵ According to Schmidt’s results, the European fresh-water eel, in order to be able to propagate, requires a depth of at least 500 fathoms, a salinity of more than 35·20 per mille, and a temperature of more than 7° C. in the required depth.

⁶ See “Journal” of Sir Joseph Banks. This and other swarms were also noticed by Darwin during the voyage of the *Beagle*.

⁷ A. Agassiz A. G. Mayer, and H. B. Bigelow.

agreement as to the cause thereof. Brandt first attributed the poverty of the plankton in the tropics to the destruction of nitrates in the sea as a result of the greater intensity of the metabolism of denitrifying bacteria in the warmer water; and various writers since then have more or less agreed that the presence of these denitrifying bacteria, by keeping down to a minimum the nitrogen concentration in tropical waters, may account for the relative scarcity of the phyto-plankton, and, consequently, of the zoo-plankton, that has been observed. But Gran, Nathansohn, Murray, Hjort, and others have shown that such bacteria are rare or absent in the open sea, that their action must be negligible, and that Brandt's hypothesis is untenable. It seems clear, moreover, that the plankton does not vary directly with the temperature of the water. Furthermore, Nathansohn has shown the influence of the vertical circulation in the water upon the nourishment of the phyto-plankton—by rising currents bringing up necessary nutrient materials, and especially carbon dioxide, from the bottom layers; and also possibly by conveying the products of the drainage of tropical lands to more polar seas so as to maintain the more abundant life in the colder water.

Pütter's view is that the increased metabolism in the warmer water causes all the available food materials to be used up rapidly, and so puts a check to the reproduction of the plankton.

According to van't Hoff's law in chemistry, the rate at which a reaction takes place is increased by raising the temperature, and this probably holds good for all biochemical phenomena, and therefore for the metabolism of animals and plants in the sea. This has been verified experimentally in some cases by J. Loeb. The contrast between the plankton of Arctic and Antarctic zones, consisting of large numbers of small crustaceans belonging to comparatively few species, and that of tropical waters, containing a great many more species generally of smaller size and fewer in number of individuals, is to be accounted for, according to Sir John Murray and others, by the rate of metabolism in the organisms. The assemblages captured in cold polar waters are of different ages and stages, young and adults of several generations occurring together in profusion,⁸ and it is supposed that the adults "may be ten, twenty, or more years of age." At the low temperature the action of putrefactive bacteria and of enzymes is very slow or in abeyance, and the vital actions of the Crustacea take place more slowly and the individual lives are longer. On the other hand, in the warmer waters of the tropics the action of the bacteria is more rapid, metabolism in general is more active, and the various stages in the life-history are passed through more rapidly, so that the smaller organisms of equatorial seas probably live only for days or weeks in place of years.

This explanation may account also for the much greater quantity of living organisms which has been found so often on the sea-floor in polar waters. It is a curious fact that the development of the polar marine animals is, in general, "direct" without larval pelagic stages, the result being that the young settle down on the floor of the ocean in the neighbourhood of the parent forms, so that there come to be enormous congregations of the same kind of animal within a limited area, and the dredge will in a particular haul come up filled with hundreds, it may be, of an Echinoderm, a Sponge, a Crustacean, a Brachiopod, or an Ascidian; whereas in warmer seas the young pass through a pelagic stage and so become

more widely distributed over the floor of the ocean. The *Challenger* Expedition found in the Antarctic certain Echinoderms, for example, which had young in various stages of development attached to some part of the body of the parents, whereas in temperate or tropical regions the same class of animals set free their eggs and the development proceeds in the open water quite independently of, and it may be far distant from, the parent.

Another characteristic result of the difference in temperature is that the secretion of carbonate of lime in the form of shells and skeletons proceeds more rapidly in warm than in cold water. The massive shells of molluscs, the vast deposits of carbonate of lime formed by corals and by calcareous seaweeds, are characteristic of the tropics; whereas in polar seas, while the animals may be large, they are for the most part soft-bodied and destitute of calcareous secretions. The calcareous pelagic Foraminifera are characteristic of tropical and sub-tropical plankton, and few, if any, are found in polar waters. Globigerina ooze, a calcareous deposit, is abundant in equatorial seas, while in the Antarctic the characteristic deposit is siliceous Diatomaceous ooze.

The part played by bacteria in the metabolism of the sea is very important and probably of wide-reaching effect, but we still know very little about it. A most promising young Cambridge biologist, the late Mr. G. Harold Drew, now unfortunately lost to science, had already done notable work at Jamaica and at Tortugas, Florida, on the effects produced by a bacillus which is found in the surface waters of these shallow tropical seas and in the mud at the bottom; and which denitrifies nitrates and nitrites, giving off free nitrogen. He found that this *Bacillus calis* also caused the precipitation of soluble calcium salts in the form of calcium carbonate ("drewite") on a large scale in the warm shallow waters. Drew's observations tend to show that the great calcareous deposits of Florida and the Bahamas previously known as "coral muds" are not, as was supposed by Murray and others, derived from broken-up corals, shells, nullipores, etc., but are minute particles of carbonate of lime which have been precipitated by the action of these bacteria.⁹

The bearing of these observations upon the formation of oolitic limestones and the fine-grained unfossiliferous Lower Palæozoic limestones of New York State, recently studied in this connection by R. M. Field,¹⁰ must be of peculiar interest to geologists, and forms a notable instance of the annectant character of oceanography, bringing the metabolism of living organisms in the modern sea into relation with palæozoic rocks.

The work of marine biologists on the plankton has been in the main *qualitative*, the identification of species, the observation of structure, and the tracing of life-histories. The oceanographer adds to that the *quantitative* aspect when he attempts to estimate numbers and masses per unit-volume of water or of area. Let me lay before you a few thoughts in regard to some such attempts, mainly for the purpose of showing the difficulties of the investigation. Modern quantitative methods owe their origin to the ingenious and laborious work of Victor Hensen, followed by Brandt, Apstein, Lohmann, and others of the Kiel school of quantitative planktologists. We may take their well-known estimations of fish-eggs in the North Sea as an example of the method.

The floating eggs and embryos of our more important food-fishes may occur in quantities in the plankton during certain months in spring, and Hensen and

⁸ Whether, however, the low temperature may not also retard reproduction is worthy of consideration.

⁹ Journ. Mar. Biol. Assoc., October, 1911.

¹⁰ Carnegie Institution of Washington. "Year Book for 1919," p. 197.

Apstein have made some notable calculations based on the occurrence of these in certain hauls taken at intervals across the North Sea, which led them to the conclusion that, taking six of our most abundant fish, such as the cod and some of the flat-fish, the eggs present were probably produced by about 1,200,000,000 spawners, enabling them to calculate that the total fish population of the North Sea (of these six species) at that time (the spring of 1895) amounted to about 10,000,000,000. Further calculations led them to the result that the fishermen's catch of these fishes amounted to about one-quarter of the total population. Now all this is not only of scientific interest, but also of great practical importance if we could be sure that the samples upon which the calculations are based were adequate and representative, but it will be noted that these samples represent only 1 square metre in 3,465,968,354. Hensen's statement, repeated in various works in slightly differing words, is to the effect that, using a net of which the constants are known hauled vertically through a column of water from a certain depth to the surface, he can calculate the volume of water filtered by the net and so estimate the quantity of plankton under each square metre of the surface; and his whole results depend upon the assumption, which he considers justified, that the plankton is evenly distributed over large areas of water which are under similar conditions. In these calculations in regard to the fish-eggs he takes the whole of the North Sea as being an area under similar conditions, but we have known since the days of P. T. Cleve and from the observations of Hensen's own colleagues that this is not the case, and they have published chart-diagrams showing that at least three different kinds of water under different conditions are found in the North Sea, and that at least five different planktonic areas may be encountered in making a traverse from Germany to the British Isles. If the argument be used that wherever the plankton is found to vary there the conditions cannot be uniform, then few areas of the ocean of any considerable size remain as cases suitable for population-computation from random samples. It may be doubted whether even the Sargasso Sea, which is an area of more than usually uniform character, has a sufficiently evenly distributed plankton to be treated by Hensen's method of estimation of the population.

In the German Plankton Expedition of 1889 Schütt reports that in the Sargasso Sea, with its relatively high temperature, the twenty-four catches obtained were uniformly small in quantity. His analysis of the volumes of these catches shows that the average was 3.33 c.c., but the individual catches ranged from 1.5 c.c. to 6.5 c.c., and the divergence from the average may be as great as +3.2 c.c.; and, after deducting 20 per cent. of the divergence as due to errors of the experiment, Schütt estimates the mean variation of the plankton at about 16 per cent. above or below. This does not seem to me to indicate the uniformity that might be expected in this "halistatic" area occupying the centre of the North Atlantic Gulf Stream circulation. Hensen also made almost simultaneous hauls with the same net in quick succession to test the amount of variation, and found that the average error was about 13 per cent.

As so much depends in all work at sea upon the weather, the conditions under which the ship is working, and the care taken in the experiment, with the view of getting further evidence under known conditions I carried out similar experiments at Port Erin on four occasions during last April and on a further occasion a month later, choosing favourable weather and conditions of tide and wind so as

to be able to maintain an approximate position. On each of four days in April the Nansen net, with No. 20 silk, was hauled six times from the same depth (on two occasions 8 fathoms and on two occasions 20 fathoms), the hauls being taken in rapid succession and the catches emptied from the net into bottles of 5 per cent. formaline, in which they remained until examined microscopically.

The results were of interest, for although they showed considerable uniformity in the amount of the catch—for example, six successive hauls from 8 fathoms being all of them 0.2 c.c., and four out of five from 20 fathoms being 0.6 c.c.—the volume was made up rather differently in the successive hauls. The same organisms occur for the most part in each haul, and the chief groups of organisms are present in much the same proportion. For example, in a series where the Copepoda average about 100, the Dinoflagellates average about 300 and the Diatoms about 8000, but the percentage deviation of individual hauls from the average may be as much as *plus* or *minus* 50. The numbers for each organism (about 40) in each of the twenty-six hauls have been worked out, and the details will be published elsewhere, but the conclusion I come to is that if on each occasion one haul only in place of six had been taken, and if one had used that haul to estimate the abundance of any one organism in that sea-area, one might have been about 50 per cent. wrong in either direction.

Successive improvements and additions to Hensen's methods in collecting plankton have been made by Lohmann, Apstein, Gran, and others, such as pumping up water of different layers through a hose-pipe and filtering it through felt, filter-paper, and other materials which retain much of the micro-plankton that escapes through the meshes of the finest silk. Use has even been made of the extraordinarily minute and beautifully regular natural filter spun by the pelagic animal Appendicularia for the capture of its own food. This grid-like trap, when dissected out and examined under the microscope, reveals a surprising assemblage of the smallest protozoa and proto-phyta, less than 30 micro-millimetres in diameter, which would all pass easily through the meshes of our finest silk nets.

The latest refinement in capturing the minutest-known organisms of the plankton (excepting the bacteria) is a culture method devised by Dr. E. J. Allen, director of the Plymouth Laboratory.¹¹ By diluting half a cubic centimetre of the sea-water with a considerable amount (1500 c.c.) of sterilised water treated with a nutrient solution, and distributing that over a large number (70) of small flasks in which after an interval of some days the number of different kinds of organisms which had developed in each flask was counted, he calculates that the sea contains 464,000 of such organisms per litre; and he gives reasons why his cultivations must be regarded as minimum results, and states that the total per litre may well be something like a million. Thus every new method devised seems to multiply many times the probable total population of the sea. As further results of the quantitative method, it may be recorded that Brandt found about 200 Diatoms per drop of water in Kiel Bay, and Hensen estimated that there are several hundred millions of Diatoms under each square metre of the North Sea or the Baltic. It has been calculated that there is approximately one Copepod in each cubic inch of Baltic water, that the annual consumption of these Copepoda by herring is about a thousand billion, and that in the sixteen square miles of a certain Baltic fishery there is Copepod food for more than 550,000,000 herring of an average weight of 60 grams.

¹¹ Journ. Mar. Biol. Assoc., vol. xii., p. 1, July, 1919.

There are many other problems of the plankton in addition to quantitative estimates—probably some that we have not yet recognised—and various interesting conclusions may be drawn from recent planktonic observations. Here is a case of the introduction and rapid spread of a form new to British seas.

Biddulphia sinensis is an exotic Diatom which, according to Ostenfeld, made its appearance at the mouth of the Elbe in 1903, and spread during successive years in several directions. It appeared suddenly in our plankton gatherings at Port Erin in November, 1909, and has been present in abundance each year since. Ostenfeld in 1908, when tracing its spread in the North Sea, found that the migration to the north along the coast of Denmark to Norway corresponded with the rate of flow of the Jutland current to the Skager Rak, viz. about 17 cm. per second—a case of plankton distribution throwing light on hydrography—and he predicted that it would soon be found in the English Channel. Dr. Marie Lebour, who recently examined the store of plankton gatherings at the Plymouth Laboratory, finds that, as a matter of fact, this form did appear in abundance in the collections of October, 1909, within a month of the time when, according to our records, it reached Port Erin. Whether or not this is an Indo-Pacific species brought accidentally by a ship from the Far East, or whether it is possibly a new mutation which appeared suddenly in our seas, there is no doubt that it was not present in our Irish Sea plankton gatherings previous to 1909, but has been abundant since that year, and has completely adopted the habits of its English relations, appearing with *B. mobiliensis* in late autumn, persisting during the winter, reaching a maximum in spring, and dying out before summer.

The Nauplius and Cypris stages of *Balanus* in the plankton form an interesting study. The adult barnacles are present in enormous abundance on the rocks round the coast, and they reproduce in winter at the beginning of the year. The newly emitted young are sometimes so abundant as to make the water in the shore-pools and in the sea close to the shore appear muddy. The Nauplii first appeared at Port Erin in 1907 in the bay gatherings on February 22 (in 1908 on February 13), and increased with ups and downs to their maximum on April 15, and then decreased until their disappearance on April 26. None were taken at any other time of the year. The Cypris stage follows on after the Nauplius. It was first taken in the bay on April 6, rose to its maximum on the same day with the Nauplii, and was last caught on May 24. Throughout the Cypris curve keeps below that of the Nauplius, the maxima being 1740 and 10,500 respectively. Probably the difference between the two curves represents the death-rate of *Balanus* during the Nauplius stage. That conclusion I think we are justified in drawing, but I would not venture to use the result of any haul, or the average of a number of hauls, to multiply by the number of square yards in a zone round our coast in order to obtain an estimate of the number of young barnacles or of the old barnacles that produced them; the irregularities are too great.

To my mind it seems clear that there must be three factors making for irregularity in the distribution of a plankton organism:

(1) The sequence of stages in its life-history, such as the Nauplius and Cypris stages of *Balanus*.

(2) The results of interaction with other organisms, as when a swarm of *Calanus* is pursued and devoured by a shoal of herring.

(3) Abnormalities in time or abundance due to the physical environment, as in favourable or unfavourable seasons.

And these factors must be at work in the open ocean as well as in coastal waters.

In many oceanographical inquiries there is a double object. There is the scientific interest and there is the practical utility—the interest, for example, of tracing a particular swarm of a Copepod like *Calanus*, and of making out why it is where it is at a particular time, tracing it back to its place of origin, finding that it has come with a particular body of water, and perhaps that it is feeding upon a particular assemblage of Diatoms; endeavouring to give a scientific explanation of every stage in its progress. Then there is the utility—the demonstration that the migration of the *Calanus* has determined the presence of a shoal of herring or mackerel that are feeding upon it, and so have been brought within the range of the fisherman and have constituted a commercial fishery.

We have evidence that pelagic fish which congregate in shoals, such as herring and mackerel, feed upon the Crustacea of the plankton, and especially upon Copepoda. A few years ago when the summer herring fishery off the south end of the Isle of Man was unusually near the land, the fishermen found large red patches in the sea where the fish were specially abundant. Some of the red stuff brought ashore by the men was examined at the Port Erin Laboratory, and found to be swarms of the Copepod, *Temora longicornis*; and the stomachs of the herring caught at the same time were engorged with the same organism. It is not possible to doubt that during these weeks of the herring fishery in the Irish Sea the fish were feeding mainly upon this species of Copepod. Some ten years ago Dr. E. J. Allen and Mr. G. E. Bullen published¹² some interesting work from the Plymouth Marine Laboratory demonstrating the connection between mackerel and Copepoda and sunshine in the English Channel; and Farran¹³ states that in the spring fishery on the West of Ireland the food of the mackerel is mainly composed of *Calanus*.

Then again, at the height of the summer mackerel fishery in the Hebrides in 1913, we found¹⁴ the fish feeding upon the large Copepod, *Calanus finmarchicus*, which was caught in the tow-net at the rate of about 6000 in a five minutes' haul, and 6000 was also the average number found in the stomachs of the fish caught at the same time.

These were cases where the fish were feeding upon the organism that was present in swarms—a monotonous plankton—but in other cases the fish are clearly selective in their diet. If the sardine of the French coast can pick out from the micro-plankton the minute Peridiniales in preference to the equally minute Diatoms which are present in the sea at the same time, there seems no reason why the herring and the mackerel should not be able to select particular species of Copepoda or other large organisms from the macro-plankton, and we have evidence that they do. Nearly thirty years ago the late Mr. Isaac Thompson, a constant supporter of the Zoological Section of this Association and one of the honorary local secretaries for the last Liverpool meeting, showed me in 1893 that young plaice at Port Erin were selecting one particular Copepod, a species of *Jonesiella*, out of many others caught in our tow-nets at the time. H. Blegvad¹⁵ showed in 1916 that young food-fishes, and also small shore-fishes, pick out certain species of Copepoda (such as Harpacticoids) and catch them individually—either lying in wait or searching for them. A couple

¹² Journ. Mar. Biol. Assoc., vol. viii. (1909), pp. 394-406.

¹³ Conseil Internat. Bull. Trimestr., 1902-8, "Planktonique," p. 89.

¹⁴ "Snølla Runiana," lili, Linn. Soc. Journ., Zoology, vol. xxxiv., p. 95, 1918.

¹⁵ Rep. Danish Biol. Stat., vol. xxiv., 1916.

of years later¹⁶ Dr. Marie Lebour published a detailed account of her work at Plymouth on the food of young fishes, proving that certain fish undoubtedly do prefer certain planktonic food.

These Crustacea of the plankton feed upon smaller and simpler organisms—the Diatoms, the Peridinians, and the Flagellates—and the fish themselves in their youngest post-larval stages are nourished by the same minute forms of the plankton. Thus it appears that our sea-fisheries ultimately depend upon the living plankton, which no doubt, in its turn, is affected by hydrographic conditions. A correlation seems to be established between the Cornish pilchard fisheries and periodic variations in the physical characters (probably the salinity) of the water of the English Channel between Plymouth and Jersey.¹⁷ Apparently a diminished intensity in the Atlantic current corresponds with a diminished fishery in the following summer. Possibly the connection in these cases is through an organism of the plankton.

It is only a comparatively small number of different kinds of organisms—both plants and animals—that make up the bulk of the plankton that is of real importance to fish. One can select about half a dozen species of Copepoda which constitute the greater part of the summer zoo-plankton suitable as food for larval or adult fishes, and about the same number of generic types of Diatoms which similarly make up the bulk of the available spring phyto-plankton year after year. This fact gives great economic importance to the attempt to determine with as much precision as possible the times and conditions of occurrence of these dominant factors of the plankton in an average year. An obvious further extension of this investigation is an inquiry into the degree of coincidence between the times of appearance in the sea of the plankton organisms and of the young fish, and the possible effect of any marked absence of correlation in time and quantity.

Just before the war the International Council for the Exploration of the Sea¹⁸ arrived at the conclusion that fishery investigations indicated the probability that the great periodic fluctuations in the fisheries are connected with the fish-larvæ being developed in great quantities only in certain years. Consequently they advised that plankton work should be directed primarily to the question whether these fluctuations depend upon differences in the plankton production in different years. It was then proposed to begin systematic investigation of the fish-larvæ and the plankton in spring and to determine more definitely the food of the larval fish at various stages.

About the same time Dr. Hjort¹⁹ made the interesting suggestion that possibly the great fluctuations in the number of young fish observed from year to year may not depend wholly upon the number of eggs produced, but also upon the relation in time between the hatching of these eggs and the appearance in the water of the enormous quantity of Diatoms and other plant plankton upon which the larval fish, after the absorption of their yolk, depend for food. He points out that if even a brief interval occurs between the time when the larvæ first require extraneous nourishment and that when such food is available, it is highly probable that an enormous mortality would result. In that case even a rich spawning season might yield but a poor result in fish in the commercial fisheries of successive years for some time to come. So that, in fact, the numbers of a year-class may depend not so much upon a favourable spawning season as upon a coincidence between the hatching of

the larvæ and the presence of abundance of phyto-plankton available as food.²⁰

The curve for the spring maximum of Diatoms corresponds in a general way with the curve representing the occurrence of pelagic fish-eggs in our seas. But is the correspondence sufficiently exact and constant to meet the needs of the case? The phyto-plankton may still be relatively small in amount during February and part of March in some years, and it is not easy to determine exactly when, in the open sea, the fish-eggs have hatched out in quantity and the larvæ have absorbed their food-yolk and started feeding on Diatoms.

If, however, we take the case of one important fish—the plaice—we can get some data from our hatching experiments at the Port Erin Biological Station, which have now been carried on for a period of seventeen years. An examination of the hatchery records for these years in comparison with the plankton records of the neighbouring sea, which have been kept systematically for the fourteen years from 1907 to 1920 inclusive, shows that in most of these years the Diatoms were present in abundance in the sea a few days at least before the fish-larvæ from the hatchery were set free, and that it was only in four years (1908, 1909, 1913, and 1914) that there was apparently some risk of the larvæ finding no phyto-plankton food or very little. The evidence so far seems to show that if fish-larvæ are set free in the sea so late as March 20 they are fairly sure of finding suitable food²¹; but if they are hatched so early as February they run some chance of being starved.

But this does not exhaust the risks to the future fishery. C. G. Joh. Petersen and Boysen-Jensen in their valuation of the Limfjord²² have shown that in the case not only of some fish, but also of the larger invertebrates on which they feed, there are marked fluctuations in the number of young produced in different seasons, and that it is only at intervals of years that a really large stock of young is added to the population.

The prospects of a year's fishery may, therefore, depend primarily upon the rate of spawning of the fish, affected, no doubt, by hydrographic and other environmental conditions; secondarily, upon the presence of a sufficient supply of phyto-plankton in the surface-layers of the sea at the time when the fish-larvæ are hatched, and that, in its turn, depends upon photosynthesis and physico-chemical changes in the water; and, finally, upon the reproduction of the stock of molluscs or worms at the bottom which constitute the fish-food at later stages of growth and development.

The question has been raised in recent years: Is there enough plankton in the sea to provide sufficient nourishment for the larger animals, and especially for those fixed forms, such as sponges, that are supposed to feed by drawing currents of plankton-laden water through the body? In a series of remarkable papers from 1907 onwards Pütter and his followers put forward the views: (1) that the carbon requirements of such animals could not be met by the amount of plankton in the volume of water that could be passed through the body in a given time, and (2) that sea-water contained a large amount of dissolved organic carbon compounds which constitute the chief, if not the only, food of a large number of marine animals. These views have given rise to

²⁰ For the purpose of this argument we may include in "phyto-plankton" the various groups of Flagellata and other minute organisms which may be present with the Diatoms.

²¹ All dates and statements as to occurrence refer to the Irish Sea round the south end of the Isle of Man. For further details see Report Lancs Sea-Fish. Lab. for 1919.

²² Report of Danish Bio'l. Station for 1919.

¹⁶ Journ. Mar. Biol. Assoc., May, 1918.

¹⁷ See E. C. Jee, "Hydrography of the English Channel," 1904-17.

¹⁸ Rapports et Proc. Verb., vol. xix., December, 1913.

¹⁹ *Ibid.*, vol. xx., 1914, p. 204.

much controversy, and have been useful in stimulating further research, but I believe it is now admitted that Pütter's samples of water from the Bay of Naples and at Kiel were probably polluted, that his figures were erroneous, and that his conclusions must be rejected, or at least greatly modified. His estimates of the plankton were minimum ones, while it seems probable that his figures for the organic carbon present represent a variable amount of organic matter arising from one of the reagents used in the analyses.²³ The later experimental work of Henze, of Raben, and of Moore shows that the organic carbon dissolved in sea-water is an exceedingly minute quantity, well within the limits of experimental error. Moore puts it at the most at one-millionth part, or 1 mgm. in a litre. At the Dundee meeting of the Association in 1912 a discussion on this subject took place, at which Pütter still adhered to a modified form of his hypothesis of the inadequacy of the plankton and the nutrition of lower marine animals by the direct absorption of dissolved organic matter. Further work at Port Erin since has shown that, while the plankton supply as found generally distributed would prove sufficient for the nutrition of such sedentary animals as Sponges and Ascidians, which require to filter only about fifteen times their own volume of water per hour, it is quite inadequate for active animals such as crustaceans and fishes. These latter are, however, able to seek out and capture their food, and are not dependent on what they may filter or absorb from the sea-water. This result accords well with recorded observations on the irregularity in the distribution of the plankton and with the variations in the occurrence of the migratory fishes which may be regarded as following and feeding upon the swarms of planktonic organisms.

This, then, like most of the subjects I am dealing with, is still a matter of controversy, still not completely understood. Our need, then, is research, more research, and *still more research*.

Our knowledge of the relations between plankton productivity and variation and the physico-chemical environment is still in its infancy, but gives promise of great results in the hands of the biochemist and the physical chemist.

Recent papers by Sørensen, Palitzsch, Witting, Moore, and others have made clear that the amount of hydrogen-ion concentration as indicated by the relative degree of alkalinity and acidity in the sea-water may undergo local and periodic variations, and that these have an effect upon the living organisms in the water and can be correlated with their presence and abundance. To take an example from our own seas, Prof. Benjamin Moore and his assistants in their work at the Port Erin Biological Station in successive years from 1912 onwards have shown²⁴ that the sea around the Isle of Man is a good deal more alkaline in spring (say April) than it is in summer (say July). The alkalinity, which gets low in summer, increases somewhat in autumn, and then decreases rapidly, to disappear during the winter; and then once more, after several months of a minimum, begins to come into evidence again in March, and rapidly rises to its maximum in April or May. This periodic change in alkalinity will be seen to correspond roughly with the changes in the living microscopic contents of the sea represented by the phyto-plankton annual curve, and the connection between the two will be seen when we realise that the alkalinity of the sea is due to the relative absence of carbon dioxide. In early spring, then, the developing myriads of

Diatoms in their metabolic processes gradually use up the store of carbon dioxide accumulated during the winter or derived from the bicarbonates of calcium and magnesium, and so increase the alkalinity of the water until the maximum of alkalinity, due to the fixation of the carbon and the reduction in amount of carbon dioxide, corresponds with the crest of the phyto-plankton curve in, say, April. Moore has calculated that the annual turnover in the form of carbon which is used up or converted from the inorganic into an organic form probably amounts to something of the order of 20,000–30,000 tons of carbon per cubic mile of sea-water, or, say, over an area of the Irish Sea measuring 16 square miles and a depth of 50 fathoms; and this probably means a production each season of about two tons of dry organic matter, corresponding to at least ten tons of moist vegetation, per acre—which suggests that we may still be very far from getting from our seas anything like the amount of possible food-matters that are produced annually.

Testing the alkalinity of the sea-water may therefore be said merely to be ascertaining and measuring the results of the photosynthetic activity of the great phyto-plankton rise in spring due to the daily increase of sunlight.

The marine biologists of the Carnegie Institution, Washington, have made a recent contribution to the subject in certain observations on the alkalinity of the sea (as determined by hydrogen-ion concentration), during which they found in tropical mid-Pacific a sudden change to acidity in a current running eastwards. Now in the Atlantic the Gulf Stream and tropical Atlantic waters generally are much more alkaline than the colder coastal water running south from the Gulf of St. Lawrence—that is, the colder Arctic water has more carbon dioxide. This suggests that the Pacific easterly set may be due to deeper water, containing more carbon dioxide (=acidity), coming to the surface at that point. The alkalinity of the sea-water can be determined rapidly by mixing the sample with a few drops of an indicator and observing the change in colour; and this method of detecting ocean currents by observing the hydrogen-ion concentration of the water might be useful to navigators as showing the time of entrance to a known current.

Oceanography has many practical applications, chiefly, but by no means wholly, on the biological side. The great fishing industries of the world deal with living organisms, of which all the vital activities and the inter-relations with the environment are matters of scientific investigation. Aquiculture is as susceptible of scientific treatment as agriculture can be; and the fisherman, who has been in the past too much the nomad and the hunter—if not, indeed, the devastating raider—must become in the future the settled farmer of the sea if his harvest is to be less precarious. Perhaps the nearest approach to cultivation of a marine product, and of the fisherman reaping what he has actually sown, is seen in the case of the oyster and mussel industries on the west coast of France, in Holland, America, and, to a less extent, on our own coast. Much has been done by scientific men for these and other similar coastal fisheries since the days when Prof. Coste in France in 1859 introduced oysters from the Scottish oyster-beds to start the great industry at Arcachon and elsewhere. Now we buy back the descendants of our own oysters from the French ostricultrists to replenish our depleted beds.

It is no small matter to have introduced a new and important food-fish to the markets of the world. The remarkable deep-water "tile-fish," new to science and

²³ See Moore, etc., *Bio-Chem. Journ.*, vol. vi., p. 266, 1912.

²⁴ "Photosynthetic Phenomena in Sea-water," *Trans. Liverpool Biol. Soc.*, vol. xxix., p. 233, 1915.

described as *Lopholatilus chamaeleonticeps*, was discovered in 1879 by one of the United States fishing schooners to the south of Nantucket, near the 100-fathom line. Several thousand pounds' weight was caught, and the matter was duly investigated by the United States Fish Commission. For a couple of years after that the fish was brought to market in quantity, and then something unusual happened at the bottom of the sea, and in 1882 millions of dead tile-fish were found floating on the surface over an area of thousands of square miles. The schooner *Navarino* sailed for two days and a night through at least 150 miles of sea thickly covered, so far as the eye could reach, with dead fish, estimated at 256,000 to the square mile. The Fish Commission sent a vessel to fish systematically over the grounds known as the "Gulf Stream slope," where the tile-fish had been so abundant during the two previous years, but she did not catch a single fish, and the associated sub-tropical invertebrate fauna was also practically obliterated.

This wholesale destruction was attributed by the American oceanographers to a sudden change in the temperature of the water at the bottom, due in all probability to a withdrawal southwards of the warm Gulf Stream water and a flooding of the area by the cold Labrador current.

I am indebted to Dr. C. H. Townsend, director of the celebrated New York Aquarium, for the latest information in regard to the reappearance in quantity of this valuable fish upon the old fishing-grounds of Nantucket and Long Island, at about 100 miles from the coast to the east and south-east of New York. It is believed that the tile-fish is now abundant enough to maintain an important fishery, which will add an excellent food-fish to the markets of the United States. It is easily caught with lines at all seasons of the year, and reaches a length of more than 3 ft. and a weight of 40-50 lb. During July, 1915, the product of the fishery was about 2,500,000 lb. weight, valued at 55,000 dollars, and in the first few months of 1917 the catch was 4,500,000 lb., for which the fishermen received 247,000 dollars.

We can scarcely hope in European seas to add new food-fishes to our markets, but much may be done through the co-operation of scientific investigators of the ocean with the administrative departments to bring about a more rational conservation and exploitation of the national fisheries.

Earlier in this address I referred to the pioneer work of the distinguished Manx naturalist, Prof. Edward Forbes. There are many of his writings and of his lectures to which I have no space to refer which have points of oceanographic interest. Take this, for example, in reference to our national sea-fisheries. We find him in 1847 writing to a friend: "On Friday night I lectured at the Royal Institution. The subject was the bearing of submarine researches and distribution matters on the fishery question. I pitched into Government mismanagement pretty strong, and made a fair case of it. It seems to me that at a time when half the country is starving we are utterly neglecting or grossly mismanaging great sources of wealth and food. . . . Were I a rich man I would make the subject a hobby for the good of the country and for the better proving that the true interests of Government are those linked with and inseparable from Science." We must still cordially approve of these last words, while recognising that our Government Department of Fisheries is now being organised on better lines, is itself carrying on scientific work of national importance, and is, I am happy to think, in complete sympathy with the work of independent scientific investigators of the sea and desirous of closer

co-operation with university laboratories and biological stations.

During recent years one of the most important and most frequently discussed of applications of fisheries investigation has been the productivity of the trawling grounds, and especially those of the North Sea. It has been generally agreed that the enormous increase of fishing power during the last forty years or so has reduced the number of large plaice, so that the average size of that fish caught in our home waters has become smaller, although the total number of plaice landed had continued to increase up to the year of the outbreak of war. Since then, from 1914 to 1919, there has of necessity been what may be described as the most gigantic experiment ever seen in the closing of extensive fishing-grounds. It is still too early to say with any certainty exactly what the results of that experiment have been, although some indications of an increase of the fish population in certain areas have been recorded. For example, the Danes, A. C. Johansen and Kirstine Smith, find that large plaice landed in Denmark are now more abundant, and they attribute this to a reversal of the pre-war tendency, due to less intensive fishing. But Dr. James Johnstone has pointed out that there is some evidence of a natural periodicity in abundance of such fish, and that the results noticed may represent phases in a cyclic change. If the periodicity noted in Liverpool Bay²⁵ holds good for other grounds, it will be necessary in any comparison of pre-war and post-war statistics to take this natural variation in abundance into very careful consideration.

In the application of oceanographic investigations to sea-fisheries problems one ultimate aim, whether frankly admitted or not, must be to obtain some kind of a rough approximation to a census or valuation of the sea—of the fishes that form the food of man, of the lower animals of the sea-bottom on which many of the fishes feed, and of the planktonic contents of the upper waters which form the ultimate organised food of the sea—and many attempts have been made in different ways to attain the desired end.

Our knowledge of the number of animals living in different regions of the sea is for the most part relative only. We know that one haul of the dredge is larger than another, or that one locality seems richer than another, but we have very little information as to the actual numbers of any kind of animal per square foot or per acre in the sea. Hensen, as we have seen, attempted to estimate the number of food-fishes in the North Sea from the number of their eggs caught in a comparatively small series of hauls of the tow-net, but the data were probably quite insufficient and the conclusions may be erroneous. It is an interesting speculation to which we cannot attach any economic importance. Heincke says of it: "This method appears theoretically feasible, but presents in practice so many serious difficulties that no positive results of real value have as yet been obtained."

All biologists must agree that to determine even approximately the number of individuals of any particular species living in a known area is a contribution to knowledge which may be of great economic value in the case of the edible fishes, but it may be doubted whether Hensen's methods, even with greatly increased data, will ever give us the required information. Petersen's method, of setting free marked plaice and then assuming that the proportion of these recaptured is to the total number marked as the fishermen's catch in the same district is to the total population, will hold good only in circumscribed areas where there is practically no migration and the fish

²⁵ See Johnstone, Report Lancs Sea-Fish. Lab. for 1917, p. 60; and Daniel, Report for 1919, p. 51.

are fairly evenly distributed. This method gives us what has been called "the fishing coefficient," and this has been estimated for the North Sea to have a probable value of about 0.33 for those sizes of fish which are caught by the trawl. Heincke,²⁶ from an actual examination of samples of the stock on the ground obtained by experimental trawling ("the catch coefficient"), supplemented by the market returns of the various countries, estimates the adult plaice at about 1,500,000,000, of which about 500,000,000 are caught or destroyed by the fishermen annually.

It is difficult to imagine any further method which will enable us to estimate any such case as, say, the number of plaice in the North Sea, where the individuals are so far beyond our direct observation and are liable to change their positions at any moment. But a beginning can be made on more accessible ground with more sedentary animals, and Dr. C. G. Joh. Petersen, of the Danish Biological Station, has for some years been pursuing the subject in a series of interesting reports on "The Evaluation of the Sea."²⁷ He uses a bottom-sampler or grab, which can be lowered down open and then closed on the bottom so as to bring up a sample square foot or square metre (or in deep water one-tenth of a square metre) of the sand or mud and its inhabitants. With this apparatus, modified in size and weight for different depths and bottoms, Petersen and his fellow-workers have made a very thorough examination of the Danish waters, and especially of the Kattegat and the Limfjord, have described a series of "animal communities" characteristic of different zones and regions of shallow water, and have arrived at certain numerical results as to the quantity of animals in the Kattegat expressed in tons—such as 5000 tons of plaice requiring as food 50,000 tons of "useful animals" (mollusca and polychæt worms), and 25,000 tons of starfish using up 200,000 tons of useful animals which might otherwise serve as food for fishes, and the dependence of all these animals directly or indirectly upon the great Beds of *Zostera*, which make up 24,000,000 tons in the Kattegat. Such estimates are obviously of great biological interest, and, even if only rough approximations, are a valuable contribution to our understanding of the metabolism of the sea and of the possibility of increasing the yield of local fisheries.

But on studying these Danish results in the light of what we know of our own marine fauna, although none of our seas have been examined in the same detail by the bottom-sampler method, it seems probable that the animal communities as defined by Petersen are not exactly applicable on our coasts, and that the estimates of relative and absolute abundance may be very different in different seas under different conditions. The work will have to be done in each great area, such as the North Sea, the English Channel, and the Irish Sea, independently. This is a necessary investigation, both biological and physical, which lies before the oceanographers of the future, upon the results of which the future preservation and further cultivation of our national sea-fisheries may depend.

It has been shown by Johnstone and others that the common edible animals of the shore may exist in such abundance that an area of the sea may be more productive of food for man than a similar area of pasture or crops on land. A Lancashire mussel-bed has been shown to have as many as 16,000 young mussels per square foot, and it is estimated that in the shallow

waters of Liverpool Bay there are from 20 to 200 animals of sizes varying from an amphipod to a plaice on each square metre of the bottom.²⁸

From these and similar data which can be readily obtained it is not difficult to calculate totals by estimating the number of square yards in areas of similar character between tide-marks or in shallow water. And from weighings of samples some approximation to the number of tons of available food may be computed. But one must not go too far. Let all the figures be based upon actual observation. Imagination is necessary in science, but in calculating a population of even a very limited area it is best to believe only what one can see and measure.

Countings and weighings, however, do not give us all the information we need. It is something to know even approximately the number of millions of animals on a mile of shore and the number of millions of tons of possible food in a sea area, but that is not sufficient. All food-fishes are not equally nourishing to man, and all plankton and bottom invertebrates are not equally nourishing to a fish. At this point the biologist requires the assistance of the physiologist and the biochemist. We want to know next the value of our food-matters in proteids, carbohydrates, and fats, and the resulting Calories. Dr. Johnstone, of the oceanography department of the University of Liverpool, has already shown us how markedly a fat summer herring differs in essential constitution from the ordinary white fish, such as the cod, which is almost destitute of fat.

Prof. Brandt at Kiel, Prof. Benjamin Moore at Port Erin, and others have similarly shown that plankton gatherings may vary greatly in their nutrient value according as they are composed mainly of Diatoms, of Dinoflagellates, or of Copepoda. And, no doubt, the animals of the "benthos," the common invertebrates of our shores, will show similar differences in analysis.²⁹ It is obvious that some contain more solid flesh, others more water in their tissues, others more calcareous matter in the exoskeleton, and that therefore, weight for weight, we may be sure that some are more nutritious than others; and this is probably at least one cause of that preference we see in some of our bottom-feeding fish for certain kinds of food, such as polychæt worms, in which there is relatively little waste, and thin-shelled lamellibranch molluscs, such as young mussels, which have a highly nutrient body in a comparatively thin and brittle shell.

My object in referring to these still incomplete investigations is to direct attention to what seems a natural and useful extension of faunistic work for the purpose of obtaining some approximation to a quantitative estimate of the more important animals of our shores and shallow water and their relative values as either the immediate or the ultimate food of marketable fishes.

Each such fish has its "food-chain" or series of alternative chains, leading back from the food of man to the invertebrates upon which it preys, and then to the food of these, and so down to the smallest and simplest organisms in the sea, and each such chain must have all its links fully worked out as to seasonal and quantitative occurrence back to the Diatoms and Flagellates, which depend upon physical conditions, and take us beyond the range of biology, but not beyond that of oceanography. The Diatoms and the Flagellates are probably more important than the more obvious seaweeds not only as food, but also in

²⁸ "Conditions of Life in the Sea," Cambridge University Press, 1908.

²⁶ F. Heincke, Cons. Per. Internat. Explor. de la Mer, "Investigations on the Plaice," Copenhagen, 1913.

²⁷ See Reports of the Danish Biological Station, and especially the Report for 1918, "The Sea Bottom and its Production of Fish Food."

²⁹ Moore and others have made analyses of the protein, fat, etc., in the soft parts of Sponge, Ascidian, Aplousia, Fusus, Echinus, and Cancer at Port Erin, and find considerable differences—the protein ranging, for example, from 8 to 51 per cent., and the fat from 2 to 14 per cent. (see *Bio-Chemical Journ.*, vol. vi., p. 291).

supplying to the water the oxygen necessary for the respiration of living protoplasm. Our object must be to estimate the rate of production and rate of destruction of all organic substances in the sea.

To attain to an approximate census and valuation of the sea—remote though it may seem—is a great aim, but it is not sufficient. We want not only to observe and to count natural objects, but also to understand them. We require to know not merely what an organism is—in the fullest detail of structure and development and affinities—where it occurs—again in full detail—and in what abundance in different circumstances, but also *how* it lives and what all its relations are to both its physical and its biological environment, and that is where the physiologist, and especially the biochemist, can help us. In the best interests of biological progress the day of the naturalist who merely collects, the day of the anatomist and histologist who merely describe, is over, and the future is with the observer and the experimenter animated by a divine curiosity to enter into the life of the organism and understand how it lives and moves and has its being. "Happy indeed is he who has been able to discover the causes of things."

Cardiff is a seaport, and a great seaport, and the Bristol Channel is a notable sea-fisheries centre of growing importance. The explorers and merchant venturers of the south-west of England are celebrated in history. What are you doing now in Cardiff to advance our knowledge of the ocean? You have here an important university centre and a great modern

national museum, and either or both of these homes of research might do well to establish an oceanographical department, which would be an added glory to your city and of practical utility to the country. This is the obvious centre in Wales for a sea-fisheries institute for both research and education. Many important local movements have arisen from British Association meetings, and if such a notable scientific development were to result from the Cardiff meeting of 1920, all who value the advance of knowledge and the application of knowledge to industry would applaud your enlightened action.

In a wider sense, it is not to the people of Cardiff alone that I appeal, but to the whole population of these islands, a maritime people who owe everything to the sea. I urge them to become better informed in regard to our national sea-fisheries and to take a more enlightened interest in the basal principles that underlie a rational regulation and exploitation of these important industries. National efficiency depends to a very great extent upon the degree in which scientific results and methods are appreciated by the people and scientific investigation is promoted by the Government and other administrative authorities. The principles and discoveries of science apply to aquiculture no less than to agriculture. To increase the harvest of the sea the fisheries must be continuously investigated, and such cultivation as is possible must be applied, and all this is clearly a natural application of the biological and hydrographical work now united under the science of oceanography.

Summaries of Addresses of Presidents of Sections of the British Association.

Mathematical and Physical Science.

PROF. EDDINGTON's presidential address to Section A deals with the investigation of the internal conditions of the stars. Most of the naked-eye stars have densities so low that they may be treated as spheres of perfect gas (giant stars). In familiar hot bodies the energy existing in the aether (radiant heat) is extremely small compared with that associated with the matter (molecular motions); conditions might exist in which this disproportion was reversed; but the stars are of just such a mass that the two kinds of energy are roughly equal. It is thought that this balance cannot be a coincidence, but determines why the masses of the stars are always close to a particular value. From astronomical data as to the masses and radiation of the stars it is possible to determine the opacity of stellar material to the radiation traversing it. The opacity turns out to be very high and of the same order of magnitude as that found for X-rays in the laboratory. (At the high temperatures in the stars the radiation consists mainly of soft X-rays.) A rather surprising result is that the opacity varies very little with the temperature of the star or wave-length of the radiation. The discussion leads to many astronomical results which appear to be generally confirmed by observation; in particular, it fixes within fairly narrow limits the period of a mechanical pulsation of any star, and this agrees in all known Cepheid variables with the observed period of light-pulsation. The question of the source of a star's heat is raised in an acute form by these investigations. It appears that the energy of gravitational contraction is quite inadequate. The recent experimental results of Aston and Rutherford seem to throw some new light on the often-discussed question whether sub-atomic energy can be made available in the stars. The address concludes with some observations on the legitimate place of speculation in scientific research.

NO. 2652, VOL. 105]

Chemistry.

MR. C. T. Heycock deals in his presidential address to Section B with the manner in which our present rather detailed knowledge of metallic alloys has been acquired, starting from the sparse information which was available thirty or forty years ago, and sketches briefly the present position of the subject. He considers chiefly the non-ferrous alloys, not because any essential difference in type exists between these and ferrous alloys, but because the whole field presented by the chemistry of the metals and their alloys is too vast to be covered in an address of reasonable length. Though Réaumur in 1722 employed the microscope to examine the fractured surfaces of white and grey cast-iron and steel, and Widmanstätten in 1808 polished and etched sections from meteorites, the founder of modern metallography is undoubtedly H. C. Sorby, whose methods of polishing and etching alloys and of vertical illumination are used to-day by all who work at this subject. The first important clue to what occurs on cooling a fused mixture of metals was given by Guthrie's experiments on cryohydrates, and these researches, with those of Sorby, undertaken as they were for the sake of investigating natural phenomena, are remarkable examples of how purely scientific experiment can lead to most important practical results. Raoult's work on the depression of the freezing point of solvents due to the addition of dissolved substances led to the establishment by van't Hoff of a general theory applicable to all solutions. Later experiments established the similarity between the behaviour of metallic solutions or alloys and that of aqueous and other solutions of organic compounds in organic solvents; and in 1897 Neville and Heycock determined the complete freezing-point curve of the copper-tin alloys, confirming and extending the work of Roberts-Austen, Stansfield, and Le Chatelier. These were probably the first of the binary alloys on