1916-17, including a high-grade copper ore from a boulder near Port Lockroy on the Neumayer Channel. The description of the varied lavas of Deception Island in both this and Mr. Tyrrell's paper is surely an argument in favour of collation and co-operation.

The reports of the British Antarctic ("Terra Nova") Expedition of 1910, published by the British Museum, are concerned, however, with another quadrant of the antarctic region. In the geological series, Nos. 3 and 4, recent and older sedimentary deposits are described by Mr. F. Debenham, from

his personal observations as geologist to the expedition. Metamorphic rocks are dealt with in No. 5; but their relationships in the field are as yet but little known. Dr. A. Smith Woodward's account (No. 2) of "Fish remains from the Upper Old Red Sandstone of Granite Harbour," including *Bothriolepis*, provides more definite information. The remoteness and perils of the district excuse the uncorrelated nature of the results, and no doubt also explain the handsome method of their presentation in the year 1921-22.

G. A. J. C.

Durability of Optical Glass.

By Dr. JAMES WEIR FRENCH.

IT is the custom of optical glass manufacturers to issue long lists of types characterised by their optical constants, without much regard to their qualities of durability, which are only occasionally indicated. To the practical computer these lengthy lists are not imposing. Experience has taught him that the number of sufficiently trustworthy types is really very limited, and that only in exceptional circumstances, that fortunately do not frequently arise, may an extension of his list be justified. But while the possibility remains that glass of an unstable kind may be used in the construction of his instruments, the optician has cause for anxiety in the knowledge that his reputation as well as the glass may become tarnished.

The British Scientific Instrument Research Association, the function of which is to provide for the industry the leaven of science, has been charged with the investigation of the durability of glass. According to the admirable report¹ that has recently been issued, "the object of the Research was to determine how far it was possible to establish simple tests by which the durability of different types and varieties of optical glass could be quickly ascertained without awaiting the results of experience by actual use over an extended period." To what extent this object has been attained may be realised from the frank confession at the conclusion of the report, that "it is not possible to recommend any simple test by which the durability of an optical glass can be determined, with such reliability as to avoid the chance of misleading users of the glass in some one or other application of it." With this pessimistic conclusion it is difficult to agree, as the object has already been attained in the workshop, if it is agreed that it is the reflecting or transmitting qualities of the surface with which the optician is concerned.

From the report it appears that numerous tests of Continental and British types of optical glass have

been examined by the iodoeosin test of Mylius, the autoclave water and steam tests at four and two atmospheres of pressure, and the "dimming" test evolved by the Royal Arsenal Directorate of Chemical Inspection, which co-operated in the research. This dimming test is really an elaboration of the Zschimmer test. The three tests as applied indicate merely to what extent alkali can be dissolved from the surface, and, as is to be expected, the flint glass types appear more durable than the crown types—a conclusion that is misleading, as the report rightly indicates. Our industrial atmosphere unfortunately contains sulphuretted hydrogen, and if in the dimming test an atmosphere more representative of reality had been employed, the flint types would have been placed more nearly in the order accorded to them by Faraday.

The report confirms the interesting fact, already known in the workshop, that in the glass-polishing process alkali is dissolved from the surface layer, which, with a few exceptions, becomes more durable. Workshop experience shows that a new cloth polisher tested with litmus will usually be found to be slightly acid; after a few hours of working it will be neutral; and thereafter it becomes strongly alkaline. A pitch polisher reacts similarly, but it does not retain the dissolved alkali to the same extent.

The optician is concerned in practice not so much with those so-called optical glasses that are visually affected by the tests referred to, as with those that are labelled as being durable and unaffected. If a well-polished specimen of the most durable crown glass be boiled in water at atmospheric pressure for two hours and its reflecting power be then tested by means of a multi-reflection photometer, a loss of $\frac{1}{2}$ per cent. per reflection may be detected. After boiling for eight hours, the loss will be about 2 per cent., but thereafter the rate diminishes.

The drastic autoclave tests adopted in the research are not necessary to demonstrate how many of the types included in the optical glass-makers' lists are vitreous substances of but little value and a source of danger to the unwary.

¹ Report of an Investigation on the Determination of the Durability of Optical Glass carried out by T. Haigh. Pp. 51 + 10 plates. (British Scientific Instrument Research Association, 26 Russell Square, W.C.1.) 7s. 6d.

Volcanic Activity in Nigeria.

REPORTS have been received by the Governor from Mr. A. A. Reading, of the Bibundi Estate, of a recent volcanic eruption in the Cameroons Mountain, near the coast of Nigeria. Repeated earthquake shocks commencing on February 3 last were followed by an eruption at an altitude of about 4000 feet on the north-west side of the mountain, giving rise to a lava stream which flowed down in the direction of the Bibundi Estate, and entered the plantation area on March 3. The lava extended, roughly, one-third of a mile out to sea, and huge

columns of steam were continually ascending. Attempts to photograph the scene failed on account of the dense smoke and ashes. The centre of the group of craters was estimated to be distant 9900 yards from the house at Bibundi Beach on a magnetic bearing of 128°, and the height above sea-level was believed to be about 4150 feet.

In May the northern stream was still advancing, and threatened to reach the sea along the water-courses on each side of Dollmanshohe. Sometimes there was a loud noise resembling that of a blast