the earliest days of solar spectroscopy, the advantages of the spectrohelioscope in quickly interpreting them must be seen to be appreciated. The line-shifter serves in somewhat the same way as the 'blink' device of a Zeiss stereocomparator, but instead of merely indicating, in a very striking way, any differences between two photographs, it shows at a glance the connexion of a series of related phenomena by linking them into a sequence. This is what I referred to in my last article when describing "a new effect of inflow," by which I meant the motion of the maximum of intensity along a flocculus due to progressive differences in radial velocity, at once interpreted by the line-shifter.

In M. Deslandres's remarkable theory of the sun, "tout se passe comme si le rayonnement corpusculaire de nos orages émane d'une couche

solaire profonde qui tourne comme un corps solide ; et cette couche offre au moins 24 volcans permanents, répartis uniformément autour de l'axe de rotation, et d'activité variable, qui rejettent au dehors la matière ionisée ou radioactive des masses intérieures. Cette division régulière est celle des corps à symétrie circulaire qui se refroidissent." This theory involves so many considerations that it cannot be discussed in the space at my disposal. In any event, such a discussion would be premature, as the theory rests upon M. Deslandres's belief in a regular sequence of terrestrial magnetic disturbances which Messrs. Chree and Stagg, after an exhaustive study of the international magnetic data for twenty years, have been unable to confirm.⁵

⁵ "Recurrence Phenomena in Terrestrial Magnetism," *Phil. Trans.*, Series A, vol. 227, pp. 21-62.

The New Vision.¹

By Prof. G. ELLIOT SMITH, F.R.S.

THE recent researches of Minkowski, Brouwer, and Woollard have brought to light the remarkable fact that in man and his nearest allies such revolutionary changes have been effected in every part of the visual system as to justify the statement that a new visual instrument has been evolved. Considering these facts from the point of view of function, we may speak of the emergence of a new vision, which differs profoundly from that enjoyed by all other living creatures. It may be of interest to discuss the nature and meaning of these changes, which are intimately related to the evolution of intelligence and the attainment of what in colloquial language we know as insight, foresight, and the wider vision.

A quarter of a century ago the term neopallium was introduced to express a new conception of the nature of the essential evolutionary changes that transformed the brain of a primitive reptile into that of a mammal, and conferred upon the latter enormously enhanced powers of learning from experience and modifying behaviour. Incidentally, the development of the new cortical area provided a means whereby vision for the first time secured representation in the cerebral cortex, which created fuller opportunities for the confluence of visual with other kinds of perception. On the anatomical side it inaugurated a series of interesting structural arrangements which marked the transference to the neopallium of functions which in all other vertebrate animals are carried on in the mid-brain and other lower centres. The new light recently thrown on this interesting process of transformation reveals the fact that the transference of functions did not occur at one time, but in progressive stages. Magnus showed experimentally that the optic-righting reflexes in mammals were not acquired until binocular vision developed. Still later the development of the macula lutea in the retina (in monkeys and man) brought about the

further changes in the brain which I am attempting to define.

In all vertebrates the nerve fibres proceeding from the retina cross (wholly or only in part in most mammals) to the other side of the brain, where they end in two masses of grey matter, the lateral geniculate body, which is part of the thalamus, and the superior quadrigeminal body, which is part of the mid-brain. The former connexion is concerned with the awareness to vision, the phenomena of consciousness, and the latter (mid-brain), with such unconscious functions as the reflex actions of the eye muscles and the general musculature of the whole body. Brouwer has shown that in a lowly mammal such as a rabbit the four quadrants of the retina have a topographical representation in the quadrigeminal body. Wilson (of Cairo) has recently demonstrated that the corresponding quadrants in a lizard's brain control definite movements or postures of the body—a kind of autonomous mechanism for the analysis and functional expression of optic influences analogous to the analytic functions of the semicircular canals in connexion with equilibration.

In mammals the lateral geniculate body, for the first time in the vertebrate series, emits a large strand of fibres (optic radiation) to provide a path for visual impulses to the cerebral cortex. But the neopallium also begins to assume some of the motor control, which hitherto has been a function of the quadrigeminal bodies. It is interesting to note that, according to Allen, this process is not completed in the rabbit. Its cerebral cortex, according to him, controls the movements of the head, forelimbs, and body, but the control of the hind-limbs is still retained by the mid-brain. In most mammals, however, the transference of motor control to the cerebral cortex is complete.

With the acquisition of binocular vision (in mammals such as the cat or, better, monkeys) the fibres of the optic tracts become rearranged. The fibres from the lateral part of each retina no longer

¹ From the Bowman Lecture delivered on April 20 to the Ophthalmological Society of the United Kingdom.

cross to the other side of the brain, but become connected with the same side, so as to bring into connexion the terminations of the fibres coming from the medial side of one retina and the lateral side of the other, which in binocular vision necessarily act together so as to merge in consciousness the two images of one object.

But this rearrangement of the optic tracts necessarily affects the endings of these tracts in the geniculate and quadrigeminal bodies. Instead of a modification of the retinal localisation in the quadrigeminal body to adapt it to the new conditions, the cerebral cortex seems more fully to usurp its motor-controlling functions. With the loss of such functions the quadrigeminal body also loses most of the direct connexions with the optic tracts, and the cerebral cortex acquires a correspondingly enhanced control of the quadrigeminal body.

In monkeys and man further profound changes occur in the whole of the visual system. A definite macula lutea develops in the retina, and each of the percipient cells in the area of acute vision transmits its impulse (indirectly) to a separate fibre of the optic nerve. In the rest of the retina and in the retinas of other mammals groups of sensory cells (rods) transmit their impulses into one granule and ganglion cell, so that there are far more percipient elements than nerve fibres in the optic nerve. Hence, when the macula develops in monkeys and man, this small area adds a contribution to the optic nerve and tract that is out of all proportion to its size. The macular fibres form more than a third of the optic nerve, and there is added to the geniculate body a new formation as a macular receptive mechanism.

With the atrophy of the quadrigeminal fibres of the optic tract and the sudden increase of the geniculate connexion in monkeys and man, practically the whole (more than ninety per cent.) of the optic fibres go to the lateral geniculate body. But with the enormous increase of the latter the body loses much of its autonomy. Its ventral nucleus, which in other vertebrates controlled the quadri-

geminal body, atrophies in the Primates. In its place the cerebral connexion is still further strengthened. The geniculate body becomes more and more an intermediary between the retina and the neopallium, and almost the whole function of visual perception becomes concentrated in the cerebral cortex.

The development of macular vision confers upon man the ability to see the world and appreciate its meaning in a way that no other living creature is able to do. His new vision depends upon powers of visual perception as distinctive as the use of articulate speech to give expression to what he sees and thinks. The late Dr. Henry Watt, of the University of Glasgow, expressed the opinion that

"Of the conditions which enrich the sensory basis of the human mind and so provide the greater wealth of material by which it attains levels beyond those of the animal mind, probably the most important are the functions of the fovea (macula lutea), of accommodation, and of static stereoscopy, and the development of a delicate skin and prehensile hands. All these make in some way for a differentiation that is the first form of abstraction. The fovea refines and distinguishes positions and forms, while accommodation sharpens the objects of attention and dissipates the rest; stereoscopy adds a new character to a group of forms that may persist for indefinite periods of observation; delicate skin gives greater sensitivity to variations of pressure, and the prehensile hand implies a very great refinement in the positions and forms of the derived articular sense. In the hand this becomes a fine mobile tridimensional sense that, like the stereoscopic eye, can go round and through things, so almost isolating them from their surroundings. At the same time the articular sense is the conscious correlate of action and of the individual's share in his experiences."

All these conditions that confer upon man the fuller vision to see and understand the world and interpret what is happening around him, can be shown to be the results directly or indirectly of those profound structural revolutions which have given man what is virtually a new instrument of sight and with it a new vision.

News and Views.

MR. G. H. WILKINS, accompanied by Mr. C. B. Eielson, left Point Barrow, Alaska, on April 15, and flew across the Arctic Ocean to Spitsbergen, a distance of 2200 miles, in a little over twenty hours. The machine used was a Lockheed monoplane with a 220 h.p. Wright Whirlwind engine. The start was difficult and a long snow runway on the frozen surface of a lagoon had to be made. For the first five hundred miles the weather was clear; then a hundred miles of cloud were passed through before the visibility again became good. In lat. 84° N. long. 75° W., dark clouds were seen to the north, described as possibly land clouds, but as this was near one of Peary's tracks, on which he reported no land, Mr. Wilkins did not turn aside. Landing-places on the pack-ice which might have served in an emergency were seen, but no attempt was made to descend, since the ascent would have been a very difficult matter. A view of Grant

Land (Ellesmere Island) was obtained, and then the course was set for Spitsbergen. This course, it will be noted, did not pass over the North Pole itself. Strong winds were experienced and there was much cloud to the south over Greenland, but one high peak of the land was seen. The temperature was then about -48° F.

WITHIN 200 miles of Spitsbergen, flying conditions became more trying. The open water and high temperature to the south were associated with great cloud masses, and even at 8000 ft. they could not be avoided. The difficulty was to locate King's Bay, Spitsbergen, where it was hoped to land. The weather became worse and flying was difficult. Petrol was running short when a landing was made on fast ice near the shore at the northern entrance to Ice Fjord. The reports mention an island, but there is