abstract geometry in such a way as to sacrifice some of the simplicity in the geometry to gain as much as possible in the physics. For example, in his theory there is no need of a gravitational force to make a planet move in its orbit, for this orbit is as natural in his geometry as is a straight line in the geometry of Euclid and Newton. This is what is meant by 'the geometrisation of physics', and we may define

physical geometry as that one of the many possible systems of abstract geometry which is most successful in giving a simple account of physical phenomena. The experience of draughtsmen and others shows that Euclidean geometry works very well indeed in ordinary terrestrial affairs, so physical geometry cannot differ very much from Euclidean.

The Origin of Adaptations.¹

By Dr. E. J. ALLEN, F.R.S.

BY an adaptation is meant nothing more than a character of an organism, which has enabled a species to survive itself as such, or to survive until it is transformed into another species. It is survival that gives the measure of the value of the adaptation. Survival can only occur if the whole organism is adapted to the environment to an extent that suffices. Organism and environment must be thought of as a unity, as interlocked and fitted closely to form that harmony which is Nature and life. Organic evolution is a phase—the crowning phase, may be—of cosmic evolution. The biological environment determines survival no less than the physical, and adaptation to both must be sufficient. The environment is not fixed, but must be thought of as in a condition of perpetual flux and change. This is true especially of the biological environment, for species once common may practically disappear, and years later may reappear abundantly with devastating effect on other organisms.

The general physical conditions under which organisms live have been well discussed by L. J. Henderson in his book "The Fitness of the Environment" (1913). Henderson discusses the unique properties of water, carbonic acid, hydrogen, and oxygen, and shows how they are specially fitted for the purposes of organic life. "There are no other compounds which share more than a small part of the qualities of fitness of water and carbonic acid; no other elements which share those of carbon, hydrogen, and oxygen." "None of the characteristics of these substances is known to be unfit or seriously inferior to the same characteristics in any other substance." "The fitness of the environment is one part of a reciprocal relationship of which the fitness of the organism is the other."

Darwin's answer to the question, how does the adaptation of organism to environment come to be, was based on three factors—heredity, variation, selection. In ultimate analysis the fact of heredity depends on the cellular structure of organisms and the phenomenon of cell division. When a living cell divides, its most essential substance, the germ plasm, separates into two portions which are almost equal. But we cannot so easily obtain an insight into the problem of variation. For simplicity's sake, consider first the formation of a germ cell from

its mother cell in an organism which is developing parthenogenetically. The researches of the colloid chemist have given us the picture. In imagination enlarge the germ mother-cell until you see the two phases; the liquid, the mass of molecular aggregates varied in size and shape; until you see the long, complex chains of atoms, building up the heavy molecules which form the aggregates; until you see the solar systems in miniature of protons and electrons which are the atoms—a seething, churning mass, active with the activity of cosmic forces, receiving matter and energy constantly from the surrounding medium, and giving them back. The preparations for cell-division begin; the molecular aggregates arrange themselves in new patterns; the separation of the cell into two parts ensues. Is it a matter for surprise that the partition of pattern and of substance is not always, perhaps is never, exact? We cannot wonder that germ cells thus produced differ in small respects among themselves. A few molecules more or less, a few atoms more or less, a few electrons even more or less, may mean large changes in the offspring into which the germ cell grows. We are, I think, safe in concluding that lack of equality in the partition of the hereditary material is one important cause of variation. If we think on similar lines of sexual development, where instead of one we have two germ cells uniting to form the zygote from which the offspring is developed, the probability of variation between parent and offspring, and between different offspring of the same parent, is obviously much increased.

Weismann was the first to draw a clear and sharp distinction between true hereditary characters and modifications of the body or soma, produced by the direct action of physical changes in the environment, and to develop the conception of the continuity of the germ-plasm. The germ-plasm is the transmitter, in unbroken continuity from generation to generation, of hereditary qualities. The body or soma is its temporary guardian, perishing when the work of transmission has been done. Blastogenic characters, as Weismann called the true hereditary characters, reappear in exactly the same form in the offspring as they show in the parent, provided both parent and offspring have grown up in the normal environment. Few now question that the nucleus is the essential organ of the germ cell which is engaged in the transmission of hereditary characters. Few

¹ Extracted from the Hooker Lecture, delivered before the Linnean Society of London on Mar. 14.

also question that the chromatin of the nucleus is the bearer of definite factors or genes, or that these factors are distributed in linear order in the chromosomes which appear at the time of cell-division. In ordinary normal development, hereditary characters are determined by the factors in the germ-plasm in response to stimuli furnished by the environment, for in the absence of a suitable environment no development at all takes place. If the environment is normal the characters are reproduced in normal form.

Hereditary *variations* are differences from the parental characters, which appear in the offspring, and are transmitted by the offspring to its descendants. We can only study them when the environmental conditions, in so far as they affect the characters concerned, remain unchanged throughout the growth of both parent and offspring, and this is the recognised basis of all breeding experiments. These hereditary or blastogenic variations we now call mutations, and mutations, according to the most recent usage of the word, may be either large or small, it being quite impossible to distinguish them from any other variations by the factor of size alone. In this respect the word mutation as now used does not convey exactly the same idea—it is not so limited—as Darwin's words 'sport' or 'monstrosity', and its meaning has been somewhat changed since it was first introduced by de Vries. The modern view is that mutations are heritable changes in the characters of organisms, which are due to definite alterations of the factors or genes, situated in the chromosomes of the germ cells. Contrasted with these mutations we have somatic modifications, the acquired characters of Weismann, the reaction of the organism to definite changes in the environment. De Vries's term 'fluctuations' is now generally employed in the same sense, and has, I think, ceased to be useful.

The variations or deviations revealed by the measurements of biometricians, which group themselves around a mean or modal value, according to the 'law of error', are probably in part small mutations which can be transmitted to descendants, and in part somatic modifications which are not so transmitted. With adequate measurements for a series of consecutive generations, the statistical tests which the biometrician applies enable him to say whether or not any of these deviations are inherited, and to give a measure of that inheritance. To take an example, it is frequently maintained that Johannsen's experiments with garden beans (*Phaseolus vulgaris nana*), which multiply by self-fertilisation and from which he obtained what he regards as pure lines, have proved that individual differences as shown by these lines are not inherited, and that therefore they cannot provide material upon which natural selection can act. Pearson, however, maintained, so long ago as 1910, that the pure line theory demands that the offspring shall be as highly correlated with the grandparent as it is with the parent, whereas Johannsen's own figures show that the coefficient of correlation between offspring and parent is

higher than that between offspring and grandparent. These experiments should be repeated with larger numbers of measurements.

Mutations may be classified into 'combination mutations', those due to rearrangement of factors or genes already present, and 'alteration mutations' due to changes in the factors themselves. Evidence is now forthcoming that the germ-plasm itself can be acted on by physical and chemical forces in the environment in such a way that mutations are produced. Heslop Harrison's work on the production of melanic forms in Geometrid moths was described (see NATURE, Jan. 22, 1927, p. 127). This work is of outstanding interest, not only on account of the fundamental importance of the results attained, but also for its perfect combination of acute and penetrating observations in the field with critical and long-sustained experimentation. Harrison has shown quite clearly that the germ-plasm can be changed by chemical substances contained in the food of an animal, or in more general terms that the germ-plasm can be altered by the environment. Another important advance in the same direction has come in H. J. Muller's account (*Science*, July 1927) of his production of mutations in *Drosophila* by irradiating spermatozoa or oöcytes with X-rays. When the correct dosage had been found, many mutations were produced, which on the whole were similar to those previously reported in *Drosophila*, such as 'white eye', 'miniature wing', and 'forked-bristles'. Most were recessive, but a number were dominant.

One further point with regard to variations must be noted. The possibilities of variation of an organism are strictly limited and circumscribed by the general physical and chemical properties of protoplasm. The essential physiological processes, upon which the life and activity of organisms depend, are comparatively few. Digestion, growth, sexual activity follow the same general lines throughout the whole animal kingdom. It is probable that the physico-chemical mechanisms alike of all muscular movement, of the movement of amœba by pseudopodia and of the movement of cilia, will fall into one general scheme. Similarly, the transmission of the nervous impulse is being shown to proceed on essentially the same lines in animals of widely separated groups. The essential physiological processes already function in the protista. If the physiological processes are few and circumscribed, variations in structure and form will be limited also, recognising that form is "a product of an inner physiological activity" (Kusnetzov. D'Arcy Thompson, "Growth and Form", 1917).

The last of the three principal factors on which Darwin based his theory of evolution was natural selection, or in Herbert Spencer's phrase, which Darwin adopted, "the Survival of the Fittest". Later, Ray Lankester suggested another formula, "the elimination of the unfit", which describes more correctly the meaning of the conception. That natural selection, acting on heritable variations, is a factor in fixing adaptations is almost a

truism. But whether it is the only factor, whether it is sufficient by itself to account for the living things we know, each fitting so perfectly its own little niche in its world, is a more difficult question. The process is of necessity so slow. But the time available is enormous, and geologists and physiologists seem satisfied that it is to be reckoned in tens, if not in hundreds of millions of years. We must consider also whether the mutations that occur are sufficiently diverse, for if adaptations are to be selected, mutations in the direction of those adaptations must occur. The known mutations of *Drosophila* amount to some 400, but the mutations so far studied are, for practical reasons, those which are large and obvious. There is increasing reason to think that they are outnumbered by small mutations which only long practice can detect. Many mutations studied are slight colour changes, because they can be distinguished with remarkable precision by the practised human eye. Correspondingly minute changes in size or shape would be very hard to detect, and to study them by breeding experiments and the methods of Mendelian analysis is not yet possible. We can only form judgments about them by analogy with results from larger mutations. There remains the statistical method of attack, by the study of mass-populations of successive generations. The method is efficient, active, and advancing, and it can only be lightly disregarded by those who have failed to grasp its meaning.

Alternative or additional theories to account for evolution are favoured by many naturalists. Darwin himself attached much importance to characters being inherited which had been produced by constant use or disuse in the parent. This is a particular case of Lamarck's conception of "the inheritance of acquired characters", or, better expressed, of somatic modifications. Some authorities consider that experimental proof of such inheritance is already available: for example, MacBride cites the work of Kammerer and of Durkhen and Brecher, the latter work being supported also by Heslop Harrison. On the other hand, Graham Kerr ("Evolution." London: Macmillan and Co., 1926) advances strong arguments on the other side, and Goodrich ("Living Organisms", 1924) takes the same view: "The real question Biology has to answer in future, as O. Hertwig has pointed out, is not 'Are modifications inherited?' but 'How are new factors acquired?'"

Even if it could be proved experimentally, without possibility of question, that somatic modifications were inherited, we should only have advanced a little way towards an understanding of our problem. The question *how* the soma influenced the factors in the germ cell would remain. In this connexion Cunningham's suggestion that hormones provided a capable instrument is of interest, and might be followed up experimentally.

The more elusive notions, which introduce the idea of some psychic or psychoid influence, controlling and regulating the processes of meta-

bolism and organic growth, it is hard to distinguish from the animisms of primitive man, who finds a spirit on every mountain, a devil in every bush. All these ideas contain a suggestion of purpose, some of them an idea of almost conscious purpose such as we know only in ourselves, or by analogy assume in higher animals, in each case associated with an elaborately differentiated nervous system. They are brought into the story at the point where knowledge based on observation and experiment ceases, at the point where it seems to many of us more satisfactory to say frankly, I do not know.

The idea of orthogenesis or nomogenesis (Berg, 1926), the idea that development takes place in a predetermined direction, is certainly unsatisfying in its elementary form. An explanation of adaptations on these lines offers special difficulty, for the theory fails to provide the flexibility necessary to produce that constant adjustment of the organism to its ever-changing environment which is imperatively demanded. If, to reach the required adjustment, a predetermined direction of variations and of evolution is postulated in the organism, a predetermined evolution of the environment on parallel lines would surely be necessary. That evolution proceeds according to laws of the same character as other laws of Nature, is the common basis of all modern evolutionary theory, and was held perhaps more strongly by Darwin, Huxley, and Weismann than it is by some writers of to-day. The physical laws in accordance with which the processes of growth are controlled, with results that we see in so many curious patterns, from the simple branching of a tree or of a nerve fibre to the elaborate spirals of a shell or of a growing plant; or again, the laws which lie behind the varied shapes, so curious and wonderful, of organisms and of their different parts,—these laws, and many others like them, still call for serious consideration and research. This is the valuable feature of the theory of orthogenesis, and in directing renewed attention to it, its followers make a valued contribution to biological thought.

There are many other aspects of the problem of the origin of adaptations that might be considered, but it has seemed better to confine ourselves to the larger questions, even at the risk of saying nothing but what was already well known. The outlook for biology to-day is as alluring and as full of hope as it was in those years of joyful enthusiasm which followed the historic paper by Darwin and Wallace, communicated to the Linnean Society by Hooker in 1858. In whatever direction we look problems bristle, problems open to successful attack; and the old qualities, insight, patience, and determination, will get them solved. But we must not limit the outlook, and all aspects of biological research must proceed hand in hand. Botany, zoology, palæontology, the work of the systematist and of the field naturalist, the study of structure and the study of function, the work of the embryologist and of the experimental physiologist, of the geneticist and of the statistician, all are necessary, and none can succeed without the others.