

Imaging Brain Phospholipase A₂-Mediated Signal Transduction in Response to Acute Fluoxetine Administration in Unanesthetized Rats

Ying Qu^{*1}, Lisa Chang¹, Justin Klaff¹, Ruth Seemann¹, and Stanley I Rapoport¹

¹Brain Physiology and Metabolism Section, National Institute on Aging, National Institutes of Health, Bethesda, MD 20892, USA

Fluoxetine, a selective serotonin (5-hydroxytryptamine, 5-HT) reuptake inhibitor, is used widely to treat depression and related disorders. By inhibiting presynaptic 5-HT reuptake, fluoxetine is thought to act by increasing 5-HT in the synaptic cleft, thus 5-HT binding to postsynaptic 5-HT_{2A/2C} receptors. These receptors can be coupled via a G-protein to phospholipase A₂ (PLA₂), which when activated releases the second messenger arachidonic acid from synaptic membrane phospholipids. To image this activation, fluoxetine (10 mg/kg) or saline vehicle was administered i.p. to unanesthetized rats, and regional brain incorporation coefficients k^* of intravenously injected radiolabeled arachidonic acid were measured after 30 min. Compared with vehicle, fluoxetine significantly increased k^* in prefrontal, motor, somatosensory, and olfactory cortex, as well as in the basal ganglia, hippocampus, and thalamus. Many of these regions demonstrate high densities of the serotonin reuptake transporter and of 5-HT_{2A/2C} receptors. Brain stem, spinal cord, and cerebellum, which showed no significant response to fluoxetine, have low densities of the transporters and receptors. The results show that it is possible to image quantitatively PLA₂-mediated signal transduction *in vivo* in response to fluoxetine.

Neuropsychopharmacology (2003) 28, 1219–1226, advance online publication, 7 May 2003; doi:10.1038/sj.npp.1300177

Keywords: serotonin; phospholipase A₂; arachidonic acid; fluoxetine; signal transduction; reuptake; imaging

INTRODUCTION

Fluoxetine is widely used to treat depression, obsessive-compulsive disorder, panic disorder, and bulimia (Fuller, 1995). It is a selective serotonin (5-hydroxytryptamine, 5-HT) reuptake inhibitor, enhancing postsynaptic serotonergic neurotransmission by increasing the concentration of 5-HT in the synaptic cleft (Fuller and Wong, 1977; Wong *et al*, 1995).

Acute administration of fluoxetine to rodents elicits anxiogenic effects in social interaction and maze tests (Kurt *et al*, 2000; To *et al*, 1999), potentiates defensive reactions such as flight and biting (Griebel *et al*, 1995), and potentiates amphetamine-induced locomotor activity (Sills *et al*, 1999). Acute drug also stimulates secretion of pituitary ACTH and adrenal corticosterone (Dinan, 1996; Li *et al*, 1993). It triggers transcription of the gene encoding the corticotrophin-release factor and its type 1 receptor (Torres

et al, 1998), thus affecting the hypothalamic–pituitary–adrenal axis that is considered hyperactive in depressed patients (Leonard, 2001).

Acute administration of fluoxetine also exerts inhibitory effects. It reduces spontaneous motor activity in 6- to 8-month-old rats, and to a lesser extent in aged rats (Stanford *et al*, 2002). It reduces regional cerebral metabolic rates for glucose (rCMR_{glc}) in awake rats (Freo *et al*, 2000) as well as in humans (Cook *et al*, 1994). Similar reductions in rCMR_{glc} in rats have been noted in response to the 5-HT_{2A/2C} receptor agonist (\pm)-2,5-dimethoxy-4-iodophenyl-2-aminopropane (DOI) (Freo *et al*, 1991).

5-HT_{2A/2C} receptors can be coupled via G-proteins to phospholipase A₂ (PLA₂) (Berg *et al*, 1998; Felder *et al*, 1990; Kim *et al*, 1999; Qu *et al*, in press), which when activated will release arachidonic acid (20:4 n-6) from the stereospecifically numbered (*sn*)-2 position of phospholipids (Axelrod, 1990). Both arachidonate and its eicosanoid metabolites are important second messengers (Fitzpatrick and Soberman, 2001). Much of the released arachidonate is not metabolized to eicosanoids, however, but rapidly re-esterified into synaptic membrane phospholipids, together with arachidonate derived from the plasma. The re-esterification process can be imaged in unanesthetized rats by injecting radiolabeled arachidonic acid intravenously following drug, and measuring its regional brain incorporation with quantitative autoradiography (Chang *et al*, 1997;

*Correspondence: Dr Y Qu, Present address: Neuroscience Johnson & Johnson Pharmaceutical Research and Development, 3210 Merryfield Row, San Diego, CA 92121, USA, Tel: +1 858 320 3418, Fax: 858 450 2040, E-mail: yqu1@prdujnj.com

Received 16 August 2002; revised 27 January 2003; accepted 28 January 2003

Online publication: 3 February 2003 at <http://www.acnp.org/citations/Npp020303314/default.pdf>

DeGeorge *et al*, 1991; Jones *et al*, 1996; Rapoport, 2001; Robinson *et al*, 1992).

In this regard, we reported that radiolabeled arachidonate could be used to image brain PLA₂ activation in rats administered the 5-HT_{2A/2C} receptor agonist, DOI (Qu *et al*, 2001a). Significant increases in tracer incorporation, noted in brain areas with high densities of 5-HT_{2A/2C} receptors, could be blocked by pretreatment with the 5-HT₂ antagonist mianserin. The increases were considered to follow the binding of DOI to 5-HT_{2A/2C} receptors, and activation of the PLA₂ to which these receptors are coupled.

We thought it of interest to see if fluoxetine also would increase incorporation of labeled arachidonate into the brain, as might be expected from its ability to increase 5-HT in the synaptic cleft and thus 5-HT occupancy of 5-HT_{2A/2C} receptors (see above). To test this expectation, we decided to measure the incorporation of radiolabeled arachidonate from plasma into the brain of unanesthetized rats, 30 min after giving fluoxetine 10 mg/kg i.p. At 30–60 min, this dose increases 5-HT levels in the rat brain, and the levels remain elevated for 3 h (Guan and McBride, 1988). Additionally, after 30 min, fluoxetine at doses of 4 mg/kg and 40 mg/kg i.p., respectively, has been reported to reduce rCMR_{glc} in 28 of 66 regions (mean global decrease was 23%) and 37 of 66 (mean global decrease was 33%) regions examined (Freo *et al*, 2000).

MATERIALS AND METHODS

Materials

Radiolabeled [5,6,8,9,11,12,14,15-³H]arachidonate ([³H]arachidonate) at a specific activity of 200 Ci/mmol was purchased from Moravsek Biochemicals (Brea, CA). Radiochemical purity as shown by thin-layer chromatography always exceeded 96%. Fluoxetine was purchased from Sigma-Research Biochemicals International (Natick, MA). Sodium pentobarbital was purchased from Richmond Veterinary Supply (Richmond, VA).

Animals

Male Fischer-344 rats (Charles River Laboratories, Wilmington, MA), weighing 290–320 g and 12-weeks old, were housed under standard laboratory conditions with a 12 h light–12 h dark cycle, with ready access to laboratory chow and water. The experimental protocol was approved by the National Institute of Child Health and Human Development Animal Care and Use Committee, and conformed to the Guide for the Care and Use of Laboratory Animals (National Institute of Health Publication 86-23). Rats were divided into two groups, eight animals in each: (1) those administered fluoxetine, 10 mg/kg i.p. in 1 ml saline, 30 min before tracer infusion and (2) controls administered 1 ml i.p. saline 30 min before tracer infusion.

Arterial and Venous Catheterization

The *in vivo* fatty acid method to image brain PLA₂ signaling has been described elsewhere (DeGeorge *et al*, 1991; Hayakawa *et al*, 2001; Qu *et al*, 2003). Briefly, rats in each of the two groups were anesthetized with halothane

(1–3% v/v in O₂). PE 50 polyethylene catheters (Clay Adams, Lincolnshire, IL) filled with heparinized saline (100 IU/ml) were surgically implanted into a femoral artery and vein, after which the incision site was infiltrated with a local anesthetic (lidocaine) and closed with wound clips. The rats were wrapped loosely in a fast-setting plaster cast, secured to a wooden block with the upper body free, and allowed to recover from anesthesia in a temperature-controlled and sound-dampened box for 4 h. Body temperature was kept at 36–37°C using a rectal thermometer and a feedback heating device.

Drug Administration and Tracer Infusion

After a rat was allowed to recover from anesthesia for 4 h, 125 µl arterial blood was withdrawn to measure pH, pO₂, and pCO₂. At 30 min after the administration of i.p. saline or fluoxetine (see above), 1.75 mCi/kg [³H]arachidonate in 2 ml of 5 mM HEPES buffer, pH 7.4, containing 50 mg/ml fatty acid-free bovine serum, was infused through the venous canula with an infusion pump (Harvard Instruments, Holliston, MA), at a rate of 400 µl/min for 5 min. Timed 125-µl arterial blood samples were collected from the beginning of infusion to 20 min, when the rats were killed with 65 mg i.v. sodium pentobarbital. The brain was immediately removed and frozen in 2-methylbutane at –70°C for subsequent quantitative autoradiography. Plasma was separated from arterial blood by centrifugation, and its lipids were extracted by the method of Folch (Folch *et al*, 1957). Radioactivity in the organic fraction was measured by liquid scintillation counting.

Autoradiography and Calculations

Frozen brains were sectioned on a cryostat at –20°C. Sets of three adjacent 20-µm sections were collected and mounted on glass coverslips at 140 µm coronal intervals and dried. The three sections were exposed together with [³H]methylmethacrylate autoradiographic standards (Amersham, Arlington Heights, IL) to [³H]phosphor imaging plates (Fuji Medical Systems, Stamford, CT) for 7 days, which were then scanned by a BAS 5000 scanner system (West Lafayette, IN) following the manufacturer's instructions. An adjacent section was collected and stained with cresyl violet to identify brain regions using a rat-brain atlas (Paxinos and Watson, 1987).

Regional brain radioactivity was measured in sextuplicate by quantitative densitometry using phosphor-imaging software (Image Gauge V3.45, Fuji). Regional brain incorporation coefficients k^* were calculated as

$$k^* = \frac{c_{\text{brain}}^*(20 \text{ min})}{\int_0^{20} c_{\text{plasma}}^* dt} \quad (1)$$

where k^* is in units of ml/s/g; $c_{\text{brain}}^*(20 \text{ min})$ is brain radioactivity at 20 min after the onset of infusion, in units of nCi/g; c_{plasma}^* is plasma fatty acid radioactivity in units of nCi per ml; and t is the time after the onset of [³H]arachidonate infusion.

Data were compared statistically using Prism software for the Macintosh (Abacus Concepts, Berkeley, CA) and are reported as means ± SEM. Student's *t*-test tests were used

to evaluate statistical significance between experimental and control means. $P \leq 0.05$ was taken as indicating statistical significance.

RESULTS

Table 1 summarizes the mean physiological parameters in unanesthetized control and fluoxetine-treated rats. The control means are similar to published values (Qu *et al*, 2003) and were not changed significantly by drug.

Figure 1 presents representative autoradiographs of coronal brain sections from a rat administered fluoxetine *i.p.* and a control rat. Fluoxetine 10 mg/kg *i.p.* caused significant widespread increases in k^* for [^3H]arachidonate. The mean regional incorporation coefficients k^* corresponding to such increases are presented in Table 2.

As illustrated by Table 2, 30 min after fluoxetine administration, 42 of 85 brain regions had significantly increased mean values for k^* (by about 30%) compared with control values. Significant increases were evident in prefrontal frontal, motor and somatosensory cortex, and olfactory and pyriform cortex, but not in the auditory or visual cortex. Regions in the caudate-putamen, nucleus accumbens, globus pallidus, amygdala, septum, thalamus, and hypothalamus also showed significant increments in k^* , whereas significant changes generally were absent in regions of the brain stem, spinal cord, and cerebellum. The hippocampus showed elevations in CA1, CA2, and CA3 areas, but not in the dentate gyrus. The choroid plexus, where the control value of k^* was much greater than in the brain parenchyma, demonstrated a 40% increase in k^* in response to fluoxetine. White matter incorporation of [^3H]arachidonate was unaffected by fluoxetine.

DISCUSSION

At 30 min following administration to unanesthetized adult male rats of fluoxetine (10 mg/kg *i.p.*), the incorporation coefficient k^* for [^3H]arachidonate was increased significantly compared with control in 45 of 85 brain regions that were examined. Significant increases were noted in regions of the prefrontal and frontal cortex, motor and somatosensory cortex, and of the basal ganglia, septum, hippocampus, thalamus, and hypothalamus, but not of the auditory or visual cortex, white matter, brain stem, spinal cord, or cerebellum (Table 2) (Appel *et al*, 1990; Li *et al*, 2001; Pazos and Palacios, 1985).

Table 1 Physiological Parameters of Rats after Surgery

	Saline	Acute fluoxetine
Body temperature ^a (°C)	36.1 ± 0.3 ^b	36.1 ± 0.3
Arterial blood pressure (mm Hg)	125 ± 2/78 ± 1	129 ± 5/74 ± 3
(systolic pressure/diastolic pressure)		
Heart rate (beats/min)	416 ± 4	413 ± 13
Arterial pH	7.41 ± 0.02	7.41 ± 0.02
Arterial blood gas (pCO ₂)	41.9 ± 1.3	39.6 ± 2.0
Arterial blood gas (pO ₂)	99.8 ± 5.0	101 ± 3.0

^aTemperature was measured with a rectal thermoprobe.

^bMean ± SEM.

Animal number = 8.




Figure 1 Coronal autoradiographs demonstrating [^3H]arachidonate incorporation coefficients k^* from brains of (a) control rat and (b) rat acutely administered fluoxetine (10 mg/kg *i.p.*). k^* is color coded.

Fluoxetine, a high-affinity, high-selectivity antagonist of the serotonin reuptake transporter (Wong *et al*, 1991) promotes accumulation of 5-HT in the synaptic cleft. Acutely administered fluoxetine has been reported to increase extracellular 5-HT in many brain areas, including the striatum (Perry and Fuller, 1993; Rutter and Auerbach, 1993), thalamus (Dailey *et al*, 1992), diencephalon (Rutter *et al*, 1993), hypothalamus (Perry *et al*, 1993), and nucleus accumbens. Increases were evident within 30–60 min and last for up to 3 h (Guan *et al*, 1988). In light of our observations that DOI increases k^* for [^3H]arachidonate and that the increases can be blocked by a 5-HT₂ antagonist (Qu *et al*, 2003), our results with fluoxetine (Table 2) suggest that increased synaptic cleft 5-HT caused by the drug indirectly increased the 5-HT occupancy of 5-HT_{2A/2C} receptors, which are coupled to PLA₂, so as to release arachidonic acid (Berg *et al*, 1998; Felder *et al*, 1990; Kim *et al*, 1999). The regional incorporation coefficient k^* for [^3H]arachidonate reflects this release (see Introduction) (Rapoport, 2001; Robinson *et al*, 1992).

While many brain areas with reported high densities of the serotonin reuptake transporter and of 5-HT_{2A/2C}

Table 2 Regional [³H]Arachidonic Acid Incorporation Coefficients k^* (ml/s/g brain $\times 10^4$) in Rat Brain in Response to Acute Fluoxetine

Brain region ^a	Saline control	Fluoxetine	Brain region ^a	Saline control	Fluoxetine
<i>Cerebral cortex</i>			<i>Thalamus and related areas</i>		
Prefrontal cortex IV	9.1 \pm 0.3	12.0 \pm 1.2*	Paratenial nuclei	8.2 \pm 0.3	11.2 \pm 1.1*
Frontal cortex layer II–III	9.5 \pm 0.5	12.0 \pm 1.0*	Anteroventral nuclei	10.2 \pm 0.4	13.0 \pm 0.8**
Frontal cortex layer IV	10.6 \pm 0.5	13.5 \pm 1.1*	Anteromedial nuclei	8.5 \pm 0.3	10.9 \pm 0.7*
Motor cortex layer II–III	9.1 \pm 0.4	12.1 \pm 1.2*	Reticular nuclei	8.4 \pm 0.5	10.8 \pm 0.9*
Motor cortex layer IV	10.5 \pm 0.5	14.1 \pm 1.3*	Paraventricular nuclei	7.3 \pm 0.6	10.0 \pm 0.6**
Motor cortex layer V–VI	8.9 \pm 0.4	12.3 \pm 1.0*	Ventroposterior medial nucleus	7.9 \pm 0.3	10.3 \pm 0.8*
Somatosensory cortex layer II–III	9.5 \pm 0.5	12.7 \pm 1.2*	Ventroposterior lateral	7.7 \pm 0.3	9.6 \pm 0.8
Somatosensory cortex layer IV	10.2 \pm 0.5	14.0 \pm 1.4*	Lateral habenular nucleus	8.6 \pm 0.3	11.9 \pm 1.2*
Somatosensory cortex layer V–VI	8.9 \pm 0.4	11.9 \pm 1.0*	Medial habenular nucleus	9.3 \pm 0.4	12.5 \pm 1.2*
Anterior cingulate cortex	9.8 \pm 0.4	11.8 \pm 1.4	Medial geniculate nucleus	10.4 \pm 1.0	11.3 \pm 1.2
Auditory cortex layer II–III	10.3 \pm 1.2	11.6 \pm 1.4	Dorsolateral geniculate nucleus	8.4 \pm 0.4	10.9 \pm 0.8*
Auditory cortex layer IV	11.9 \pm 1.5	13.1 \pm 1.2	Parafascicular nucleus	7.84 \pm 0.30	10.3 \pm 0.75*
Auditory cortex layer V–VI	9.6 \pm 1.1	10.1 \pm 0.8	Inferior colliculus	11.9 \pm 0.97	12.9 \pm 0.99
Visual cortex layer II–III	10.0 \pm 1.0	11.6 \pm 1.1	Superior colliculus	11.7 \pm 1.9	12.4 \pm 1.5
Visual cortex layer IV	10.7 \pm 1.0	12.3 \pm 1.1			
Visual cortex layer V–VI	10.0 \pm 0.9	13.6 \pm 2.4			
<i>White matter</i>			<i>Hypothalamus</i>		
Corpus callosum	4.4 \pm 0.4	5.4 \pm 0.3	Supraoptic nucleus	13.0 \pm 2.5	16.1 \pm 2.8
Internal capsule	4.2 \pm 0.4	5.2 \pm 0.3	Subfornical organ	9.1 \pm 0.6	12.9 \pm 1.3*
Anterior commissure	5.3 \pm 0.5	6.5 \pm 0.4	Lateral nuclei	6.7 \pm 0.5	8.7 \pm 0.7*
			Anterior nuclei	6.8 \pm 0.8	9.3 \pm 0.9*
			Periventricular nucleus	8.0 \pm 0.5	9.2 \pm 0.8
<i>Olfactory system</i>			Arcuate nucleus	6.8 \pm 0.4	8.2 \pm 0.8*
Olfactory cortex	11.1 \pm 0.5	14.6 \pm 1.1**	Ventromedial nucleus	6.5 \pm 0.4	8.9 \pm 0.9*
Pyramidal cortex	9.7 \pm 0.4	14.5 \pm 1.2**	Posterior nucleus	7.6 \pm 0.3	10.1 \pm 0.8*
			Medial forebrain bundle	6.8 \pm 0.4	9.2 \pm 0.8
<i>Basal ganglia and related areas</i>			Mammillary body	7.4 \pm 0.4	10.7 \pm 1.0*
Nucleus accumbens	8.1 \pm 0.4	10.9 \pm 0.9*	Median eminence	14.0 \pm 2.5	13.5 \pm 2.6
Caudate-putamen dorsal	8.1 \pm 0.3	10.5 \pm 0.7*			
Caudate-putamen ventral	8.3 \pm 0.3	10.4 \pm 0.9			
Caudate-putamen lateral	8.0 \pm 0.4	10.6 \pm 0.8*	<i>Brainstem and spinal cord</i>		
Caudate-putamen medial	7.9 \pm 0.3	9.5 \pm 0.8	Raphe magnus nuclei	7.5 \pm 0.6	8.9 \pm 1.2
Bed nucleus stria preoptic nucl.	7.4 \pm 0.3	9.7 \pm 0.7*	Raphe pallidus nuclei	8.5 \pm 0.7	9.6 \pm 1.4
Suprachiasmatic nucleus	7.7 \pm 0.3	10.9 \pm 0.6*	Raphe median nuclei	8.5 \pm 0.8	10.2 \pm 1.2
Bed nucleus stria terminalis	6.6 \pm 0.4	8.6 \pm 0.7*	Raphe dorsal nuclei	9.1 \pm 0.7	10.4 \pm 0.8
Entopeduncular nucleus	6.2 \pm 0.5	7.9 \pm 0.7	Locus coeruleus	9.9 \pm 1.0	10.5 \pm 0.7
Globus pallidus	6.2 \pm 0.5	8.0 \pm 0.7*	Cochlear nucleus	14.0 \pm 2.5	16.7 \pm 3.8
Subthalamic nucleus	7.5 \pm 0.4	9.8 \pm 0.8*	Vestibular nucleus (medial)	11.7 \pm 0.8	13.8 \pm 1.2
Amygdala basolateral/basomedial nucl.	6.6 \pm 0.3	9.2 \pm 1.0*	Pretectal area	12.6 \pm 1.5	15.1 \pm 2.0
Substantia nigra			Pedunculopontine nucleus	7.2 \pm 0.7	8.2 \pm 0.8
Pars reticulata	8.5 \pm 0.8	9.6 \pm 0.7	Deep layers of superior colliculus	13.0 \pm 1.8	14.8 \pm 1.6
Pars compacta	8.2 \pm 0.8	7.9 \pm 0.9	Interpeduncular nucleus	10.3 \pm 0.8	18.0 \pm 6.1
<i>Septum</i>			Spinal Tract V nucleus	8.3 \pm 0.8	8.8 \pm 0.7
Lateral septal nucleus	6.5 \pm 0.3	8.1 \pm 0.7			
Medial septal nucleus	7.3 \pm 0.4	10.0 \pm 0.8*	<i>Cerebellum</i>		
Dorsal diagonal band	7.6 \pm 0.5	9.9 \pm 0.8*	Cerebellar gray matter	10.4 \pm 0.9	11.5 \pm 0.8
Ventral diagonal band	7.6 \pm 0.4	9.9 \pm 0.7*	Molecular layer, gray matter	10.7 \pm 0.9	11.9 \pm 0.8
			Granular layer, gray matter	11.9 \pm 0.7	14.3 \pm 1.2
<i>Hippocampal formation</i>			Flocculus	10.3 \pm 0.7	12.3 \pm 1.3
Ammon's horn CA1	9.2 \pm 0.3	12.1 \pm 1.0*	Cerebellar white matter	5.3 \pm 0.6	5.5 \pm 0.3
Ammon's horn CA2	8.5 \pm 0.4	12.3 \pm 1.3*			
Ammon's horn CA3	8.7 \pm 0.5	12.3 \pm 1.0**			
Dentate gyrus	9.9 \pm 1.2	11.6 \pm 1.2	<i>Choroid plexus</i>		
				46.9 \pm 3.1	66.6 \pm 5.3**

k^* values are means \pm SEM ($n=8$). k^* in rats given fluoxetine (10 mg/kg i.p.) was compared with k^* in rats given i.p. saline (controls).

Mean significantly different from control mean, * $p < 0.05$; ** $p < 0.01$

^aFrom Paxinos (1987).

receptors demonstrated significant increases in k^* for [³H]arachidonate in response to fluoxetine (Table 2), exceptions were evident. High densities of the serotonin reuptake transporter have been noted in the olfactory tubercle, lateral septal nucleus, olfactory tubercle, lateral septal nucleus, hypothalamic and thalamic nuclei, globus pallidus, central gray, superior colliculus, substantia nigra, interpeduncular nucleus, dorsal and lateral raphé, locus

coeruleus. Lesser but nevertheless high densities are present in the frontal cortex, caudate-putamen, ventral pallidum, and hippocampus (Choi *et al*, 2000; De Souza and Kuyatt, 1987; McReynolds and Meyer, 1998; Moll *et al*, 2000). k^* was elevated significantly in many of these regions, but not in all of them (eg lateral septal nucleus, superior colliculus, median, and dorsal raphé). High 5-HT₂ receptor densities are reported in the cerebral cortex (particularly layer IV),

olfactory and pyriform cortex, nucleus accumbens, caudate-putamen body and tail (Appel *et al*, 1990; Li *et al*, 2001; Pazos and Palacios, 1985). Frontal and motor cortical regions have higher densities than other cortical regions. Densities are not as high in the caudate-putamen head, globus pallidus, red nuclei, septal nuclei, and most parts of the hippocampus, thalamus, and hypothalamus. Spinal cord, brain stem, and cerebellum have few transporter or 5-HT_{2A/2C} receptor sites, and as expected k^* was not elevated significantly in these areas.

Increases in k^* may not occur at all brain sites containing the serotonin reuptake transporter because transporter sites and postsynaptic 5-HT_{2A/2C} receptors are not always colocalized. The transporter can be found at axons distant from 5-HT₂ synapses (Zhou *et al*, 1998), and some 5-HT₂ synapses may lack the transporter altogether (Brown and Molliver, 2000). Additionally, fluoxetine can increase brain dopamine levels (Yoshino *et al*, 2002) to activate PLA₂ at dopaminergic D₂ receptor sites rather than at 5-HT_{2A/2C} receptor sites (Hayakawa *et al*, 2001; Vial and Piomelli, 1995). Finally, fluoxetine can noncompetitively inhibit cholinergic muscarinic (Owens *et al*, 1997) and nicotinic receptors (Fryer and Lukas, 1999) and thereby reduce release of arachidonic acid.

Indeed, PLA₂ can be activated via a G-protein when an agonist binds to cholinergic muscarinic receptors and dopaminergic D₂ receptors as well as to 5-HT_{2A/2C} receptors (Bayon *et al*, 1997; Cooper *et al*, 1996; DeGeorge *et al*, 1991; Felder *et al*, 1990; Hayakawa *et al*, 2001; Kim *et al*, 1999; Vial *et al*, 1995). PLA₂ also may be activated by Ca²⁺ entry into cells when glutamate acts at NMDA receptors or acetylcholine acts at nicotinic receptors. Thus, baseline k^* for [³H]arachidonate in control rats represents the sum of baseline activation of PLA₂ via these different receptor subtypes, as well as incorporation because of membrane synthesis (particularly relevant for white matter myelin) (Rapoport *et al*, 1997).

A 10-fold higher baseline value of k^* for [³H]arachidonate in the choroid plexus, as well as a significant increment in k^* in response to DOI or fluoxetine (Table 2) are consistent with high densities of 5-HT_{2C}-binding sites in the choroid plexus (Kaufman *et al*, 1995; Li *et al*, 2001; Qu *et al*, 2003) and with activation of PLA₂ via these receptors. Fluoxetine has been reported to bind to 5-HT_{2C} sites in the choroid plexus (Palvimaki *et al*, 1996). The high baseline k^* in the choroid plexus is consistent with high rates of uptake of radiolabeled fatty acids into median eminence, subfornical organ, pineal gland, adenohypophysis, and neurohypophysis (Noronha *et al*, 1990; Noronha *et al*, 1989; Qu *et al*, 2003). These regions lack the continuous capillary bed of the blood-brain barrier (Rapoport, 1976), allowing access of labeled fatty acid both in its unbound unesterified form (as for the brain parenchyma), and when bound to albumin (Robinson *et al*, 1992). High values of k^* for labeled arachidonate also have been noted in the heart (Murphy *et al*, 2000).

Fluoxetine can induce hyperthermia in rats (Lin *et al*, 1998). The activity of PLA₂ as well that of other enzymes might be expected to increase with an increase in temperature (Bell *et al*, 1996). However, it is unlikely that hyperthermia accounted for the findings in this paper, as increased values for k^* were not found in regions having

low densities of the serotonin reuptake transporter, and because body temperature was maintained between 36 and 37°C in this study (see Materials and methods). Additionally hyperthermia has been shown to produce widespread increases in rCMR_{glc} in awake rats (Mickley *et al*, 1997), whereas fluoxetine, when administered under the same conditions in our study, has been shown either to not change or to reduce rCMR_{glc} (Freo *et al*, 2000).

Serotonin release from presynaptic elements can be controlled by two types of autoreceptors (Cerrito and Raiteri, 1980). 5-HT_{1A} autoreceptors are found at the somatodendritic region of cell bodies in midline and raphe nuclei in the pons and upper brain stem, whereas 5-HT_{1B} autoreceptors are located at axon terminals, where they control local 5-HT synthesis and release (Cooper *et al*, 1996; Hervas *et al*, 2000; Martin and Sanders-Bush, 1982). Acutely administered fluoxetine has been reported to inhibit electrical activity of serotonin neurons by activating somatodendritic 5-HT_{1A} autoreceptors (Czachura and Rasmussen, 2000; Fuller, 1995), thus reducing 5-HT released into the synaptic cleft. With chronic fluoxetine, inhibition of 5-HT unit activity is less because of desensitization of the 5-HT_{1A} autoreceptor. Desensitization and recovery of firing of 5-HT neurons develop slowly and concurrently, and may contribute to the delayed onset of therapeutic efficacy (Bergqvist *et al*, 1999; Blier and De Montigny, 1983).

The widespread increments in k^* for [³H]arachidonate in response to acute fluoxetine (Table 2) contrast with reports of no change or decrements in rCMR_{glc} under comparable experimental conditions (Freo *et al*, 2000). A similar discrepancy between increased radiolabeled arachidonate incorporation but decreased or no change in rCMR_{glc} follows DOI or methiothepin administration to unanesthetized rats (Freo *et al*, 1991; Qu *et al*, 2001b; Ricchieri *et al*, 1987). These discrepancies likely arise because [³H]arachidonate incorporation localizes the postsynaptic PLA₂-mediated release of arachidonic acid at the serotonergic neuron, whereas rCMR_{glc} represents ATP consumption by the downstream firing of presynaptic axon terminals of the neuron (Ashby *et al*, 1990; Purdon and Rapoport, 1998; Qu *et al*, 2003; Sokoloff, 1999).

The frontal cortex and hippocampus have been implicated as sites of action of antidepressant drugs (Duman *et al*, 1997; Jacobson and Sapolsky, 1991). In this study (Table 2), incorporation coefficients k^* for [³H]arachidonate were increased by fluoxetine in these regions, suggesting that to signaling via PLA₂ at these sites contributes to fluoxetine's antidepressant action. In major depression, serotonergic neurotransmission may be disturbed and processes of cellular immunity may be activated, suggesting a relation between the two processes. Indeed, there appears to be a constant 'crosstalk' between the immune endocrine, central, and peripheral nervous systems which can involve serotonin and arachidonic acid metabolites (eg prostaglandins) (Leonard, 2001; Maier and Watkins, 1998).

In conclusion, we have demonstrated in unanesthetized rats that acute fluoxetine activates PLA₂ signaling and increases labeled arachidonate incorporation from plasma into brain regions having high serotonin reuptake transporter densities, with some exceptions. Increased tracer incorporation likely arises because fluoxetine indirectly

increases binding of 5-HT to PLA₂-coupled 5-HT_{2A/2C} receptors. Identifying the exact mechanisms of fluoxetine's acute action will entail using specific receptor antagonists with it (Hayakawa *et al*, 2001; Mazzola-Pomietto *et al*, 1997; Rabiner *et al*, 2002), or studying 5-HT_{2A} or 5-HT_{2C} knockout mice (Lira *et al*, 2001).

REFERENCES

- Appel NM, Mitchell WM, Gaslick RK, Glennon RA, Teitler M, De Souza EB (1990). Autoradiographic characterization of (+)-1-(2,5-dimethoxy-4-[125I]iodophenyl)-2-aminopropane ([125I]DOI) binding to 5-HT₂ and 5-HT_{1C} receptors in rat brain. *J Pharmacol Exp Ther* 255: 843–857.
- Ashby Jr CR, Jiang LH, Kasser RJ, Wang RY (1990). Electrophysiological characterization of 5-hydroxytryptamine₂ receptors in the rat medial prefrontal cortex. *J Pharmacol Exp Ther* 252: 171–178.
- Axelrod J (1990). Receptor-mediated activation of phospholipase A₂ and arachidonic acid release in signal transduction. *Biochem Soc Trans* 18: 503–507.
- Bayon Y, Hernandez M, Alonso A, Nunez L, Garcia-Sancho J, Leslie C, Sanchez Crespo M, Nieto ML (1997). Cytosolic phospholipase A₂ is coupled to muscarinic receptors in the human astrocytoma cell line 1321N1: characterization of the transducing mechanism. *Biochem J* 323: 281–287.
- Bell JD, Burnside M, Owen JA, Royall ML, Baker ML (1996). Relationships between bilayer structure and phospholipase A₂ activity: interactions among temperature, diacylglycerol, lysolecithin, palmitic acid, and dipalmitoylphosphatidylcholine. *Biochemistry* 35: 4945–4955.
- Berg KA, Maayani S, Goldfarb J, Scaramellini C, Leff P, Clark WP (1998). Effector pathway-dependant relative efficacy at serotonin type 2A and 2C receptors: Evidence for agonist-directed trafficking of receptor stimulus. *Mol Pharmacol* 54: 94–104.
- Bergqvist PB, Bouchard C, Blier P (1999). Effect of long-term administration of antidepressant treatments on serotonin release in brain regions involved in obsessive-compulsive disorder. *Biol Psychiatry* 45: 164–174.
- Blier P, De Montigny C (1983). Electrophysiological investigations on the effect of repeated zimelidine administration on serotonergic neurotransmission in the rat. *J Neurosci* 3: 1270–1278.
- Brown P, Molliver ME (2000). Dual serotonin (5-HT) projections to the nucleus accumbens core and shell: relation of the 5-HT transporter to amphetamine-induced neurotoxicity. *J Neurosci* 20: 1952–1963.
- Cerrito F, Raiteri M (1980). Presynaptic autoreceptors control serotonin release from central nerve endings. *Pharmacol Res Commun* 12: 593–597.
- Chang MCJ, Arai T, Freed LM, Wakabayashi S, Channing MA, Dunn BB, Der MG, Bell JM, Sasaki T, Herscovitch P, Eckelman WC, Rapoport SI (1997). Brain incorporation of [1-¹¹C]-arachidonate in normocapnic and hypercapnic monkeys, measured with positron emission tomography. *Brain Res* 755: 74–83.
- Choi SR, Hou C, Oya S, Mu M, Kung MP, Siciliano M, Acton PD, Kung HF (2000). Selective *in vitro* and *in vivo* binding of [(125I)]ADAM to serotonin transporters in rat brain. *Synapse* 38: 403–412.
- Cook Jr EH, Metz J, Leventhal BL, Lebovitz M, Nathan M, Semerdjian SA, Brown T, Cooper MD (1994). Fluoxetine effects on cerebral glucose metabolism. *Neuroreport* 5: 1745–1748.
- Cooper JR, Bloom FE, Roth RH (1996). *The Biochemical Basis of Neuropharmacology*, 7th edn. Oxford University Press: Oxford.
- Czachura JF, Rasmussen K (2000). Effects of acute and chronic administration of fluoxetine on the activity of serotonergic neurons in the dorsal raphe nucleus of the rat. *Naunyn-Schmiedeberg's Arch Pharmacol* 362: 266–275.
- Dailey JW, Yan QS, Mishra PK, Burger RL, Jobe PC (1992). Effects of fluoxetine on convulsions and on brain serotonin as detected by microdialysis in genetically epilepsy-prone rats. *J Pharmacol Exp Ther* 260: 533–540.
- DeGeorge JJ, Nariai T, Yamazaki S, Williams WM, Rapoport SI (1991). Arecoline-stimulated brain incorporation of intravenously administered fatty acids in unanesthetized rats. *J Neurochem* 56: 352–355.
- De Souza EB, Kuyatt BL (1987). Autoradiographic localization of 3H-paroxetine-labeled serotonin uptake sites in rat brain. *Synapse* 1: 488–496.
- Dinan TG (1996). Serotonin: current understanding and the way forward. *Int Clin Psychopharmacol* 11(Suppl 1): 19–21.
- Duman RS, Heninger GR, Nestler EJ (1997). A molecular and cellular theory of depression. *Arch Gen Psychiatry* 54: 597–606.
- Felder CC, Kanterman RY, Ma AL, Axelrod J (1990). Serotonin stimulates phospholipase A₂ and the release of arachidonic acid in hippocampal neurons by a type 2 serotonin receptor that is independent of inositol phospholipid hydrolysis. *Proc Natl Acad Sci USA* 87: 2187–2191.
- Fitzpatrick F, Soberman R (2001). Regulated formation of eicosanoids. *J Clin Invest* 107: 1347–1351.
- Folch J, Lees M, Sloane Stanley GH (1957). A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 226: 497–509.
- Freo U, Ori C, Dam M, Merico A, Pizzolato G (2000). Effects of acute and chronic treatment with fluoxetine on regional glucose cerebral metabolism in rats: implications for clinical therapies. *Brain Res* 854: 35–41.
- Freo U, Soncrant TT, Holloway HW, Rapoport SI (1991). Dose- and time-dependent effects of 1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropane (DOI), a serotonergic 5-HT₂ receptor agonist, on local cerebral glucose metabolism. *Brain Res* 541: 63–69.
- Fryer JD, Lukas RJ (1999). Antidepressants noncompetitively inhibit nicotinic acetylcholine receptor function. *J Neurochem* 72: 1117–1124.
- Fuller RW (1995). Serotonin uptake inhibitors: uses in clinical therapy and in laboratory research. *Prog Drug Res* 45: 167–204.
- Fuller RW, Wong DT (1977). Inhibition of serotonin reuptake. *Fed Proc* 36: 2154–2158.
- Griebel G, Blanchard DC, Agnes RS, Blanchard RJ (1995). Differential modulation of antipredator defensive behavior in Swiss-Webster mice following acute or chronic administration of imipramine and fluoxetine. *Psychopharmacology (Berl)* 120: 57–66.
- Guan XM, McBride WJ (1988). Fluoxetine increases the extracellular levels of serotonin in the nucleus accumbens. *Brain Res Bull* 21: 43–46.
- Hayakawa T, Chang MC, Rapoport SI, Appel NM (2001). Selective dopamine receptor stimulation differentially affects [3H]arachidonic acid incorporation, a surrogate marker for phospholipase A₂-mediated neurotransmitter signal transduction, in a rodent model of Parkinson's disease. *J Pharmacol Exp Ther* 296: 1074–1084.
- Hervas I, Queiroz CM, Adell A, Artigas F (2000). Role of uptake inhibition and autoreceptor activation in the control of 5-HT release in frontal cortex and dorsal hippocampus of the rat. *Br J Pharmacol* 130: 160–166.
- Jacobson L, Sapolsky R (1991). The role of the hippocampus in feedback regulation of the hypothalamic-pituitary-adrenocortical axis. *Endocr Rev* 12: 118–134.
- Jones CR, Arai T, Bell JM, Rapoport SI (1996). Preferential *in vivo* incorporation of [³H]arachidonic acid from blood into rat brain synaptosomal fractions before and after cholinergic stimulation. *J Neurochem* 67: 822–829.
- Kaufman MJ, Hartig PR, Hoffman BJ (1995). Serotonin 5-HT_{2C} receptor stimulates cyclic GMP formation in choroid plexus. *J Neurochem* 64: 199–205.

- Kim HY, Edsall L, Garcia M, Zhang H (1999). The release of polyunsaturated fatty acids and their lipoxygenation in the brain. *Adv Exp Med Biol* **447**: 75–85.
- Kurt M, Arik AC, Celik S (2000). The effects of sertraline and fluoxetine on anxiety in the elevated plus-maze test in mice. *J Basic Clin Physiol Pharmacol* **11**: 173–180.
- Leonard BE (2001). Stress, norepinephrine and depression. *J Psychiatry Neurosci* **26**(Suppl): S11–S16.
- Li Q, Levy AD, Cabrera TM, Brownfield MS, Battaglia G, Van de Kar LD (1993). Long-term fluoxetine, but not desipramine, inhibits the ACTH and oxytocin responses to the 5-HT_{1A} agonist, 8-OH-DPAT, in male rats. *Brain Res* **630**: 148–156.
- Li Q, Ma L, William K, Murphy DL (2001). 5-HT_{2A} and 5-HT_{2C} receptors are differently regulated in 5-HT transporter mice: studies on the density and gene expression of the receptors. *Soc Neurosci Abstr* **26**: 380.5.
- Lin MT, Tsay HJ, Su WH, Chueh FY (1998). Changes in extracellular serotonin in rat hypothalamus affect thermoregulatory function. *Am J Physiol* **274**: R1260–R1267.
- Lira A, Bradley-Moore M, Zhou M, Fairhurst S, Menzaghi F, Brunner D, Gingrich JA (2001). Cognitive performance in 5-HT_{2A} receptor knockout mice. *Soc Neurosci Abstr* **27**: 380.4.
- Maier SF, Watkins LR (1998). Cytokines for psychologists: implications of bidirectional immune-to-brain communication for understanding behavior, mood, and cognition. *Psychol Rev* **105**: 83–107.
- Martin LL, Sanders-Bush E (1982). Comparison of the pharmacological characteristics of 5-HT₁ and 5-HT₂ binding sites with those of serotonin autoreceptors which modulate serotonin release. *Naunyn Schmiedebergs Arch Pharmacol* **321**: 165–170.
- Mazzola-Pomietto P, Aulakh CS, Tolliver T, Murphy DL (1997). Functional subsensitivity of 5-HT_{2A} and 5-HT_{2C} receptors mediating hyperthermia following acute and chronic treatment with 5-HT_{2A/2C} receptor antagonists. *Psychopharmacology (Berl)* **130**: 144–151.
- McReynolds AM, Meyer JS (1998). Effects of prenatal cocaine exposure on serotonin and norepinephrine transporter density in the rat brain. *Ann NY Acad Sci* **846**: 412–414.
- Mickley GA, Cobb BL, Farrell ST (1997). Brain hyperthermia alters local cerebral glucose utilization: a comparison of hyperthermic agents. *Int J Hyperthermia* **13**: 99–114.
- Moll GH, Mehnert C, Wicker M, Bock N, Rothenberger A, Ruther E, Huether G (2000). Age-associated changes in the densities of presynaptic monoamine transporters in different regions of the rat brain from early juvenile life to late adulthood. *Brain Res Dev Brain Res* **119**: 251–257.
- Murphy EJ, Rosenberger TA, Patrick CB, Rapoport SI (2000). Intravenously injected [¹⁴C]arachidonic acid targets phospholipids, and [¹⁴C]palmitic acid targets neutral lipids in hearts of awake rats. *Lipids* **35**: 891–898.
- Noronha JG, Bell JM, Rapoport SI (1990). Quantitative brain autoradiography of [^{9,10-³H}]palmitic acid incorporation into brain lipids. *J Neurosci Res* **26**: 196–208.
- Noronha JG, Larson DM, Rapoport SI (1989). Regional cerebral incorporation of plasma [¹⁴C]palmitate, and cerebral glucose utilization in water-deprived Long-Evans and Brattleboro rats. *Exp Neurol* **103**: 267–276.
- Owens MJ, Morgan WN, Plott SJ, Nemeroff CB (1997). Neurotransmitter receptor and transporter binding profile of antidepressants and their metabolites. *J Pharmacol Exp Ther* **283**: 1305–1322.
- Palvimaki EP, Roth BL, Majasuo H, Laakso A, Kuoppamaki M, Syvalahti E, Hietala J (1996). Interactions of selective serotonin reuptake inhibitors with the serotonin 5-HT_{2c} receptor. *Psychopharmacology (Berl)* **126**: 234–240.
- Paxinos G, Watson C (1987). *The Rat Brain in Stereotaxic Coordinates*, 3rd edn. Academic Press: New York.
- Pazos A, Palacios JM (1985). Quantitative autoradiographic mapping of serotonin receptor in the rat brain. I. Serotonin-1 receptors. *Brain Res* **346**: 205–230.
- Perry KW, Fuller RW (1993). Extracellular 5-hydroxytryptamine concentration in rat hypothalamus after administration of fluoxetine plus L-5-hydroxytryptophan. *J Pharmacol* **45**: 759–761.
- Purdon AD, Rapoport SI (1998). Energy requirements for two aspects of phospholipid metabolism in mammalian brain. *Biochem J* **335**: 313–318.
- Qu Y, Chang L, Klaff J, Balbo A, Rapoport SI (2001a). Imaging phospholipase A2 mediated signal transduction in response to a 5-HT₂ agonist in brain of awake rats. *J Neurochem* **78**(Suppl 1): 144.
- Qu Y, Chang L, Klaff J, Balbo A, Rapoport SI (2001b). *In vivo* neuroimaging of serotonin-mediated phospholipase A2 signal transduction in awake rats: effect of methiothepin. *J Neurochem (Abstr)* **78**(Suppl 1): 152.
- Qu Y, Chang L, Klaff J, Balbo A, Rapoport SI (2003). Imaging brain phospholipase A2 activation in awake rat in response to 5-HT_{2A/2C} agonist, (+)-2,5-dimethoxy-4-iodophenyl-2-aminopropane (DOI). *Neuropsychopharmacology* **28**: 244–252.
- Rabiner EA, Wilkins MR, Turkheimer F, Gunn RN, De Haes JU, De Vries M, Grasby PM (2002). 5-Hydroxytryptamine (1A) receptor occupancy by novel full antagonist 2-[4-[4-(7-chloro-2,3-dihydro-1,4-benzodioxyn-5-yl)-1-piperazinyl]butyl]-1, 2-benzisothiazol-3-(2H)-one-1,1-dioxide: A [(11)C][O-methyl-3H]-N-(2-(4-(2-methoxyphenyl)-1-piperazinyl)ethyl)-N-(2-pyridinyl)cyclohexanecarboxamide trihydrochloride (WAY-100635) positron emission tomography study in humans. *J Pharmacol Exp Ther* **301**: 1144–1150.
- Rapoport SI (1976). *Blood-Brain Barrier in Physiology and Medicine*. Raven Press: New York.
- Rapoport SI (2001). *In vivo* fatty acid incorporation into brain phospholipids in relation to plasma availability, signal transduction and membrane remodeling. *J Mol Neurosci* **16**: 243–261.
- Rapoport SI, Purdon D, Shetty HU, Grange E, Smith Q, Jones C, Chang MCJ (1997). *In vivo* imaging of fatty acid incorporation into brain to examine signal transduction and neuroplasticity involving phospholipids. *Ann NY Acad Sci* **820**: 56–74.
- Ricchieri GL, Soncrant TT, Holloway HW, Rapoport SI (1987). Methiothepin reduces glucose utilization in forebrain regions of awake rats. *Psychopharmacology (Berl)* **93**: 449–456.
- Robinson PJ, Noronha J, DeGeorge JJ, Freed LM, Nariai T, Rapoport SI (1992). A quantitative method for measuring regional *in vivo* fatty-acid incorporation into and turnover within brain phospholipids: review and critical analysis. *Brain Res Rev* **17**: 187–214.
- Rutter JJ, Auerbach SB (1993). Acute uptake inhibition increases extracellular serotonin in the rat forebrain. *J Pharmacol Exp Ther* **265**: 1319–1324.
- Sills TL, Greenshaw AJ, Baker GB, Fletcher PJ (1999). Acute fluoxetine treatment potentiates amphetamine hyperactivity and amphetamine-induced nucleus accumbens dopamine release: possible pharmacokinetic interaction. *Psychopharmacology (Berl)* **141**: 421–427.
- Sokoloff L (1999). Energetics of functional activation in neural tissues. *Neurochem Res* **24**: 321–329.
- Stanford JA, Currier TD, Gerhardt GA (2002). Acute locomotor effects of fluoxetine, sertraline, and nomifensine in young versus aged Fischer 344 rats. *Pharmacol Biochem Behav* **71**: 325–332.
- To CT, Anheuer ZE, Bagdy G (1999). Effects of acute and chronic fluoxetine treatment of CRH-induced anxiety. *Neuroreport* **10**: 553–555.
- Torres G, Horowitz JM, Laflamme N, Rivest S (1998). Fluoxetine induces the transcription of genes encoding c-fos, corticotropin-

- releasing factor and its type 1 receptor in rat brain. *Neuroscience* **87**: 463–477.
- Vial D, Piomelli D (1995). Dopamine D2 receptors potentiate arachidonate release via activation of cytosolic, arachidonic-specific phospholipase A2. *J Neurochem* **64**: 2765–2772.
- Wong DT, Bymaster FP, Engleman EA (1995). Prozac (fluoxetine, Lilly 110140), the first selective serotonin uptake inhibitor and an antidepressant drug: twenty years since its first publication. *Life Sci* **57**: 411–441.
- Wong DT, Threlkeld PG, Robertson DW (1991). Affinities of fluoxetine, its enantiomers, and other inhibitors of serotonin uptake for subtypes of serotonin receptors. *Neuropsychopharmacology* **5**: 43–47.
- Yoshino T, Nisijima K, Katoh S, Yui K, Nakamura M (2002). Tansospirone potentiates the fluoxetine-induced increases in extracellular dopamine via 5-HT(1A) receptors in the rat medial frontal cortex. *Neurochem Int* **40**: 355–360.
- Zhou FC, Tao-Cheng JH, Segu L, Patel T, Wang Y (1998). Serotonin transporters are located on the axons beyond the synaptic junctions: anatomical and functional evidence. *Brain Res* **805**: 241–254.