

Olanzapine Attenuates the Okadaic Acid-Induced Spatial Memory Impairment and Hippocampal Cell Death in Rats

Jue He^{1,3}, Yi Yang^{2,3}, Haiyun Xu^{1,3}, Xia Zhang¹ and Xin-Min Li^{*1}

¹Neuropsychiatry Research Unit, Department of Psychiatry, University of Saskatchewan, Saskatoon, SK, Canada; ²Department of Neurology, First Teaching Hospital, Jilin University, Chang Chun, China

It is hypothesized that olanzapine, an atypical antipsychotic drug, has beneficial effects on cognitive impairment and neuropathological changes in treating neurodegenerative diseases. In the present study, we investigated the effects of chronic administration of olanzapine on the spatial memory impairment and hippocampal cell death induced by the direct injection of okadaic acid (OA), a potent neurotoxin, into the rat hippocampus. After being pretreated with olanzapine (0.5 or 2 mg/kg/day, i.p.) for 2 weeks, the rats were unilaterally microinjected with OA (100 ng) into the hippocampus, and then were continuously administered with olanzapine for an additional week. The rats were trained on a spatial memory task in an eight-arm radial maze before OA administration, and tested on the same task 18 h after the last olanzapine injection. After the behavioral test, the rats were killed for Nissl staining and terminal deoxynucleotidyl transferase-mediated biotinylated UTP nick end labeling staining. OA significantly induced spatial memory impairment, and caused pyramidal cell loss in the CA1 and apoptotic cell death in the hippocampus. Olanzapine significantly attenuated OA-induced spatial memory impairment and the OA-induced neuropathological changes in the hippocampus. These findings suggest that olanzapine may have therapeutic effects in treatment of cognitive impairment and neuropathological changes of schizophrenia and other neurodegenerative diseases.

Neuropsychopharmacology (2005) **30**, 1511–1520, advance online publication, 11 May 2005; doi:10.1038/sj.npp.1300757

Keywords: olanzapine; okadaic acid; memory; hippocampus; cell death; neuroprotection

INTRODUCTION

Neuroanatomical and clinical studies of schizophrenia suggest that progressive neuropathological changes (such as neuronal atrophy and/or cell death) occur over the course of the disease (Woods *et al*, 1990; Waddington *et al*, 1991; Arnold and Trojanowski, 1996; DeLisi *et al*, 1997). Cognitive deficits tend to occur early in the course of schizophrenia, and the severity of deficits is predictive of the long-term treatment outcome for patients (Green, 1996). Neural injury or neurodegeneration may cause cognitive deficits in schizophrenia (Kraepelin, 1919; Harvey, 1998).

Olanzapine, an atypical antipsychotic drug, has been widely used in the treatment of schizophrenia and other psychoses without causing agranulocytosis (Fulton and Goa, 1997). It effectively treats positive and negative

symptoms and has a low association with tardive dyskinesia or parkinsonism (Beasley *et al*, 1996, 1997; Casey, 1997). Olanzapine is also associated with wide-ranging benefits on cognitive functioning, and may effectively alleviate cognitive impairment in schizophrenia patients (Purdon *et al*, 2000; Cuesta *et al*, 2001; Bilder *et al*, 2002; Harvey *et al*, 2003). Furthermore, we have previously demonstrated that olanzapine is effective in reducing PC12 cell death induced by hydrogen peroxide, β -amyloid peptide, or *N*-methyl-4-phenylpyridinium ions (MPP⁺) (Qing *et al*, 2003; Wei *et al*, 2003a,b). We have also shown that olanzapine effectively reduces the methamphetamine (METH)-induced mortality, and attenuates the METH-induced decrease of tyrosine hydroxylase immunoreactivity in the striatum in rats (He *et al*, 2004). We hypothesized that olanzapine has beneficial effects on cognitive impairment and neuropathological changes in treating neurodegenerative diseases.

Okadaic acid (OA), a selective and potent inhibitor of the serine/threonine phosphatases 1 and 2A (Ishihara *et al*, 1989; Cohen *et al*, 1990), has been reported to cause neuronal cell death *in vitro* (Cagnolli *et al*, 1996) and *in vivo* (Van Dam *et al*, 1998; He *et al*, 2001). OA may have rapid metabolic consequences leading to cell death by altering rates of phosphorylation–dephosphorylation *in vivo* (Arias

*Correspondence: Dr X-M Li, Neuropsychiatry Research Unit, Department of Psychiatry, University of Saskatchewan, 103 Wiggins Road, Saskatoon, SK, Canada SK 57N 5E4, Tel: +1 306 966 8573, Fax: +1 306 966 8830, E-mail: xin-min.li@usask.ca

³These authors contributed equally to this work

Received 5 January 2005; revised 23 March 2005; accepted 24 March 2005

Online publication: 1 April 2005 at <http://www.acnp.org/citations/Npp040105050001/default.pdf>

et al, 1998). In cultured rat cortical neurons, OA may induce cell death by increasing phosphorylation of microtubule-associated protein 2 and tau concomitant with early changes in the neuronal cytoskeleton (Arias *et al*, 1993). OA may also induce cellular injury and death by inactivating Bcl-2 by phosphorylation (Halder *et al*, 1995), activating caspase-3 (Hong *et al*, 2002), or forming high reactive free radicals (Matias *et al*, 1999) *in vitro*. Infusion of OA into the rat brain results in severe memory impairment, accompanied by remarkable neuropathological changes including hippocampal neurodegeneration, a paired helical filament-like phosphorylation of tau protein, and the formation of β /A4-amyloid containing plaque-like structures in gray and white matter areas (Arendt *et al*, 1995, 1998; Zhao *et al*, 1995; He *et al*, 2001; Tian *et al*, 2004).

In the present study, unilateral microinjection of OA into the dorsal hippocampus in rats was used as a memory impairment and hippocampal neurodegeneration animal model. The possible beneficial effect of olanzapine on the OA-induced spatial working and reference memory impairment was examined in a radial arm maze task, and the neuroprotective potential of olanzapine on OA-induced lesion in the hippocampus was evaluated by Nissl staining and terminal deoxynucleotidyl transferase-mediated biotinylated UTP nick end labeling (TUNEL) staining.

MATERIALS AND METHODS

Animals

Male Sprague-Dawley rats (Charles River, St Constant, QC) weighing 225–250 g at the beginning of the experiments were used. They were housed four or five per cage with free access to food and water under controlled laboratory conditions (a 12:12 light:dark cycle at room temperature (19–21°C)). The rats were acclimated to the laboratory for 7 days before the experiments. All procedures with animals were performed in accordance with the guidelines established by the Canadian Council on Animal Care and were approved by the University of Saskatchewan Animal Care Committee.

Preliminary Experiment

In order to decide the dose of OA for olanzapine experiment, different doses (0, 50, and 100 ng/ μ l saline) of OA were injected into the right hippocampus (the same injection method described in Surgery and microinjection) after spatial reference/working memory task training in the rats (see Radial arm maze task). The rats were tested for spatial memory (see Radial arm maze task) 1 week after OA injection.

Drug Treatment

Olanzapine (Eli Lilly and Company, Indianapolis, IN) was freshly dissolved in 0.2% glacial acetic acid in saline. The ammonium salt of OA (Research Biochemicals International, Natick, MA) was dissolved to a concentration of 122 μ M (100 ng/ μ l) in saline. The total experimental period was 22 days. The rats were pretreated with olanzapine (0.5

or 2 mg/kg/day) or 0.2% glacial acetic acid in saline as vehicle 1 by intraperitoneal (i.p.) injection once a day for 2 weeks. On day 15, the rats were microinjected with OA (100 ng in 1 μ l saline) or saline (vehicle 2; 1 μ l) into the right hippocampus (see Surgery and microinjection). At 1 h after the microinjection of OA or saline, the rats were continuously administrated with olanzapine (0.5 or 2 mg/kg/day) or vehicle 1 by i.p. injection (1.0 ml/kg) once a day for 1 week. These procedures produced the following four groups of rats: vehicle 1 + vehicle 2 (CON), vehicle 1 + OA (OA), olanzapine 0.5 mg/kg/day + OA (OLA0.5 + OA), and olanzapine 2.0 mg/kg/day + OA (OLA2.0 + OA).

Surgery and Microinjection

On day 15, the rats were anesthetized with isoflurane inhalation (2.0% isoflurane in oxygen for induction, 1.5% isoflurane in oxygen for maintenance), and restrained in a stereotactic apparatus. OA (100 ng in 1 μ l saline) or saline (1 μ l) was injected into the right dorsal hippocampus (A: –3.8 mm, L: 2.5 mm from bregma, and V: 3.2 mm) according to the atlas of Paxinos and Watson (1986). OA or saline was injected over a 2 min period, and the injection cannula was left in place for an additional 5 min to allow for diffusion of OA away from the injection site. The dosage of OA (100 ng) was based on our preliminary data (Figure 1) and the published reports (Arias *et al*, 1998; He *et al*, 2001). The position of the injection site was examined in Nissl-stained hippocampal sections. The rats that had injection sites away from the dorsal hippocampus were omitted from data analysis.

Radial Arm Maze Task

The radial arm maze used in the present study consisted of eight arms, numbered from 1 to 8 (48 \times 12 cm), extending radially from a central area (32 cm in diameter), with a 5 cm edge around the apparatus. The maze was placed 40 cm above the floor and was surrounded by various extramaze cues such as different laboratory benches and posters. Rats were trained for the spatial reference/working memory task, as described previously (Zou *et al*, 1998; Mizuno *et al*, 2000; He *et al*, 2002a, b). From day 2 to 5, the rats were shaped in the maze to run to the end of the arms and consume the bait, in groups of four rats. The bait was initially available throughout the maze, but was gradually restricted to the food cup. Following this shaping period, each rat was placed individually in the center of the maze and subjected to the reference and working memory task training from day 6 to 14 (9 days, 3 trials/day with a 5 min interval). Finally, each rat was individually tested for the reference and working memory task on day 22 (1 week after OA or saline microinjection). The first training trial of each day and the test trial (on day 22) were performed 18 h after the injection of olanzapine or vehicle 1. Olanzapine administered 18 h before the radial maze training and test did not evoke any sedating effect during maze experiments in rats. In all training and test trials, the same four arms (nos. 1, 2, 4, and 7) were baited. The other four arms (nos. 3, 5, 6, and 8) were never baited. The training or test trial continued until all four baits in the food cups had been consumed or

until 5 min had elapsed. Reference memory is regarded as a long-term memory for information that remains constant over repeated trials (memory for the positions of baited arms), while working memory is considered as a short-term memory in which the information to be remembered changes in every trial (memory for the positions of arms that had already been visited in each trial; Olton *et al*, 1979). Thus, measures were made of the number of reference memory errors (entering an arm that was not baited) and working memory errors (entering an arm containing food but previously entered).

Locomotor Speed and Food Consumption

We measured locomotor speed and food consumption to see whether motor function and/or motivation were affected by the treatments (OA and olanzapine), because the animals' motor function and motivation may have an impact on their performance in the radial arm maze task. Locomotor speed was calculated by total time/number of total entries in the test trial of the spatial memory task on day 22. On the same day (day 22), after the spatial memory test, rats were measured for food consumption. The rats were individually placed in a home cage, and then 10 baits, which were the same as the one used in the radial arm maze task, were provided. The time taken to consume all 10 baits in the food cup was recorded, with a cutoff time of 180 s (Zou *et al*, 1998).

Nissl Staining and TUNEL Staining

On day 22, the rats were deeply anesthetized with chloral hydrate (400 mg/kg, *i.p.*) after the reference/working memory test, and were perfused through the ascending aorta with 250–300 ml of 0.1 M phosphate-buffered saline (PBS, pH 7.4) followed by 300–350 ml of a fixative solution of 4% paraformaldehyde in PBS. The brains were removed, postfixed in the same fixative for 24 h, and then cryoprotected in 30% sucrose in PBS. The brains were cut into 40- μ m-thick coronal sections on a cryostat. The sections throughout the hippocampus were mounted on gelatin-coated slides and allowed to air-dry overnight.

The surviving pyramidal cells in the hippocampus of rats were visualized by Nissl staining (He *et al*, 2001; Zhang *et al*, 2002). The mounted sections were rehydrated in distilled water, and submerged in 0.5% cresyl violet solution for about 1 min until the desired depth of staining was achieved.

Apoptotic nuclei showing nucleosomal DNA fragmentation can be detected by TUNEL staining (Gavrieli *et al*, 1992). The apoptotic cells in the hippocampus of rats were stained by a TUNEL staining kit according to the manufacturer's instructions (Trevigen, Gaithersburg, MD; Zhang *et al*, 2002). TUNEL staining relies on the detection of fragmented DNA strands, but because fragmentation can occur via nonapoptotic mechanisms, it is not absolutely specific for apoptosis (Wang *et al*, 2000).

Cell Counting and Data Analysis

The number of the CA1 pyramidal cells in Nissl staining sections (three sections of the right hemisphere of each rat

at levels of -3.60 , -3.80 , and -4.00 mm; Paxinos and Watson, 1986) was counted at $\times 400$ magnification. Cells were counted only if structures of appropriate size and shape were demonstrated clearly. Questionable structures were examined under $\times 1000$ magnification and were not counted if identification remained uncertain. Similarly, the number of TUNEL-positive cells in the sections of the whole right hippocampus of each rat at levels of -3.64 , -3.84 , and -4.04 mm (Paxinos and Watson, 1986) was also counted at $\times 400$ magnification. Although we did not use a stereological method for cell counting, our cell counting analysis provided complementary data to our qualitative observation on the hippocampal sections.

Statistical Analysis

All results were expressed as means \pm SEM. A two-tailed *t*-test for independent samples was used for the CON and OA group comparison. The significant difference was determined by a one-way ANOVA, followed by Bonferroni's test for multiple comparisons. The significant difference of the spatial memory formation before OA injection in the CON, OA, OLA0.5 + OA, and OLA2.0 + OA groups was determined by a two-way ANOVA factoring time (daily training of 9 days) and drug treatment (olanzapine). A *P*-value of less than 0.05 was regarded as statistically significant.

RESULTS

Preliminary Experiment: OA (100 ng) Induced Spatial Memory Impairment in the Radial Arm Maze Task

The number of working and reference memory errors measured premicroinjection and 1 week after microinjection in the saline-injected (CON), OA (50 ng)-injected (OA50), and OA (100 ng)-injected (OA100) groups is shown in Figure 1. One-way ANOVA showed that there was no difference in the number of working ($F(2,12) = 0.169$, $P = 0.846$) or reference ($F(2,12) = 0.300$, $P = 0.746$) memory errors on the last training day (premicroinjection) among the CON, OA50, and OA100 groups. However, one-way ANOVA showed that OA had an effect on the number of working ($F(2,12) = 7.176$, $P = 0.009$) and reference ($F(2,12) = 17.765$, $P < 0.001$) memory errors after microinjection of OA, and a *post hoc* analysis with Bonferroni's test indicated that the number of working memory errors in the OA100 group was significantly higher than that in the CON group ($P < 0.01$; Figure 1a), and that the number of reference memory errors in the OA100 group was significantly higher than that in the CON group ($P < 0.001$; Figure 1b) and the OA50 group ($P < 0.01$; Figure 1b).

Chronic Administration of Olanzapine did not Affect the Spatial Memory Formation of Rats before OA Injection

Figures 2a and b show the time course (total 9 days; from day 6 to 14) of working memory (Figure 2a) and reference memory (Figure 2b) formation during the training period of the radial arm maze task performed

before OA injection. Two-way ANOVA disclosed that daily training (time) significantly decreased the number of both working ($F(8,24)=39.707$, $P<0.001$) and refer-




Figure 1 Dose effect of OA on the working (a) and reference (b) memory in the radial arm maze task. The working and reference memory was measured 1 week after the microinjection of OA or saline into the right hippocampus. Results are expressed as means \pm SEM ($n=5$ in each group). ** $P<0.01$ and *** $P<0.001$ vs CON, and ## $P<0.01$ vs OA50.

ence ($F(8,24)=132.979$, $P<0.001$) memory errors, and that olanzapine produced no effect on the number of working ($F(3,24)=1.606$, $P=0.217$) or reference ($F(3,24)=1.936$, $P=0.153$) memory errors with no interaction between the two factors (time and olanzapine) in the number of working ($F(8,24)=1.189$, $P=0.258$) or reference ($F(8,24)=0.966$, $P=0.513$) memory errors. One-way ANOVA showed that there was no difference in the number of working ($F(3,22)=0.612$, $P=0.614$) or reference ($F(3,22)=0.334$, $P=0.801$) memory errors on the last training day among the CON, OA, OLA0.5 + OA, and OLA2.0 + OA groups before saline or OA injection.

Chronic Administration of Olanzapine Attenuated the OA-Induced Spatial Memory Impairment in the Radial Arm Maze Task

The number of working and reference memory errors in the rats was measured 1 week after OA or saline microinjection and the data are shown in Figures 2c and d. A two-tailed t -test for independent samples disclosed that the number of working ($P<0.001$; Figure 2c) and reference ($P<0.001$; Figure 2d) memory errors in the OA group is significantly higher than that in the CON. One-way ANOVA showed that the treatment (olanzapine) had an effect on the number of working ($F(2,16)=5.094$, $P=0.019$) and reference ($F(2,16)=12.827$, $P<0.001$) memory errors in OA-injected rats, and a *post hoc* analysis with Bonferroni's test indicated that the number of working memory errors in the OLA2.0 + OA group is significantly lower than that in the OA group ($P<0.05$; Figure 2c), and that the number of reference memory errors in the OLA2.0 + OA group is significantly




Figure 2 Olanzapine did not affect the spatial working (a) and reference (b) memory formation before OA or saline microinjection in the radial arm maze task in rats. However, olanzapine significantly attenuated the OA-induced impairment in animals' working (c) and reference (d) memory, which was measured by the radial arm maze task 1 week after the microinjection of OA or saline into the right hippocampus. Results are expressed as means \pm SEM ($n=6-7$ in each group). *** $P<0.001$ vs CON, # $P<0.05$ and ### $P<0.001$ vs OA, and + $P<0.05$ vs OLA0.5 + OA.

lower than that in the OA group ($P < 0.001$; Figure 2d) and the OLA0.5 + OA group ($P < 0.05$; Figure 2d).

Chronic Administration of Olanzapine and OA Injection had no Effect on Food Consumption and Locomotor Speed of the Rats

The treatments (OA and olanzapine) had no effect on locomotor speed and food consumption in the rats. A two-tailed t -test for independent samples disclosed that there was no difference in locomotor speed and food consumption between the OA and CON groups. One-way ANOVA showed that the treatment (olanzapine) had no effect on locomotor speed ($F(2,16) = 1.862$, $P = 0.188$) and food consumption ($F(2,16) = 0.316$, $P = 0.734$) in OA-injected rats.

Chronic Administration of Olanzapine Attenuated the OA-Induced Decrease in the Number of Surviving Pyramidal Cells in the CA1 Region of Hippocampus

Nissl staining of hippocampal sections was used to evaluate the effect of olanzapine on the OA-induced decrease in the number of surviving pyramidal cells in the CA1 region of the injected hippocampus. Representative photomicrographs of Nissl staining in the hippocampus in each group are shown in Figure 3a.

As shown in Figure 3b, the number of surviving pyramidal cells in the CA1 region of the injected hippocampus was counted in each group. A two-tailed t -test for independent samples disclosed that the number of pyramidal cells in the CA1 region of the injected hippocampus significantly decreased in the OA group compared with that in the CON group ($P < 0.001$). One-




Figure 3 (a) Representative photomicrographs of Nissl staining in the injected hippocampus of the rats in the CON, OA, OLA0.5 + OA, and OLA2.0 + OA groups. In the OA-injected hippocampi (OA, OLA0.5 + OA, and OLA2.0 + OA), the lesion is obvious. The lesion in the rats treated with OA alone (OA) is more extensive, fragile, and easier to be peeled off when the brains were cut compared with that in the rats treated with both olanzapine and OA (OLA2.0 + OA). The scale bar is 300 μ m. (b) Quantitative analysis of the effect of olanzapine on the OA-induced pyramidal cell loss in the CA1 region of hippocampus. The number of CA1 pyramidal cells in Nissl staining sections was counted at $\times 400$ magnification. Results are expressed as means \pm SEM ($n = 6-7$ in each group). *** $P < 0.001$ vs CON and ## $P < 0.01$ vs OA.

way ANOVA showed that the treatment (olanzapine) had an effect ($F(2,16) = 7.489$, $P = 0.005$) in the OA-injected rats, and a *post hoc* analysis with Bonferroni's test indicated that

the number of pyramidal cells in the CA1 region of the injected hippocampus in the OLA2.0 + OA group is significantly higher than that in the OA group ($P < 0.01$).




Figure 4 (a–d, a'–d') Representative photomicrographs of TUNEL staining in the injected hippocampus of the rats in the CON (a, a'), OA (b, b'), OLA0.5 + OA (c, c'), and OLA2.0 + OA (d, d') groups. The high magnification of right photomicrographs (a', b', c', and d') were picked up from a, b, c, and d, respectively. The arrows on the low-magnification panels indicate the location of the high-magnification images. In the hippocampi of OA-injected groups (b, b'; c, c'; d, d'), the TUNEL-positive cells are visible in different frequency, whereas there are almost no TUNEL-positive cells in the hippocampus of the control rat (a, a'). The scale bar represents 300 μm in a–d, and represents 30 μm in a'–d'. (e) Quantitative analysis of the effect of olanzapine on the OA-induced increase of TUNEL-positive cells in the injected hippocampus. The number of TUNEL-positive cells in the hippocampus was counted at $\times 400$ magnification. Results are expressed as means \pm SEM ($n = 6$ –7 in each group). *** $P < 0.001$ vs CON, ## $P < 0.01$ vs OA, and + $P < 0.05$ vs OLA0.5 + OA.




Figure 5 Scatter plot of the number of working (a) or reference (b) memory errors against the number of TUNEL-positive cells in the hippocampus after OA or saline injection in each individual rat in the CON, OA, OLA0.5 + OA, and OLA2.0 + OA groups ($n = 6-7$ in each group).

Chronic Administration of Olanzapine Attenuated the Number of OA-Induced TUNEL-Positive Cells in the Hippocampus

The hippocampal sections prepared by TUNEL staining were used to evaluate the effect of olanzapine on OA-induced apoptotic cell death in the injected hippocampus. Representative photomicrographs of TUNEL staining in the hippocampus in each group are shown in Figures 4a-d. Under a higher magnification ($\times 400$), many TUNEL-positive cells were observed in the hippocampus of the OA-injected rats (Figures 4b', 4c', and 4d') but not in that of the CON rats (Figure 4a').

As shown in Figure 4e, the number of TUNEL-positive cells in the injected hippocampus was counted in each group. A two-tailed *t*-test for independent samples disclosed that the number of TUNEL-positive cells in the injected hippocampus significantly increased in the OA group compared with that in the CON group ($P < 0.001$). One-way ANOVA showed that the treatment (olanzapine) had an effect ($F(2,16) = 9.247$, $P = 0.002$) in the OA-injected rats, and a *post hoc* analysis with Bonferroni's test indicated that the number of TUNEL-positive cells in the injected hippocampus in the OLA2.0 + OA group is significantly lower than that in the OA group ($P < 0.01$) and that in the OLA0.5 + OA group ($P < 0.05$).

Spatial Memory Errors Increased with the Increasing Number of TUNEL-Positive Cells in the Hippocampus

The relationship between the number of working memory errors and the number of TUNEL-positive cells in the hippocampus was significant ($r^2 = 0.6292$, $P < 0.0001$;

Figure 5a). There was also a significant direct relationship between the number of reference memory errors and the number of TUNEL-positive cells in the hippocampus ($r^2 = 0.6835$, $P < 0.0001$; Figure 5b).

DISCUSSION

Memory impairment induced by microinjection of OA into animal brains has been reported (Arendt *et al*, 1995; He *et al*, 2001; Bennett *et al*, 2003). Similarly, in the present study, unilateral microinjection of OA (100 ng) into the dorsal hippocampus induced impairment in spatial working and reference memory. Consistent with the previous reports (Arias *et al*, 1998; He *et al*, 2001; Ramirez-Munguia *et al*, 2003), Nissl staining in the present study showed that the direct microinjection of OA into the dorsal hippocampus robustly decreased the number of surviving pyramidal neurons in the CA1 region of the hippocampus. Microinjection of OA into the hippocampus also induced apoptosis as revealed by TUNEL staining. This is in accordance with previous studies that showed the OA-induced apoptosis in HL60 leukemia (Riordan *et al*, 1998), neuroblastoma (Cabado *et al*, 2004), and mammalian cells (Boe *et al*, 1991). Because the intact hippocampus is required for the recall, item recognition, and associative recognition memory of animals (Alvarado and Bachevalier, 2005; Sweatt, 2004), the OA-induced spatial memory impairment may be attributed, at least in part, to the hippocampal cell death caused by the drug.

Chronic administration of olanzapine significantly attenuated the OA-induced spatial memory impairment. This improvement is parallel to the alleviative effect of olanzapine on the OA-induced cell death in the hippocampus, suggesting an association between the neuroprotective effect and the memory-improving effect exerted by olanzapine.

The attenuating effect of chronic administration of olanzapine on the OA-induced neurodegeneration and apoptosis provides the direct evidence supporting the neuroprotective action of olanzapine. Although the mechanisms responsible for the neuroprotective effect remain to be elucidated, our previous studies have provided some insights into this issue. In PC12 cells, olanzapine is effective in reducing the cell death induced by H_2O_2 , $A\beta$, or MPP^+ , which may cause oxidative stress thus killing the treated cells (Qing *et al*, 2003; Wei *et al*, 2003a,b). We demonstrated that these cytoprotective actions in the cells might involve the upregulation of superoxide dismutase-1, one of the endogenous antioxidant enzymes, by the drug (Li *et al*, 1999; Qing *et al*, 2003; Wei *et al*, 2003a). In addition, olanzapine can regulate the translocation and expression of pro- and antiapoptotic proteins Bax and Bcl-X_L in PC12 cells (Wei *et al*, 2003b). In animal studies, chronic administration of olanzapine was shown to upregulate the expression of Bcl-2 and BDNF in the hippocampus (Bai *et al*, 2003, 2004) and to help restore the repeated restraint stress-induced decrease in these two neuroprotective proteins in hippocampal neurons (Luo *et al*, 2004). Furthermore, the prevention of Bcl-2 decrease has been associated with the protective effects of olanzapine on methamphetamine-induced neurotoxicity (He *et al*, 2004).

Thus our own previous data, together with the findings by other investigators that OA-induced apoptosis is associated with downregulation of Bcl-2 and can be prevented by upregulation of Bcl-2 (Benito *et al*, 1997; Nuydens *et al*, 2000; Cabado *et al*, 2004), suggest that Bcl-2 plays an important role in the neuroprotective effects of olanzapine on OA-induced neurodegeneration and apoptosis. Olanzapine may also attenuate OA-induced neurotoxicity by upregulating superoxide dismutase (Manna *et al*, 1998; Li *et al*, 1999; Matias *et al*, 1999), and perform protective effects on OA-induced apoptotic cell death by modulating the expression of pro- and antiapoptotic proteins such as Bax and Bcl-X_L (Wei *et al*, 2003b; Cabado *et al*, 2004). However, further studies are necessary to elucidate whether olanzapine attenuated OA-induced neurotoxicity by directly affecting the activation of phosphatases or caspases.

The time point (1 week after OA injection) for spatial memory measurement ruled out the direct effect of OA (as an inhibitor of PP1 and PP2A) on memory processes in the present study (He *et al*, 2001; Bennett *et al*, 2003). It is likely that OA-induced spatial memory impairment in the present paradigm is due to the secondary effect of OA-induced hippocampal cell death (He *et al*, 2001). Unilateral microinjection of OA (100 ng), which induced spatial memory impairment, induced a massive lesion of the normal brain tissue (Figure 3a) in the OA-injected rats. The lesion in the rats treated with OA alone was more extensive, fragile, and easier to be peeled off when the brains were cut on a cryostat compared with that in the rats treated with both olanzapine and OA (Figure 3a). Chronic olanzapine not only significantly attenuated OA-induced decrease of the number of pyramidal cells in the CA1 region, but also attenuated OA-induced increase in apoptotic cell death in the hippocampus. Therefore, the improving effects of olanzapine on OA-induced spatial memory impairment in rats may be subsequent to its attenuating effects on OA-induced hippocampal cell death.

It should be noted that there must be some molecules linking the attenuating effects of OA-induced hippocampal cell death (neuroprotective effect) and memory improvement. Among possible molecules, BDNF is a stronger candidate. It has been shown that chronic administration of olanzapine upregulates the expression of BDNF mRNA in the hippocampus (Bai *et al*, 2003), and improves memory in different maze tasks in animal studies (Nowakowska *et al*, 1999; Wolff and Leander, 2003). Indeed, the involvement of BDNF in spatial memory formation and maintenance in a radial arm maze test has been demonstrated in rats (Mizuno *et al*, 2000; Radecki *et al*, 2005).

Chronic administration of olanzapine and OA injection had no effect on locomotor speed and food consumption of the animals. This result rules out the possibility that olanzapine may affect motor function or/and motivation for food thus improving their performance in the radial arm maze task. However, it is still possible that OA-induced behavioral deficits are not limited to memory deficits, and the capacity of olanzapine to reverse other types of behavioral deficits remains unknown.

Although the above explanation is helpful for us to understand the mechanisms responsible for the improving effect of olanzapine on the OA-induced memory impairment, there are other experimental data meriting our

attention. It was reported that injection of OA into the nucleus basalis of Meynert, which is a major source of cholinergic innervation of the cerebral cortex and hippocampus, induced a decreased level of acetylcholine (ACh) in the brain and spatial memory deficit in rats (Tian *et al*, 2004). On the other hand, olanzapine in rats induces an increase of ACh release in the medial prefrontal cortex and hippocampus, a possible factor contributing to cognitive improvement in schizophrenia (Ichikawa *et al*, 2002; Shirazi-Southall *et al*, 2002). Therefore, the effects of olanzapine on ACh may be another possible contributor to its improving effect on the OA-induced memory impairment.

Although chronic administration of olanzapine (14 days, 0.5 or 2 mg/kg/day) had no effects on spatial memory formation (Figures 2a and b), we still cannot completely exclude the possibility that olanzapine itself affects memory performance because there was no olanzapine control group (chronic olanzapine treatment (21 days, 0.5 or 2 mg/kg/day) + intrahippocampal vehicle microinjection) in the present study. However, because of the length of the training and the interval between the training and test, it is possible that even if olanzapine by itself had some memory-enhancing effect, this effect could not be detected in the olanzapine control group. In this context, the alternative possibility that olanzapine may reverse OA effect on memory through a nonspecific memory-enhancing effect should be considered. Moreover, further studies are necessary to elucidate whether the effects of olanzapine observed in the present model system are also shared by other atypical antipsychotic drugs.

In conclusion, unilateral microinjection of OA significantly induced spatial working and reference memory impairment, and caused neurodegeneration and apoptosis in the injected hippocampus. Chronic pre- and postlesion administration of olanzapine effectively attenuated OA-induced spatial memory impairment and the OA-induced neuropathological changes in the hippocampus. These findings suggest that olanzapine may have therapeutic effects in treatment of cognitive impairment and neuropathological changes of schizophrenia and other neurodegenerative diseases.

ACKNOWLEDGEMENTS

This work was supported by the Saskatchewan Health Research Foundation Postdoctoral Fellowship to Jue He. We thank Dr Augusto V Juorio and Dr Sergey Fedoroff for their helpful comments during the preparation of this manuscript. We are also grateful to Gabriel Stegeman for her excellent technical assistance.

REFERENCES

- Alvarado MC, Bachevalier J (2005). Selective neurotoxic damage to the hippocampal formation impairs performance of the transverse patterning and location memory tasks in rhesus macaques. *Hippocampus* 15: 118–131.
- Arendt T, Holzer M, Fruth R, Bruckner MK, Gartner U (1995). Paired helical filament-like phosphorylation of tau, deposition of β /A4-amyloid and memory impairment in rat induced by

- chronic inhibition of phosphatase 1 and 2A. *Neuroscience* **69**: 691–698.
- Arendt T, Holzer M, Fruth R, Bruckner MK, Gartner U (1998). Phosphorylation of tau, A β -formation, and apoptosis after *in vivo* inhibition of PP-1 and PP-2A. *Neurobiol Aging* **19**: 3–13.
- Arias C, Becerra-Garcia F, Arrieta I, Tapia R (1998). The protein phosphatase inhibitor okadaic acid induces heat shock protein expression and neurodegeneration in rat hippocampus *in vivo*. *Exp Neurol* **153**: 242–254.
- Arias C, Sharma N, Davies P, Shafit-Zagardo B (1993). Okadaic acid induced early changes in microtubule-associated protein 2 and tau phosphorylation prior to neurodegeneration in cultured cortical neurons. *J Neurochem* **61**: 673–682.
- Arnold SE, Trojanowski JQ (1996). Recent advances in defining the neuropathology of schizophrenia. *Acta Neuropathol* **92**: 217–231.
- Bai O, Chlan-Fourney J, Bowen R, Keegan D, Li XM (2003). Expression of brain-derived neurotrophic factor mRNA in rat hippocampus after treatment with antipsychotic drugs. *J Neurosci Res* **71**: 127–131.
- Bai O, Zhang H, Li XM (2004). Antipsychotic drugs clozapine and olanzapine upregulate bcl-2 mRNA and protein in rat frontal cortex and hippocampus. *Brain Res* **1010**: 81–86.
- Beasley CM, Sanger T, Satterlee W, Tollefson GD, Tran PV, Hamilton S (1996). Olanzapine vs. placebo: results of a double-blind, fixed-dose olanzapine trial. *Psychopharmacology* **124**: 159–167.
- Beasley CM, Tollefson GD, Tran PV (1997). Efficacy of olanzapine: an overview of pivotal clinical trials. *J Clin Psychiatry* **58**: 7–12.
- Benito A, Lerga A, Silva M, Leon J, Fernandez-Luna JL (1997). Apoptosis of human myeloid leukemia cells induced by an inhibitor of protein phosphatases (okadaic acid) is prevented by Bcl-2 and Bcl-X(L). *Leukemia* **11**: 940–944.
- Bennett PC, Moutsoulas P, Lawen A, Perini E, Ng KT (2003). Novel effects on memory observed following unilateral intracranial administration of okadaic acid, cyclosporin A, FK506 and [MeVal⁴]CyA. *Brain Res* **988**: 56–58.
- Bilder RM, Goldman RS, Volavka J, Czobor P, Hoptman M, Sheitman B et al (2002). Neurocognitive effects of clozapine, olanzapine, risperidone, and haloperidol in patients with chronic schizophrenia or schizoaffective disorder. *Am J Psychiatry* **159**: 1018–1028.
- Boe R, Gjertsen BT, Vintermyr OK, Houge G, Lanotte M, Doskeland SO (1991). The protein phosphatase inhibitor okadaic acid induces morphological changes typical of apoptosis in mammalian cell. *Exp Cell Res* **195**: 237–246.
- Cabado AG, Leira F, Vieytes MR, Vieites JM, Botana LM (2004). Cytoskeletal disruption is the key factor that triggers apoptosis in okadaic acid-treated neuroblastoma cells. *Arch Toxicol* **78**: 74–85.
- Cagnoli CM, Kharlamov E, Atabay C, Uz T, Manev H (1996). Apoptosis induced in neuronal cultures by either the phosphatase inhibitor okadaic acid or the kinase inhibitor staurosporine is attenuated by isoquinolinesulfonamides H-7, H-8 and H-9. *J Mol Neurosci* **7**: 65–76.
- Casey DE (1997). The relationship of pharmacology to side effects. *J Clin Psychiatry* **58**: s55–s62.
- Cohen P, Holmes CF, Tsukitani Y (1990). Okadaic acid: a new probe for the study of cellular regulation. *Trends Biol Sci* **15**: 98–102.
- Cuesta MJ, Peralta V, Zarzuela A (2001). Effects of olanzapine and other antipsychotics on cognitive function in chronic schizophrenia: a longitudinal study. *Schizophr Res* **48**: 17–28.
- deLisi LE, Sakuma M, Rew W, Kushner M, Hoff AL, Grimson R (1997). Schizophrenia as a chronic active brain process: a study of progressive brain structural change subsequent to the onset of schizophrenia. *Psychiatry Res* **74**: 129–140.
- Fulton B, Goa KL (1997). Olanzapine. A review of its pharmacological properties and therapeutic efficacy in the management of schizophrenia and related psychoses. *Drugs* **53**: 281–298.
- Gavrieli Y, Sherman Y, Ben-Sasson SA (1992). Identification of programmed cell death *in situ* via specific labeling of nuclear DNA fragmentation. *J Cell Biol* **119**: 493–501.
- Green MF (1996). What are the functional consequences of neurocognitive deficits in schizophrenia? *Am J Psychiatry* **153**: 321–330.
- Haldar S, Jena N, Croce CM (1995). Inactivation of Bcl-2 by phosphorylation. *Proc Natl Acad Sci USA* **92**: 4507–4511.
- Harvey PD (1998). Schizophrenia and related disorders in late life. In: Bellack AS, Hersen M (eds). *Comprehensive Clinical Psychology*. Elsevier: Amsterdam. Vol 8, pp 74–86.
- Harvey PD, Green MF, McGurk SR, Meltzer HY (2003). Changes in cognitive functioning with risperidone and olanzapine treatment: a large-scale, double-blind, randomized study. *Psychopharmacology* **169**: 404–411.
- He J, Xu H, Yang Y, Zhang X, Li XM (2004). Neuroprotective effects of olanzapine on methamphetamine-induced neurotoxicity are associated with an inhibition of hyperthermia and prevention of Bcl-2 decrease in rats. *Brain Res* **1018**: 186–192.
- He J, Yamada K, Nabeshima T (2002a). A role of Fos expression in the CA3 region of the hippocampus in spatial memory formation in rats. *Neuropsychopharmacology* **26**: 259–268.
- He J, Yamada K, Zou L, Nabeshima T (2001). Spatial memory deficit and neurodegeneration induced by the direct injection of okadaic acid into the hippocampus in rats. *J Neural Transm* **108**: 1435–1443.
- He J, Yamada K, Zou L, Nakajima A, Kamei H, Nabeshima T (2002b). Learning and memory in two different reward tasks in a radial arm maze in rats. *Behav Brain Res* **134**: 139–148.
- Hong HN, Yoon SY, Suh J, Lee JH, Kim D (2002). Differential activation of caspase-3 at two maturational stages during okadaic acid-induced rat neuronal death. *Neurosci Lett* **334**: 63–67.
- Ichikawa J, Dai J, O'Laughlin IA, Fowler WL, Meltzer HY (2002). Atypical, but not typical, antipsychotic drugs increase cortical acetylcholine release without an effect in the nucleus accumbens or striatum. *Neuropsychopharmacology* **26**: 325–339.
- Ishihara H, Martin BL, Brautigam DL, Karaki H, Ozaki Y et al (1989). Calyculin A and okadaic acid: inhibitors of protein phosphatase activity. *Biochem Biophys Res Commun* **159**: 871–877.
- Kraepelin E (1919). *Dementia Praecox and Paraphrenia*. E & S Livingstone: Edinburgh.
- Li X-M, Chlan-Fourney J, Bennett V, Shrikhande S, Juorio AV, Keegan D et al (1999). Differential effects of Olanzapine on the gene expression of superoxide dismutase and low affinity nerve growth factor receptor. *J Neurosci Res* **56**: 72–75.
- Luo C, Xu H, Li XM (2004). Post-stress changes in BDNF and Bcl-2 immunoreactivities in hippocampal neurons: effect of chronic administration of olanzapine. *Brain Res* **1025**: 194–202.
- Manna SK, Zhang HJ, Yan T, Oberley LW, Aggarwal BB (1998). Overexpression of manganese superoxide dismutase suppresses tumor necrosis factor-induced apoptosis and activation of nuclear transcription factor-kappaB and activated protein-1. *J Biol Chem* **273**: 13245–13254.
- Matias WG, Traore A, Bonini M, Sanni A, Creppy EE (1999). Oxygen reactive radicals production in cell culture by okadaic acid and their implication in protein synthesis inhibition. *Hum Exp Toxicol* **18**: 634–639.
- Mizuno M, Yamada K, Olariu A, Nawa H, Nabeshima T (2000). Involvement of BDNF in spatial memory formation and maintenance in a radial arm maze test in rats. *J Neurosci* **20**: 7116–7121.
- Nowakowska E, Chodera A, Kus K (1999). Influence of olanzapine on cognitive functions and catalepsy in rats after single and chronic administration. *Pol J Pharmacol* **51**: 295–300.
- Nuydens R, Dispersyn G, Van Den Keiboom G, de Jong M, Connors R, Ramaekers F et al (2000). Bcl-2 protects against

- apoptosis-related microtubule alterations in neuronal cells. *Apoptosis* 5: 43–51.
- Olton DS, Becker JT, Handelmann GE (1979). Hippocampus, space, and memory. *Behav Brain Sci* 2: 313–365.
- Paxinos G, Watson C (1986). *The Rat Brain in Stereotaxic Coordinates*. Academic Press: London.
- Purdon SE, Jones BDW, Stip E, Labelle A, Addington D, David SR et al (2000). Neuropsychological change in early phase schizophrenia during 12 months of treatment with olanzapine, risperidone, or haloperidol. *Arch Gen Psychiatry* 57: 249–258.
- Qing H, Xu H, Wei Z, Gibson K, Li XM (2003). The ability of atypical antipsychotic drugs vs haloperidol to protect PC12 cells against MPP⁺-induced apoptosis. *Eur J Neurosci* 17: 1563–1570.
- Radecki DT, Brown LM, Martinez J, Teyler TJ (2005). BDNF protects against stress-induced impairments in spatial learning and memory and LTP. *Hippocampus* 15: 246–253.
- Ramirez-Munguia N, Vera G, Tapia R (2003). Epilepsy, neurodegeneration, and extracellular glutamate in the hippocampus of awake and anesthetized rats treated with okadaic acid. *Neurochem Res* 28: 1517–1524.
- Riordan FA, Foroni L, Hoffbrand AV, Mehta AB, Wickremasinghe RG. (1998). Okadaic acid-induced apoptosis of HL60 leukemia cells is preceded by destabilization of bcl-2 mRNA and down-regulation of bcl-2 protein. *FEBS Lett* 435: 195–198.
- Shirazi-Southall S, Rodriguez DE, Nomikos GG (2002). Effects of typical and atypical antipsychotics and receptor selective compounds on acetylcholine efflux in the hippocampus of the rat. *Neuropsychopharmacology* 26: 583–594.
- Sweatt JD (2004). Hippocampal function in cognition. *Psychopharmacology* 174: 99–110.
- Tian Q, Lin Z-Q, Wang X-C, Chen J, Wang Q, Gong C-X et al (2004). Injection of okadaic acid into the meynert nucleus basalis of rat brain induces decreased acetylcholine level and spatial memory deficit. *Neuroscience* 126: 277–284.
- Van Dam A-M, Bol JGJM, Binnekade R, Van Muiswinkel FL (1998). Acute or chronic administration of okadaic acid to rats induces brain damage rather than Alzheimer-like neuropathology. *Neuroscience* 85: 1333–1335.
- Waddington JL, O'Callaghan E, Buckley P, Larkin C, Redmond O, Stack J et al (1991). The age differences of MRI abnormalities in schizophrenia suggest early ventricular enlargement but later prominence of cortical atrophy. *Schizophr Res* 5: 188–189.
- Wang C, Kaufmann JA, Sanchez-Ross MG, Johnson KM (2000). Mechanisms of N-methyl-D-aspartate-induced apoptosis in phencyclidine-treated cultured forebrain neurons. *J Pharmacol Exp Ther* 294: 287–295.
- Wei Z, Bai O, Richardson JS, Mousseau DD, Li X-M (2003a). Olanzapine protects PC12 cells from oxidative stress induced by hydrogen peroxide. *J Neurosci Res* 73: 364–368.
- Wei Z, Mousseau DD, Richardson JS, Dyck LE, Li X-M (2003b). Atypical antipsychotics attenuate neurotoxicity of β -amyloid (25–35) by modulating Bax and Bcl-X_{L/S} expression and localization. *J Neurosci Res* 74: 942–947.
- Wolff MC, Leander JD (2003). Comparison of the effects of antipsychotics on a delayed radial maze task in the rat. *Psychopharmacology* 168: 410–416.
- Woods BT, Yurgelun-Todd D, Benes FM, Frankenburg FR, Pope HG, McSparren J (1990). Progressive ventricular enlargement in schizophrenia: comparison to bipolar disorder and correlation with clinical course. *Biol Psychiatry* 27: 341–352.
- Zhang X, Cui SS, Wallace AE, Hannesson DK, Schmued LC, Saucier DM et al (2002). Relations between brain pathology and temporal lobe epilepsy. *J Neurosci* 22: 6052–6061.
- Zhao WQ, Bennett P, Sedman GL (1995). The impairment of long-term memory formation by the phosphatase inhibitor okadaic acid. *Brain Res Bull* 36: 557–561.
- Zou L-B, Yamada K, Tanaka K, Kameyama T, Nabeshima T (1998). Nitric oxide synthase inhibitors impair reference memory formation in a radial arm maze task in rats. *Neuropharmacology* 37: 323–330.