

# Differential Effects of $\sigma_1$ Receptor Blockade on Self-Administration and Conditioned Reinstatement Motivated by Cocaine vs Natural Reward

Rémi Martin-Fardon<sup>\*1</sup>, Tanguy Maurice<sup>2</sup>, Harinder Aujla<sup>1</sup>, Wayne D Bowen<sup>3</sup> and Friedbert Weiss<sup>1</sup>

<sup>1</sup>Molecular and Integrative Neurosciences Department, The Scripps Research Institute, La Jolla, CA, USA; <sup>2</sup>INSERM U. 710, EPHE, University of Montpellier II, Montpellier, France; <sup>3</sup>Department of Molecular Pharmacology, Physiology and Biotechnology, Brown University, Providence, RI, USA

Growing evidence suggests a role for sigma<sub>1</sub> ( $\sigma_1$ ) receptors in cognitive function, anxiety, depression, regulation of stress responses, and, recently, the appetitive effects of cocaine as measured by conditioned place preference. This study was designed to extend understanding of the role of  $\sigma_1$  receptors in addiction-relevant conditioned effects of cocaine by testing the effects of a potent and selective  $\sigma_1$  receptor antagonist, BD1047, on conditioned reinstatement of cocaine-seeking. To determine whether modification of conditioned reinstatement by BD1047 is selective for drug-directed behavior or reflects general suppressant effects on motivated behavior, BD1047 was tested also on reinstatement induced by stimuli conditioned to a natural reward, sweetened condensed milk (SCM). Additionally, because  $\sigma_1$  receptors have been implicated also in processes linked to the acute reinforcing actions of cocaine, tests of the effects of BD1047 on cocaine self-administration—including a comparison with the  $\sigma_1$  antagonist effects on SCM self-administration—were conducted as well. Cocaine self-administering male Wistar rats were trained to associate a discriminative stimulus ( $S^D$ ) with the availability of cocaine or SCM, and then subjected to reinstatement tests following extinction of cocaine or SCM-reinforced behavior. BD1047 (1–30 mg/kg) reversed response reinstatement induced by the cocaine  $S^D$  at 20 and 30 mg/kg but did not modify SCM  $S^D$ -induced responding at all but the highest 30 mg dose, at which responding was reversed to extinction levels. BD1047 did not modify responding reinforced directly by SCM or cocaine. The findings support a role for  $\sigma_1$  receptors in regulating conditioned responses to cocaine-related contextual stimuli and identify this receptor as a potential treatment target for the prevention of craving and relapse.

*Neuropsychopharmacology* (2007) **32**, 1967–1973; doi:10.1038/sj.npp.1301323; published online 31 January 2007

**Keywords:** addiction; BD1047; self-administration; drug-seeking behavior; relapse; drug cues

## INTRODUCTION

The sigma<sub>1</sub> ( $\sigma_1$ ) receptor is a 223-amino-acid protein localized in several organs, such as liver, heart, testis, gastrointestinal tract, and brain (Maurice *et al*, 2002). Immunohistochemical studies confirmed earlier *in vitro* binding assays indicating that  $\sigma_1$  receptors are present in numerous brain structures, with particularly high concentrations in specific areas within the limbic system as well as brainstem motor structures (for review, see Maurice *et al*, 2002). The highest levels of  $\sigma_1$  receptor immunostaining have been observed in the granular layer of the olfactory bulb, hypothalamic nuclei, hippocampus, and pyramidal layers of the hippocampus, caudate putamen, septum, nucleus accumbens, and amygdala (Alonso *et al*, 2000; but

see also Maurice *et al*, 2002). At the subcellular level, the  $\sigma_1$  receptor is present in neurons, on the endoplasmic reticulum as well as mitochondrial, nuclear, and plasma membranes (Maurice *et al*, 2002; Su and Hayashi, 2003). Physiologically,  $\sigma_1$  receptor activation leads to cellular restructuring by translocating cholesterol and cytoskeletal proteins from the endoplasmic reticulum to the plasma membrane and nucleus, and regulates synaptic transmission by modulating intracellular  $Ca^{2+}$  mobilization (Hayashi and Su, 2001, 2003a,b; Monnet *et al*, 2003). The endogenous ligand of  $\sigma_1$  receptors has not been fully identified yet. However, several lines of evidence suggest that neuroactive steroids such as dehydroepiandrosterone (DHEA) and progesterone act as potent endogenous  $\sigma_1$  receptor modulators (for review, see Monnet and Maurice, 2006). Behaviorally,  $\sigma_1$  receptors have been implicated in cognitive function, anxiety, depression, and regulation of stress responses (eg Maurice *et al*, 2001; Urani *et al*, 2001). Additionally,  $\sigma_1$  receptors have been found to modulate several neurobehavioral effects of cocaine, including the drug's subjective (Katz *et al*, 2003), psychomotor stimulant (Menkel *et al*, 1991; Ujike *et al*, 1996), rewarding (Romieu

\*Correspondence: Dr R Martin-Fardon, Molecular and Integrative Neurosciences Department, The Scripps Research Institute, SP30-2120, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA, Tel: +1 858 784 7154, Fax: +1 858 784 7146, E-mail: rmartinf@scripps.edu  
Received 1 August 2006; revised 3 November 2006; accepted 29 November 2006

*et al.*, 2000), and toxic (Matsumoto *et al.*, 2001) actions. Of interest with regard to a  $\sigma_1$  receptor role in cocaine addiction is that pharmacological blockade of this receptor attenuates expression of cocaine-conditioned place preference (CPP) (Romieu *et al.*, 2000, 2002), suggesting that  $\sigma_1$  receptors participate in mediating the conditioned incentive effects of cocaine-related environmental stimuli. It is well established that such stimuli can evoke craving or lead to relapse in abstinent individuals (eg O'Brien *et al.*, 1998) and, in animals, consistently elicit reinstatement of extinguished cocaine-seeking (See, 2002; Shaham *et al.*, 2003; Weiss, 2005).

To extend our understanding of the role of  $\sigma_1$  receptors in addiction-relevant conditioned effects of cocaine, the effects of a potent selective  $\sigma_1$  receptor antagonist, BD1047 (Matsumoto *et al.*, 1995; McCracken *et al.*, 1999b), were examined on reinstatement of cocaine-seeking induced by drug-related contextual stimuli. To establish whether BD1047 preferentially modifies drug-directed behavior or exerts general suppressant effects on motivated behavior, BD1047 effects were tested also on responding induced by stimuli conditioned to a potent conventional reinforcer, sweetened condensed milk (SCM). Because  $\sigma_1$  receptors have been implicated not only in the expression but also acquisition (Romieu *et al.*, 2000, 2002) of cocaine CPP (ie a process linked to the acute reinforcing actions of cocaine), tests of BD1047 effects on self-administration of cocaine and SCM were conducted as well.

## MATERIALS AND METHODS

### Subjects

Eighty-seven male Wistar rats (Charles River, Wilmington, MA; 200–250 g upon arrival) were housed 2–3/cage in a temperature- and humidity-controlled vivarium on a reverse 12:12 h light/dark cycle with *ad libitum* access to food and water. All procedures were conducted in strict adherence to the National Institutes of Health Guide for the Care and Use of Laboratory Animals and approved by the Institutional Animal Care and Use Committee of The Scripps Research Institute.

### Drugs

Cocaine hydrochloride (National Institute on Drug Abuse, Bethesda, MD) was dissolved in sterile physiological saline. Cocaine or saline vehicle was intravenously (i.v.) infused at a volume of 0.1 ml over 4 s. *N*-[2-(3,4-dichlorophenyl)ethyl]-*N*-methyl-2-(dimethylamino)ethylamine (BD1047), obtained from Dr WD Bowen (Brown University, Providence, RI), was dissolved in distilled water and intraperitoneally (i.p.) administered in a volume of 1 ml/kg.

### Effects of BD1047 on Conditioned Reinstatement: Procedures

Behavioral training and testing were conducted according to previously described procedures (Baptista *et al.*, 2004; Caine *et al.*, 1993; Weiss *et al.*, 2000). Briefly, operant responding maintained by food or SCM was initially established using a continuous reinforcement schedule in daily 60- and 40-min sessions, respectively. Rats designated for cocaine self-

administration were surgically prepared with jugular catheters (Caine *et al.*, 1993; Weiss *et al.*, 2000) and given 7 days of recovery before commencing self-administration training. Self-administration of cocaine (0.25 mg/0.1 ml; i.v.) or SCM (2:1 v/v in distilled water; 0.1 ml delivered into a 0.2 ml receptacle) began on a fixed ratio 1 (FR1) schedule of reinforcement in daily 120-min cocaine or 40-min SCM sessions, 5 days/week. Responses at the right, active lever were reinforced and followed by a 20-s time-out (TO) period signaled by illumination of a cue light above the lever. During this time, the lever remained inactive. Responses at the left, inactive lever had no scheduled consequences.

Following 2 weeks of training, responses at the active lever were differentially reinforced in the presence of distinct discriminative stimuli ( $S^D$ ). A constant 70 dB white noise ( $S^+$ ) signaled availability of the reinforcer whereas illumination of a 2.8 W house light ( $S^-$ ), located at the top of the chamber's front panel, signaled the absence of the reinforcer (ie, saline solution instead of cocaine or no consequence instead of SCM; see Baptista *et al.*, 2004). Three daily sessions (lasting 1 h for the cocaine group, and restricted to 20 min for the SCM group to avoid satiety effects by excessive ingestion of SCM), separated by 30-min intervals, were conducted, with two 'reward' sessions and one 'non-reward' session sequenced in random order. Sessions were initiated by presentation of the respective  $S^D$  and extension of the levers. The  $S^D$  remained present until termination of the session by retraction of the levers. After eight training days (ie, a total of 16 'reward' and 8 'non-reward' sessions), both the cocaine and SCM groups were placed on extinction conditions in daily 1-h sessions during which the reinforcers and  $S^D$  were withheld until a criterion of  $\leq 4$  responses/session for 3 consecutive days was reached. One day after each animal reached the extinction criterion, reinstatement tests began. These 1-h tests were conducted under extinction conditions, but with reintroduction of the  $S^D$  as during the conditioning phase. Rats were tested first in the presence of the  $S^-$  to verify the behavioral selectivity of the  $S^D$ . Two days later, tests of BD1047 (0, 1, 3, 10, 20, or 30 mg/kg, i.p.) on  $S^+$ -induced reinstatement began. BD1047 was administered 10 min before the onset of sessions. Each animal was tested with only one dose of BD1047 according to a between-subjects design.

### Effect of BD1047 on Self-Administration: Procedures

After acquisition of cocaine- or SCM-reinforced responding on an FR1 schedule as above, but without being subjected to conditioning procedures, rats received further daily 120-min access to cocaine on an FR 5 schedule or 30-min access to SCM on an FR1 schedule until stable intake developed ( $\pm 10\%$ ). Effects of BD1047 (0, 1, 3, 10, 20, and 30 mg/kg, i.p.) on cocaine or SCM self-administration were then determined. Each rat was tested once with each BD1047 dose on separate days according to a Latin square design ( $n = 6$  animals/group). Each drug test was preceded by five baseline cocaine or SCM self-administration sessions.

### Statistics

Differences in responding at the active lever between the respective reward and non-reward conditions during the

training phase were analyzed by paired *t*-tests. Differences in the number of responses during the extinction and reinstatement phases, including the effects of BD1047 on reinstatement responses, were analyzed separately for the cocaine and SCM groups by one-way ANOVA. Cumulative responses were analyzed by mixed-factorial ANOVA, followed by simple effects analysis. Effects of BD1047 on responses reinforced directly by cocaine or SCM were analyzed separately by one-way within-subjects ANOVA. Significant omnibus tests were followed by Fisher's PLSD *post hoc* tests.

## RESULTS

### Effect of BD1047 on Conditioned Reinstatement

**Cocaine.** All rats ( $n=33$ ) acquired cocaine-reinforced responding, maintained stable cocaine self-administration during the conditioning phase, and ceased responding during non-reward (saline) sessions (Figure 1a, left panel). Following initiation of the extinction contingency, rats required, on average,  $9.6 \pm 1.3$  (mean  $\pm$  SEM) sessions to reach the criterion. During subsequent reinstatement tests, the cocaine  $S^+$  (in BD1047 vehicle-treated rats) but not the saline-associated  $S^-$  elicited strong recovery of responding ( $p < 0.001$ , Fisher PLSD tests after ANOVA:  $F_{2,8} = 31.2$ ;  $p < 0.001$ ; Figure 1a). BD1047 dose-dependently reduced the response reinstatement induced by the  $S^+$  (Figure 1a, right panel) with significant effects at the 20 ( $p < 0.05$ ) and 30 mg/kg ( $p < 0.01$ ) doses (Fisher PLSD tests after ANOVA:  $F_{5,27} = 3.2$ ;  $p < 0.05$ ). Further examination of the BD1047 effects revealed that this agent modified the cumulative response profile associated with  $S^+$ -induced reinstatement (Figure 1a, inset) as reflected by a main effect for BD1047 doses ( $F_{5,27} = 4.2$ ;  $p < 0.01$ ) and a dose  $\times$  time (10 min intervals) interaction ( $F_{25,135} = 1.9$ ;  $p < 0.01$ ). Moreover, this analysis indicated that only the 20 mg/kg and 30 mg/kg BD1047 doses decreased responding at all time points compared to vehicle-treated rats (simple effects,  $p < 0.05$ ; Figure 1a, inset). Inactive lever responses remained low ( $\leq 4$  responses) throughout the experiment and unaltered by BD1047.

**Sweetened condensed milk.** As in the case of cocaine, all rats ( $n=42$ ) acquired robust SCM-reinforced responding, maintained stable responding during the conditioning phase, and ceased responding during non-reward sessions (Figure 1b, left panel). Rats required on average  $10.3 \pm 0.7$  (mean  $\pm$  SEM) sessions to reach the extinction criterion. Presentation of the SCM  $S^+$  (in vehicle-injected rats), but not the  $S^-$ , produced reliable reinstatement ( $p < 0.001$ , Fisher PLSD tests after ANOVA:  $F_{2,12} = 53.3$ ;  $p < 0.001$ ; Figure 1b, right panel). Only the highest dose of BD1047 attenuated the  $S^+$ -induced response reinstatement ( $p < 0.01$ , Fisher PLSD tests after ANOVA:  $F_{5,36} = 2.6$ ;  $p < 0.05$ ; Figure 1b). Examination of BD1047's effects on cumulative responses confirmed that the  $\sigma_1$  antagonist modified the shape of the cumulative response profile (main effect of BD1047 dose:  $F_{5,36} = 3.5$ ;  $p < 0.05$ ), but only at the highest dose (Figure 1b, inset). The failure to obtain a significant dose  $\times$  time interaction reflects the suppressant effects on reinstatement of only a single (ie the 30 mg/kg) dose that persisted across the entire session

(Figure 1b, inset). Responses at the inactive lever remained low (4 responses) throughout training and testing and were not modified by BD1047.

### Effect of BD1047 on Cocaine and SCM-Reinforced Behavior


All rats acquired stable cocaine ( $n=6$ ) or SCM self-administration ( $n=6$ ) after 14 days of training. BD1047 did not alter either cocaine ( $F_{5,25} = 1.9$ ;  $p > 0.05$ ; Figure 2a) or SCM-reinforced responding ( $F_{5,25} = 1.7$ ;  $p > 0.05$ ; Figure 2b). Responses at the inactive lever remained low ( $\leq 2$  responses) throughout testing, and were not modified by BD1047.

## DISCUSSION

BD1047 dose-dependently reduced reinstatement induced by a cocaine-related contextual stimulus, whereas responding elicited by a stimulus conditioned to a palatable conventional reinforcer, SCM, was attenuated at the highest BD1047 dose only. The  $\sigma_1$  antagonist did not modify self-administration of cocaine or SCM. This observation suggests that pharmacological blockade of  $\sigma_1$  receptors does not modify the acute reinforcing effects of cocaine or palatable natural reward, but selectively attenuates the incentive-motivational effects of reward-paired contextual stimuli with a preferential action on stimuli conditioned to cocaine *vs* natural reward.

The preferential interference by BD1047 with cocaine  $S^+$ -induced reinstatement resembles earlier observations where a Group II metabotropic glutamate receptor agonist (LY379268) selectively attenuated reinstatement induced by stimuli conditioned to cocaine *vs* conventional reinforcers including food and SCM (Baptista *et al*, 2004). This effect of BD1047 is unlikely to be the result of differences in response-reinstating efficacy between the cocaine and SCM  $S^+$  *per se*. While SCM conditioning sessions were restricted to 20 min to avoid satiety and, thus, shorter than cocaine conditioning sessions, the number of responses per session was considerably higher in the SCM than in the cocaine group such that these animals experienced a greater number of reinforcer presentations in the presence of the SCM  $S^+$  compared to the cocaine  $S^+$  condition. Nonetheless, as in previous reports where stimuli conditioned to conventional reinforcers were less effective in eliciting reward-seeking than cocaine-associated stimuli (Baptista *et al*, 2004; Grimm *et al*, 2002), the SCM  $S^+$  produced weaker reinstatement than the cocaine  $S^+$ . Moreover, the SCM  $S^+$  induced and maintained responding only during the first 10–20 min of the reinstatement test, whereas the cocaine  $S^+$  produced strong and sustained responding throughout the 60 min session.


BD1047 produced dose-dependent effects on reinstatement only in the case of the cocaine  $S^+$ . However, at the highest dose, reinstatement elicited by the SCM  $S^+$  was significantly suppressed as well. Further scrutiny of the time course of responding during the reinstatement tests (Figure 1a and b, insets) shows that the highest doses of BD1047 induced a delayed onset of responding. Specifically, the effects of the 20 mg/kg dose appeared to dissipate 30 min into the test (Figure 1a), whereas the effects of the 30 mg/kg dose did not show signs of dissipation until 50–60 min after the beginning of the test (see insets of Figure 1a and b).



**Figure 1** Effects of BD1047 on reinstatement induced by discriminative stimuli associated with (a) cocaine and (b) SCM. (a) Left panels: active lever responses during conditioning sessions in the presence of stimuli paired with cocaine (upper panel: COC/S<sup>+</sup>) or SCM (lower panel: SCM/S<sup>+</sup>) vs non-availability of these reinforcers (upper panel: SAL/S<sup>-</sup>; lower panel: non-SCM/S<sup>-</sup>). Center panels: Extinction (EXT) responses at criterion and responses during an initial reinstatement test in the presence of the stimulus paired with reward non-availability (S<sup>-</sup>). Right panels: reinstatement responses in the presence of the stimuli previously associated with reward (cocaine: upper panel; SCM: lower panel) availability (S<sup>+</sup>) in vehicle-treated rats (0), and modification of conditioned reinstatement across doses of BD1047. Insets: cumulative number of responses throughout the 60-min reinstatement periods (error bars omitted for clarity). Upper panels: \*\*\*paired *t*-test  $t_{32} = 14.3$ ;  $p < 0.001$  vs COC/S<sup>+</sup>; & $p < 0.001$  vs S<sup>+</sup>; \* $p < 0.05$  and \*\* $p < 0.01$  vs vehicle; +  $p < 0.05$  and ++  $p < 0.01$  vs BD1047 1 mg/kg. Lower panels: \*\*\*paired *t*-test  $t_{42} = 91.1$ ;  $p < 0.001$  vs SCM/S<sup>+</sup>; & $p < 0.001$  vs S<sup>+</sup>; \*\* $p < 0.01$  vs vehicle; ++  $p < 0.01$  vs BD1047 1 mg/kg. Numbers inside bars represent sample sizes.

The pharmacokinetic profile of BD1047 has not yet been precisely described. However, these observations in conjunction with earlier data (Matsumoto *et al*, 1995; Romieu *et al*, 2004; Urani *et al*, 2001) strongly suggest that the behavioral effects of BD1047 last for at least 30 min following administration and that BD1047's duration of action may vary as a function of dose. The reason for the failure of BD1047 to alter SCM S<sup>+</sup>-induced behavior at low doses, paired with the full suppression of this behavior only at the highest dose is presently unclear. This effect cannot be

attributed to motor impairment or sedation, because BD1047 did not interfere, even at the highest dose, with the high rate of responding in rats self-administering SCM (Figure 2b), confirming previous reports that  $\sigma_1$  receptor antagonists are devoid of general suppressant effects on behavior (McCracken *et al*, 1999a; Romieu *et al*, 2006). In particular, while BD1047 reduces the acute locomotor stimulant effects of cocaine as well as cocaine-induced convulsions and lethality, the  $\sigma_1$  receptor antagonist does not interfere with spontaneous locomotor activity (McCracken *et al*, 1999a).



**Figure 2** Effects of BD1047 on self-administration of cocaine or SCM. Cocaine (a) and SCM- (b) reinforced responses after vehicle vs BD1047 administration.

As BD1047 reliably attenuated conditioned cocaine-seeking, one might expect that this drug would exert at least similar, if not greater, effects on behavior that is less robust (ie conditioned SCM-seeking). One interpretation of this finding is that  $\sigma_1$  receptors participate in mediating behavior motivated by drug-related but not non-drug-related stimuli and therefore, possibly, point toward the existence of separate neural substrates for drug-related vs conventional learning. It is also possible that  $\sigma_1$  receptors participate in regulating conditioning processes independent of the nature of the unconditioned stimulus (ie cocaine vs SCM), but that the effects of non-drug cues are more resistant to disruption than the effects of drug cues. A more likely interpretation for the differential effects of BD1047 on cocaine  $S^+$  vs SCM  $S^+$ -induced reinstatement is related to upregulation of  $\sigma_1$  receptor function after cocaine treatment. Acute cocaine administration has been shown to upregulate  $\sigma_1$  receptor gene expression and protein levels in whole brain, striatum, and cortex (Liu *et al*, 2005). Moreover, 4 days of intermittent cocaine exposure during place conditioning increased *in vivo* binding levels of the  $\sigma_1$  receptor agonist [<sup>3</sup>H](+)-SKF-10,047 in the olfactory bulb, hippocampus, hypothalamus, cortex, and striatum—a change that was sustained during extinction of CPP—indicative of persistent functional  $\sigma_1$  receptor upregulation (Romieu *et al*, 2004). It is possible then that over the course of cocaine self-administration and cue conditioning, followed by a prolonged extinction period, sustained upregulation of  $\sigma_1$  receptor function may have occurred, an effect that would not be expected to develop in the cocaine-naïve SCM group. Consequently, increases in  $\sigma_1$  receptor expression or affinity may have amplified the effects of BD1047 in the cocaine-exposed but not drug-naïve SCM group, providing a possible explanation for the differential efficacy of BD1047 in reversing behavior induced by the cocaine vs SCM  $S^+$ . Several endogenous systems have been shown to interact with the  $\sigma_1$  receptor, including peptides of the neuropeptide Y and calcitonin gene-related peptide families and neuroactive steroids that have attracted much interest recently as putative endogenous ligands for the  $\sigma_1$  receptor (for reviews, see Maurice, 2004; Monnet and Maurice, 2006). Specifically, it has been shown that pregnenolone and DHEA act as  $\sigma_1$  receptor agonists, progesterone being the most potent endogenous  $\sigma_1$  receptor antagonist known to date (Maurice, 2004; Maurice *et al*, 2002). These neuroactive steroids have been shown to modulate cocaine-induced locomotor stimulation or CPP through a direct interaction with  $\sigma_1$  receptors (Romieu *et al*, 2003), and *in vivo* assay also showed that

circulating levels of cortisol and DHEA sulfate were increased following days of abstinence in cocaine addicts (Buydens-Branchey and Branchey, 2004; Buydens-Branchey *et al*, 2002). Finally, studies in rats have confirmed that DHEA attenuated cocaine self-administration and cocaine-seeking, an effect possibly resulting from an interaction with the  $\sigma_1$  receptor (Doron *et al*, 2006; Maayan *et al*, 2006). On the basis of these findings, it would seem justified to speculate that cocaine during self-administration training and conditioning produced persistent neuroendocrine perturbations, which could in turn have resulted in upregulation of  $\sigma_1$  receptors increasing susceptibility to relapse (ie, conditioned reinstatement) in the cocaine group.

BD1047 did not modify the primary reinforcing effects of cocaine and SCM at a dose range that progressively reduced the conditioned effects of the cocaine  $S^+$ . Sigma<sub>1</sub> receptors are highly expressed in the hippocampus (Alonso *et al*, 2000), a brain site with an established role in the occasion-setting action of contextual stimuli (eg Holland and Bouton, 1999). Indeed, transient inactivation of the dorsal hippocampus was shown to specifically block contextual reinstatement of cocaine-seeking, but not reinstatement induced by cocaine or discrete cocaine-paired cues (Fuchs *et al*, 2005). Therefore, one may speculate that, in the present study, the  $\sigma_1$  receptor antagonist interfered with the processing of reward-related contextual information at the hippocampal level, reducing its motivating impact, an effect that would not interfere with the direct reinforcing actions of drug or natural reward. It is important to note, however, that BD1047 has been reported to also block cocaine priming-induced reactivation of cocaine CPP (Romieu *et al*, 2004)—an effect thought to be mediated by the nucleus accumbens shell and the ventral tegmental area (see Kalivas and McFarland, 2003 for review)—suggesting that  $\sigma_1$  receptors in the nucleus accumbens and the ventral tegmental area may play a role in reinstatement induced by cocaine priming manipulations (Alonso *et al*, 2000; Maurice *et al*, 2002; McFarland *et al*, 2003). However, a more comprehensive understanding of the pharmacological profile of BD1047 relevant for reinstatement and relapse, including its effects on priming-induced reinstatement as well as the assessment of specific links between effects of  $\sigma_1$  manipulations in anatomically distinct brain regions on cocaine-seeking associated with distinct risk factors (ie contextual vs discrete cocaine cues vs cocaine priming vs stress exposure) will remain for further research.

The lack of BD1047 effect on cocaine self-administration observed here is in apparent contradiction to earlier findings showing that BD1047 blocks the acquisition of

cocaine CPP, suggesting that the drug can antagonize the acute reinforcing actions of cocaine (Romieu *et al*, 2000, 2002). Several explanations can be offered for this discrepancy. First, acquisition of cocaine CPP was obtained with involuntary cocaine administration rather than self-administration. It is well established that the behavioral and neurochemical effects of voluntary *vs* involuntary cocaine administration differ (eg Hemby *et al*, 1997; Jacobs *et al*, 2003; Wise, 2000). Moreover, the duration and amount of cocaine exposure in the self-administering rats of the present study was considerably greater compared to that in the CPP studies. Therefore, both the reinforcing quality and relevant neuroadaptive changes are likely to differ in rats subjected to involuntary *vs* self-administration of cocaine. Consistent with this hypothesis is recent evidence on differential regulation of the  $\sigma_1$  receptor as a function of the mode of administration of methamphetamine (active *vs* passive). Rats self-administering methamphetamine showed increased expression of  $\sigma_1$  receptor mRNA in the hippocampus and decreased expression in the frontal cortex compared to yoked-methamphetamine and yoked-saline controls (Stefanski *et al*, 2004). Second, rats in the present study self-administered cocaine for 2 h after receiving BD1047. Therefore, the effects of BD1047 may have been surmountable by continued response-contingent administration of cocaine in the present case, but not in the CPP studies in which rats received a single i.p. cocaine dose before conditioning sessions. A possible alternative explanation for these discrepant findings may be that self-administration and CPP procedures also differ with respect to the significance of contextual learning in the acquisition of cocaine-reinforced behavior. The contextual associative learning component relevant for CPP through which environmental stimuli in the drug-paired environment eventually establish secondary reinforcing properties is less critical in the establishment of drug-reinforced behavior in self-administration studies. In fact, self-administration depends on intact operant learning abilities of the organism (ie animals learn the association between a response and an outcome rather than between a previously neutral set of environmental stimuli and drug-induced internal state). According to this possibility, BD1047 may be effective in disrupting the acquisition of associations between previously neutral cues now paired with cocaine, but does not disrupt the maintenance of response–outcome behavior. A comparative study testing the effects of chronic BD1047 administration during the induction of CPP *vs* acquisition of cocaine self-administration will be needed to pursue this hypothesis.

What may constrain interpretation of the findings in terms of a specific role of  $\sigma_1$  receptors in conditioned cocaine reinstatement is that only a single  $\sigma_1$  antagonist was tested. However, few  $\sigma_1$ -selective ligands are presently available, and BD1047, which has nanomolar affinity for the  $\sigma_1$  receptor, is the most selective and potent  $\sigma_1$  receptor antagonist described to date (Daniels *et al*, 2006; Matsu-moto *et al*, 1995; McCracken *et al*, 1999a,b). Moreover, although BD1047 also has moderate affinity for  $\sigma_2$  receptors and  $\beta$ -adrenergic receptors, it shows substantial selectivity for  $\sigma_1$  receptors ( $\approx 50$  times over  $\sigma_2$  receptors, and  $\approx 150$  times over  $\beta$ -adrenergic receptors), providing confidence that, at the dose range used, the behavioral effects obtained

here are attributable to an action at  $\sigma_1$  receptors (Matsuo-moto *et al*, 1995). In addition, evidence exists that reduction of  $\sigma_1$  receptor expression following antisense treatment prevents the acquisition of cocaine CPP, a finding that directly implicates  $\sigma_1$  receptors in contextual cocaine conditioning (Romieu *et al*, 2000). Moreover, BD1047 produced effects on CPP highly similar to those of  $\sigma_1$  antisense treatment (Romieu *et al*, 2000). Together, these observations provide strong support for the interpretation that the present findings resulted from an action of BD1047 at  $\sigma_1$  receptors.

In summary, the selective  $\sigma_1$  receptor antagonist, BD1047, attenuated conditioned reinstatement by reward-paired contextual stimuli with a preferential action on stimuli conditioned to cocaine *vs* natural reward. BD1047 had little effect on consummatory behavior maintained by drug or natural reward. The presumably selective  $\sigma_1$  receptor antagonist action of BD1047, therefore, appears to attenuate the motivating effects of cocaine cues without interfering with normal motivational function. Thus, the results identify BD1047 as a promising agent for further scrutiny with regard to therapeutic potential and, by inference, that  $\sigma_1$  receptors may represent a promising novel target for the prevention of cocaine craving and relapse.

#### ACKNOWLEDGEMENTS

This is publication number 17316-NP from The Scripps Research Institute. This research was supported by NIH/NIDA Grants DA07348 and DA08467 (FW). We thank ND Stuempfig for technical assistance, M Arends for help with manuscript preparation, and Dr P Romieu (INSERM U. 575, Strasbourg, France) for constructive suggestions.

#### REFERENCES

- Alonso G, Phan V, Guillemain I, Saunier M, Legrand A, Anoaï M *et al* (2000). Immunocytochemical localization of the sigma(1) receptor in the adult rat central nervous system. *Neuroscience* **97**: 155–170.
- Baptista MA, Martin-Fardon R, Weiss F (2004). Preferential effects of the metabotropic glutamate 2/3 receptor agonist LY379268 on conditioned reinstatement versus primary reinforcement: comparison between cocaine and a potent conventional reinforcer. *J Neurosci* **24**: 4723–4727.
- Buydens-Branchey L, Branchey M (2004). Cocaine addicts with conduct disorder are typified by decreased cortisol responsivity and high plasma levels of DHEA-S. *Neuropsychobiology* **50**: 161–166.
- Buydens-Branchey L, Branchey M, Hudson J, Dorota Majewska M (2002). Perturbations of plasma cortisol and DHEA-S following discontinuation of cocaine use in cocaine addicts. *Psychoneuroendocrinology* **27**: 83–97.
- Caine SB, Lintz R, Koob GF (1993). Intravenous drug self-administration techniques in animals. In: Sahgal A (ed). *Behavioral Neuroscience: A Practical Approach*, Vol. 2. University Press: Oxford. pp 117–143.
- Daniels A, Ayala E, Chen W, Coop A, Matsumoto RR (2006). *N*-[2-(*m*-methoxyphenyl)ethyl]-*N*-ethyl-2-(1-pyrrolidinyl)ethylamine (UMB 116) is a novel antagonist for cocaine-induced effects. *Eur J Pharmacol*, print copy in press (originally published online April 5, 2006, at [www.sciencedirect.com](http://www.sciencedirect.com)).
- Doron R, Fridman L, Gispán-Herman I, Maayan R, Weizman A, Yadid G (2006). DHEA, a neurosteroid, decreases cocaine

- self-administration and reinstatement of cocaine-seeking behavior in rats. *Neuropsychopharmacology* 31: 2231–2236.
- Fuchs RA, Evans KA, Ledford CC, Parker MP, Case JM, Mehta RH et al (2005). The role of the dorsomedial prefrontal cortex, basolateral amygdala, and dorsal hippocampus in contextual reinstatement of cocaine seeking in rats. *Neuropsychopharmacology* 30: 296–309.
- Grimm JW, Shaham Y, Hope BT (2002). Effect of cocaine and sucrose withdrawal period on extinction behavior, cue-induced reinstatement, and protein levels of the dopamine transporter and tyrosine hydroxylase in limbic and cortical areas in rats. *Behav Pharmacol* 13: 379–388.
- Hayashi T, Su TP (2001). Regulating ankyrin dynamics: roles of sigma-1 receptors. *Proc Natl Acad Sci USA* 98: 491–496.
- Hayashi T, Su TP (2003a). Intracellular dynamics of sigma-1 receptors (sigma(1) binding sites) in NG108-15 cells. *J Pharmacol Exp Ther* 306: 726–733.
- Hayashi T, Su TP (2003b). Sigma-1 receptors (sigma(1) binding sites) form raft-like microdomains and target lipid droplets on the endoplasmic reticulum: roles in endoplasmic reticulum lipid compartmentalization and export. *J Pharmacol Exp Ther* 306: 718–725.
- Hemby SE, Co C, Koves TR, Smith JE, Dworkin SI (1997). Differences in extracellular dopamine concentrations in the nucleus accumbens during response-dependent and response-independent cocaine administration in the rat. *Psychopharmacology* 133: 7–16.
- Holland PC, Bouton ME (1999). Hippocampus and context in classical conditioning. *Curr Opin Neurobiol* 9: 195–202.
- Jacobs EH, Smit AB, de Vries TJ, Schoffelmeer AN (2003). Neuroadaptive effects of active versus passive drug administration in addiction research. *Trends Pharmacol Sci* 24: 566–573.
- Kalivas PW, McFarland K (2003). Brain circuitry and the reinstatement of cocaine-seeking behavior. *Psychopharmacology (Berl)* 168: 44–56.
- Katz JL, Libby TA, Kopajtic T, Husbands SM, Newman AH (2003). Behavioral effects of rimcazole analogues alone and in combination with cocaine. *Eur J Pharmacol* 468: 109–119.
- Liu Y, Chen GD, Lerner MR, Brackett DJ, Matsumoto RR (2005). Cocaine up-regulates Fra-2 and sigma-1 receptor gene and protein expression in brain regions involved in addiction and reward. *J Pharmacol Exp Ther* 314: 770–779.
- Maayan R, Lotan S, Doron R, Shabat-Simon M, Gispán-Herman I, Weizman A et al (2006). Dehydroepiandrosterone (DHEA) attenuates cocaine-seeking behavior in the self-administration model in rats. *Eur Neuropsychopharmacol* 16: 329–339.
- Matsumoto RR, Bowen WD, Tom MA, Vo VN, Truong DD, De Costa BR (1995). Characterization of two novel sigma receptor ligands: antidystonic effects in rats suggest sigma receptor antagonism. *Eur J Pharmacol* 280: 301–310.
- Matsumoto RR, McCracken KA, Pouw B, Miller J, Bowen WD, Williams W et al (2001). N-alkyl substituted analogs of the sigma receptor ligand BD1008 and traditional sigma receptor ligands affect cocaine-induced convulsions and lethality in mice. *Eur J Pharmacol* 411: 261–273.
- Maurice T (2004). Neurosteroids and sigma1 receptors, biochemical and behavioral relevance. *Pharmacopsychiatry* 37(Suppl 3): S171–S182.
- Maurice T, Martin-Fardon R, Romieu P, Matsumoto RR (2002). Sigma(1) (sigma(1)) receptor antagonists represent a new strategy against cocaine addiction and toxicity. *Neurosci Biobehav Rev* 26: 499–527.
- Maurice T, Phan VL, Privat A (2001). The anti-amnesic effects of sigma(1) (sigma(1)) receptor agonists confirmed by *in vivo* antisense strategy in the mouse. *Brain Res* 898: 113–121.
- McCracken KA, Bowen WD, de Costa BR, Matsumoto RR (1999a). Two novel sigma receptor ligands, BD1047 and LR172, attenuate cocaine-induced toxicity and locomotor activity. *Eur J Pharmacol* 370: 225–232.
- McCracken KA, Bowen WD, Matsumoto RR (1999b). Novel sigma receptor ligands attenuate the locomotor stimulatory effects of cocaine. *Eur J Pharmacol* 365: 35–38.
- McFarland K, Lapish CC, Kalivas PW (2003). Prefrontal glutamate release into the core of the nucleus accumbens mediates cocaine-induced reinstatement of drug-seeking behavior. *J Neurosci* 23: 3531–3537.
- Menkel M, Terry P, Pontecorvo M, Katz JL, Witkin JM (1991). Selective sigma ligands block stimulant effects of cocaine. *Eur J Pharmacol* 201: 251–252.
- Monnet FP, Maurice T (2006). The  $\sigma_1$  protein as a target for the non-genomic effects of neuro(active)steroids: molecular, physiological, and behavioral aspects. *J Pharmacol Sci* 100: 93–118.
- Monnet FP, Morin-Surun MP, Leger J, Combettes L (2003). Protein kinase C-dependent potentiation of intracellular calcium influx by sigma(1) receptor agonists in rat hippocampal neurons. *J Pharmacol Exp Ther* 307: 705–712.
- O'Brien CP, Childress AR, Ehrman R, Robbins SJ (1998). Conditioning factors in drug abuse: can they explain compulsion? *J Psychopharmacol* 12: 15–22.
- Romieu P, Lucas M, Maurice T (2006). Sigma(1) receptor ligands and related neuroactive steroids interfere with the cocaine-induced state of memory. *Neuropsychopharmacology* 31: 1431–1443.
- Romieu P, Martin-Fardon R, Bowen WD, Maurice T (2003). Sigma 1 receptor-related neuroactive steroids modulate cocaine-induced reward. *J Neurosci* 23: 3572–3576.
- Romieu P, Martin-Fardon R, Maurice T (2000). Involvement of the sigma(1) receptor in the cocaine-induced conditioned place preference. *Neuroreport* 11: 2885–2888.
- Romieu P, Meunier J, Garcia D, Zozime N, Martin-Fardon R, Bowen WD et al (2004). The sigma(1) (sigma(1)) receptor activation is a key step for the reactivation of cocaine conditioned place preference by drug priming. *Psychopharmacology (Berl)* 175: 154–162.
- Romieu P, Phan VL, Martin-Fardon R, Maurice T (2002). Involvement of the sigma(1) receptor in cocaine-induced conditioned place preference: possible dependence on dopamine uptake blockade. *Neuropsychopharmacology* 26: 444–455.
- See RE (2002). Neural substrates of conditioned-cued relapse to drug-seeking behavior. *Pharmacol Biochem Behav* 71: 517–529.
- Shaham Y, Shalev U, Lu L, De Wit H, Stewart J (2003). The reinstatement model of drug relapse: history, methodology and major findings. *Psychopharmacology (Berl)* 168: 3–20.
- Stefanski R, Justinova Z, Hayashi T, Takebayashi M, Goldberg SR, Su TP (2004). Sigma(1) receptor upregulation after chronic methamphetamine self-administration in rats: a study with yoked controls. *Psychopharmacology (Berl)* 175: 68–75.
- Su TP, Hayashi T (2003). Understanding the molecular mechanism of sigma-1 receptors: towards a hypothesis that sigma-1 receptors are intracellular amplifiers for signal transduction. *Curr Med Chem* 10: 2073–2080.
- Ujike H, Kuroda S, Otsuki S (1996). sigma Receptor antagonists block the development of sensitization to cocaine. *Eur J Pharmacol* 296: 123–128.
- Urani A, Roman FJ, Phan VL, Su TP, Maurice T (2001). The antidepressant-like effect induced by sigma(1)-receptor agonists and neuroactive steroids in mice submitted to the forced swimming test. *J Pharmacol Exp Ther* 298: 1269–1279.
- Weiss F (2005). Neurobiology of craving, conditioned reward and relapse. *Curr Opin Pharmacol* 5: 9–19.
- Weiss F, Maldonado-Vlaar CS, Parsons LH, Kerr TM, Smith DL, Ben-Shahar O (2000). Control of cocaine-seeking behavior by drug-associated stimuli in rats: effects on recovery of extinguished operant responding and extracellular dopamine levels in amygdala and nucleus accumbens. *Proc Natl Acad Sci USA* 97: 4321–4326.
- Wise RA (2000). Addiction becomes a brain disease. *Neuron* 26: 27–33.