

Interleukin-6 Upregulates Neuronal Adenosine A₁ Receptors: Implications for Neuromodulation and Neuroprotection

Knut Biber^{*1,6}, A Pinto-Duarte^{2,3,6}, MC Wittendorp¹, AM Dolga⁴, CC Fernandes^{2,3},
J Von Frijtag Drabbe Künzel⁵, JN Keijser⁴, R de Vries¹, AP Ijzerman⁵, JA Ribeiro^{2,3}, U Eisel⁴,
AM Sebastião^{2,3} and HWGM Boddeke¹

¹Department of Medical Physiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands; ²Institute of Pharmacology and Neurosciences, Faculty of Medicine, University of Lisbon, Lisbon, Portugal; ³Institute of Molecular Medicine, University of Lisbon, Lisbon, Portugal; ⁴Department of Molecular Neurobiology, University of Groningen, Groningen, The Netherlands; ⁵Leiden/Amsterdam Center for Drug Research, Gorlaeus Laboratories, Leiden, The Netherlands

The immunological response in the brain is crucial to overcome neuropathological events. Some inflammatory mediators, such as the immunoregulatory cytokine interleukin-6 (IL-6) affect neuromodulation and may also play protective roles against various noxious conditions. However, the fundamental mechanisms underlying the long-term effects of IL-6 in the brain remain unclear. We now report that IL-6 increases the expression and function of the neuronal adenosine A₁ receptor, with relevant consequences to synaptic transmission and neuroprotection. IL-6-induced amplification of A₁ receptor function enhances the responses to readily released adenosine during hypoxia, enables neuronal rescue from glutamate-induced death, and protects animals from chemically induced convulsing seizures. Taken together, these results suggest that IL-6 minimizes the consequences of excitotoxic episodes on brain function through the enhancement of endogenous adenosinergic signaling.

Neuropsychopharmacology (2008) **33**, 2237–2250; doi:10.1038/sj.npp.1301612; published online 7 November 2007

Keywords: brain inflammation; cytokines; excitotoxicity; seizures; hypoxia; adenosine A₁ receptor

INTRODUCTION

In the past years important functions were ascribed to cytokines within the central nervous system. Despite the constitutive expression of these mediators in the brain is usually low, their secretion increases dramatically in response to specific stimuli, such as infection or injury. Due to numerous reports showing the augmented expression of the immunoregulatory cytokine interleukin-6 (IL-6) in a variety of neurodegenerative diseases, this molecule is now recognized as a key player in brain pathology (Gadient and Otten, 1997; Gruol and Nelson, 1997). However, the role of IL-6 in that phenomenon has been regarded as a double-edged sword, since both neuroprotective and detrimental properties were associated to its expression. On one hand it has been shown that IL-6 protects cultured neurons (Carlson *et al*, 1999; Ali *et al*, 2000; Pizzi *et al*, 2004) and is beneficial in several models of neurodegenerative diseases

(Bensadoun *et al*, 2001; Bolin *et al*, 2002; Penkowa *et al*, 2001; Loddick *et al*, 1998; Ali *et al*, 2000; Swartz *et al*, 2001; Herrmann *et al*, 2003); on the other hand, chronic treatment of cultured neurons with IL-6 or chronic expression of IL-6 in astrocytes might result in neuronal death and neurodegeneration (Qiu *et al*, 1998; Nelson *et al*, 2004; Conroy *et al*, 2004; Campbell *et al*, 1993).

Despite the large body of data on the effects of IL-6 in brain, the molecular events that lead either to IL-6-dependent protection or death of neurons remain unclear. We have previously shown that IL-6 induces the upregulation of mRNA transcripts for the adenosine A₁ receptor in the brain (Biber *et al*, 2001). Functional adenosine A₁ receptors fulfill important neuroprotective properties during excitotoxic conditions, causing the inhibition of glutamate release from the presynaptic terminal (Barrie and Nicholls, 1993), the downregulation of postsynaptic NMDA receptor-mediated currents (de Mendonça *et al*, 1995; Sebastião *et al*, 2001) and the stabilization of the postsynaptic cell membrane (Trussell and Jackson, 1985; Gerber and Gahwiler, 1994). Despite the fact that adenosine A₁ receptor has been recurrently discussed as a potential target for the treatment of neurodegenerative diseases (Ribeiro, 2005), adenosine A₁ receptor agonists are not used with therapeutic purposes, mostly due to prominent

*Correspondence: Dr K Biber, Department of Medical Physiology, University Medical Center Groningen, University of Groningen, Ant. Deusinglaan 1, Groningen 9713 AV, The Netherlands, Tel: +31 50 3632706, Fax: +31 50 3632751, E-mail: k.p.h.biber@med.umcg.nl

⁶These authors contributed equally to this study.

Received 18 December 2006; revised 18 September 2007; accepted 29 September 2007

peripheral side effects, short half-lives, and/or receptor desensitization (Yan *et al*, 2003; Stone, 2002; Dalpiaz and Manfredini, 2002). Interestingly, it has been suggested that the enhancement of adenosine A₁ receptor expression might efficiently induce adenosine A₁ receptor-mediated neuroprotection (Rudolphi and Schubert, 1996).

Thus, we now investigated if IL-6 influences the expression and function of neuronal adenosine A₁ receptors *in vitro* and *in vivo*. Looking at three different pathological conditions in which adenosine has proven to be implicated as a protective agent—hypoxia, glutamate-induced cell death, and seizures—we evaluated if the beneficial properties attributed to IL-6 could be due, at least in part, to the amplification of adenosine A₁ receptor-mediated actions.

The results reported here underscore the crucial role played by IL-6 in the regulation of neuronal adenosine A₁ receptors and demonstrate that such mechanism is involved in the enhancement of A₁ receptor-mediated signaling in the brain under excitotoxic situations, with a beneficial impact on neuronal survival.

MATERIALS AND METHODS

Chemicals

Amphotericin B, pentylenetetrazole (PTZ), poly-L-lysine, RNase-free DNase and adenosine deaminase, (\pm)- α -amino-3-hydroxy-5-methylisoxazole-4-propionic acid hydrate (AMPA) were obtained from Sigma-Aldrich (Zwijndrecht, The Netherlands and Sintra, Portugal). Neurobasal medium, B27 supplement and gentamycin were from Gibco (Breda, The Netherlands). Recombinant IL-6 with a specific activity of $>5 \times 10^7$ U/mg was purchased from R&D systems (London, UK) or PeprTech EC Ltd (London, UK). In most experiments 10 U/ml IL-6 have been used which is comparable to <200 pg/ml. 1,3-Dipropyl-8-cyclopentylxanthine (DPCPX), N⁶-cyclopentyladenosine (CPA), 6-cyano-2,3-dihydroxy-7-nitro-quinoxaline (CNQX), DL-2-amino-5-phosphonovaleric acid (APV), bicuculline methochloride and cycloheximide (CHX), and tetrodotoxin citrate (TTX) were from Tocris Cookson Ltd (Bristol, UK). N⁶-cyclohexyladenosine ([³H]-CHA) was purchased from Moravex Biochemicals (Brea, CA, USA). iQ SYBR Green supermix at Bio-Rad (Veenendaal, The Netherlands). Rabbit polyclonal antibodies for the adenosine A₁ receptor were from Abcam (Cambridge, UK) or Affinity Bioreagents (Golden, CO, USA); monoclonal and polyclonal antibodies for β -actin were purchased from MP Biomedicals (Amsterdam, The Netherlands) and Abcam, respectively.

Animals

All experiments were performed with 3-month-old (except for neuronal cultures, see below) male mice (C57Bl6 and IL-6-deficient mice on C57Bl6 background; Jackson Laboratory, Bar Harbor, USA). The animals were handled according to European Community guidelines and in accordance with the regulations of the committee for use of experimental animals of the universities of Groningen and Lisbon. The animals were housed in groups of maximum five per cage in a room with controlled light/dark (12/12 h) cycle and temperature (21°C). Food and

water were supplied *ad libitum*. Mice were anesthetized with halothane before decapitation.

Electrophysiological Recordings in Hippocampal Slice Preparations

Slice preparation was carried out as previously described (Diógenes *et al*, 2004). Field excitatory postsynaptic potentials (fEPSPs) and whole-cell postsynaptic currents were recorded from the CA1 area of hippocampal slices (300–400 μ m) upon stimulation of the Schaffer collateral fibers (rectangular pulses of 0.1–0.2 ms duration delivered each 15 s). Drugs were applied to the perfusion solution (3 ml/min), which contained NaCl 125 mM, KCl 3 mM, NaH₂PO₄ 1.25 mM, NaHCO₃ 25 mM, CaCl₂ 2 mM, MgSO₄ 1 mM, and glucose 10 mM. Bath temperature was 32°C throughout fEPSPs recording experiments. To elicit paired-pulse facilitation (PPF; see eg Citri and Malenka, 2007), the Schaffer collaterals were stimulated twice with 50 ms interpulse interval. Synaptic facilitation was quantified as the ratio between the slopes of the fEPSP elicited by the second and the first stimuli. While evaluating the influence of a drug (CPA) on PPF, the difference in PPF values obtained in the presence and in the absence of that drug in the same slice (Δ PPF) was calculated. Hypoxia was induced substituting the artificial cerebrospinal fluid (aCSF) by an identical aCSF preequilibrated with 95% N₂/5% CO₂ for 3 min. This manipulation reduces bath oxygen tension in the recording chamber from ≈ 600 to ≈ 250 mm Hg (Sebastião *et al*, 2000). Each slice was subjected to a single period of hypoxia, since the effects of hypoxia may be modified by subsequent episodes in the same slice (Schurr *et al*, 1986). To minimize individual variation between slices, the effects of hypoxia were compared in control and in test slices with less than 1 h of interval.

Patch clamp experiments were performed at room temperature (22–25°C) in CA1 pyramidal cells identified under infrared visualization and electrophysiologically characterized (Staff *et al*, 2000). Patch electrodes (5–7 M Ω) were filled with 125 mM potassium gluconate, 11 mM KCl, 0.1 mM CaCl₂, 2 mM MgCl₂, 1 mM EGTA, 10 mM HEPES, 2 mM NaATP, 0.3 mM NaGTP, and 10 mM TRIS phosphocreatine; pH 7.3 adjusted with KOH, 280–290 mOsm. Every cell was recorded from a separate slice. Resting membrane potentials were between -60 and -70 mV. Junction potentials were not corrected. Whole-cell postsynaptic currents were recorded in voltage-clamp mode ($V_h = -70$ mV). Averages of four consecutive excitatory postsynaptic potentials (EPSCs) or inhibitory postsynaptic currents (IPSCs) were continuously acquired. IPSCs were obtained in the presence of CNQX (20 μ M) and APV (10 μ M) and were completely blocked after superfusion of bicuculline (20 μ M). AMPA receptor-mediated postsynaptic currents were evoked by pressure ejection of AMPA (20 μ M) to the dendrites in the presence of TTX (1 μ M). Data were digitally acquired in voltage-clamp mode using an EPC-7 amplifier (List Biologic, Campbell, CA), Digidata 1322A A-D converter and LTP program or Win LTP (Anderson and Collingridge, 2001, 2007). A small voltage step (5 mV, 50 ms) was delivered before evoking each postsynaptic current to monitor membrane and series resistances; if one

of both or holding current changed significantly, the experiment was rejected.

Preparation of Hippocampal Synaptosomes

Hippocampal slices kept in the absence or in the presence of exogenously added IL-6 for 6 h were collected into a sucrose solution (320 mM sucrose, 1 mM EDTA, 5 mM HEPES, and 1 mg/ml BSA, pH 7.4) and homogenized by four up-and-down strokes. The homogenate was then centrifuged at 3000g for 10 min (at 4°C), the supernatant was collected and centrifuged again at 14 000g for 12 min (at 4°C) to prepare the synaptosomal P2 fraction, as described previously (Pinto-Duarte *et al*, 2005).

Primary Neuronal Cultures

Cortical neurons were prepared from embryonic mouse brains (E15–16) as described before (de Jong *et al*, 2005). Neurons were used after 6 days in culture. Cells were incubated for 6 or 24 h with 10 U/ml IL-6 and then either stimulated for 1 h with glutamate at various concentrations or incubated for 15 min with CPA (1–100 nM) or 100 nM DPCPX followed by glutamate challenge for 1 h. The effect of test drugs (IL-6, CPA, or DPCPX) on glutamate-induced neurotoxicity was always evaluated comparing the effect of glutamate in the absence and presence of the test drug in the same batch of neurons.

Determination of Neuronal Viability

Neuronal viability was determined by the colorimetric MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay as described previously (Marchetti *et al*, 2004).

Seizure Induction by Pentylentetrazole Treatment

Male wild-type (WT) C57Bl6 and IL-6 knockout mice weighing 23–29 g were used. Mice received once daily, for 2 consecutive days, one intraperitoneal injection with 55 mg/kg PTZ dissolved in 0.9% saline. Control animals received saline only. Latency of seizure onset was timed and animals were subsequently observed for 1 h. Animals were assigned a seizure score according to the following criteria (Kondziella *et al*, 2002): 0 = normal behavior, 1 = myoclonic jerks, 2 = minimal seizures without Straub-tail, 3 = minimal seizures with Straub-tail, 4 = generalized tonic-clonic seizures, 5 = like 4 with loss of consciousness and postictal phase, 6 = like 5 with rotation on their axis, and 7 = like 6 and death. All PTZ-treated animals showed at least score 2 (minimal seizures) within 5 min. Mice were killed 26 h after the second injection.

Brain Tissue Preparation from PTZ-Injected Mice

Brains were removed, the two hemispheres were separated, and subsequently frozen and stored at –80°C until use. Both groups that received PTZ injections (WT and IL-6 knockout) were further divided according to the severity of the seizures. The ‘PTZ-high’ groups consisted of animals that showed tonic-clonic seizures (score 4) after at least one of the two PTZ injections. All other animals that received PTZ injections were assigned to the ‘PTZ-low’ groups. WT and IL-6 knockout control groups contained 15 animals, while the four PTZ-treated groups (PTZ-high and PTZ-low) consisted of 6–8 animals per group.

The right hemispheres were prepared for RNA analysis (see below) and radioligand binding studies (Lohse *et al*, 1984), while the left hemispheres were used for autoradiography.

RNA Isolation and Reverse Transcription–Polymerase Chain Reaction

Tissue was lysed in guanidinium isothiocyanate/mercapto-ethanol buffer and total RNA was extracted with one phenol/chloroform step (Chomczynski and Sacchi, 1987) and transcribed into cDNA as described (Biber *et al*, 2001).

Real-Time Polymerase Chain Reaction

Adenosine A₁ receptor mRNA expression in the brains of WT and IL-6 knockout mice was analyzed by real-time PCR using the iCycler (Bio-Rad) and the iQ SYBR Green supermix. Mouse ribosomal protein L32–3A (rpL32A) primers and hypoxanthine guanine phosphoribosyl transferase (HPRT1) primers were used for normalization to housekeeping genes. These genes did not show variations in response to the experimental treatment (see Table 1 for primer sequences). The comparative C_t method (amount of target amplicon X in sample S, normalized to a reference R and related to a control sample C, calculated by $2^{-((C_t X, S - C_t R, S) - (C_t X, C - C_t R, C))}$) was used to determine the relative expression levels (Livak and Schmittgen, 2001).

Western Blotting

Primary cortical neurons (2×10^6 cells per well) were washed twice with ice-cold phosphate-buffered saline, subsequently lysed by the addition of 0.15 ml of lysis buffer (20 mM Tris, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1% Triton, 2.5 mM sodium pyrophosphate, and 1 mM sodium orthovanadate and protease inhibitors). The homogenates were centrifuged for 10 min at 12 000 r.p.m. and the supernatants were collected. P2 synaptosomal fraction was

Table 1 Sequence of the Primers Used for Real-Time PCR

Gene	Accession number	Forward primer (5'-3')	Backward primer (5'-3')
rpL32A	X06483	GCTGGAGGTGCTGCTGATGT	ACTCTGATGGCCAGCTGTGC
HPRT1	X62085	GACTTGCTCGAGATGTCA	TGTAATCCAGCAGGTCAG
Ado A ₁ R	AJ555877	CCTCTCCGGTACAAGACAGT	GGTGTCCAGGCTACCACAAG

lysed in RIPA buffer (50 mM Tris, 150 mM NaCl, 1 mM EDTA, 1% NP-40, 0.1% SDS, 1 mM sodium orthovanadate and protease and phosphatase inhibitors, adjusted to pH = 8). Samples were boiled for 5 min in Laemmli's buffer (2% SDS, 5% DTT). A total of 20 µg protein in each sample was separated by 10–12% SDS-PAGE and transferred to Immobilon P polyvinylidene difluoride membranes (Millipore Corporation). After blocking for 1 h in TBS with 0.1% Tween 20 with 5% low-fat milk, membranes were incubated overnight at 4°C with a rabbit anti-adenosine A₁R antibody (1:1000, Affinity Bioreagents or 1:1000, Abcam) and then with a secondary antibody conjugated with horseradish peroxidase or with an alkaline phosphatase labeled secondary antibody for 1 h at room temperature. Afterwards it was processed for protein detection using Super Signal™ substrate (Pierce) or using the ECL detection system (Tropix) according to the manufacturer's instructions. β-Actin (1:10 000, Abcam or 1:10 000, MP Biomedicals) was used as a loading control. We recently cloned and expressed mouse adenosine A₁ receptor in CHO cells (Wittendorp *et al*, 2004). Accordingly, CHO cells expressing mouse adenosine A₁ receptor have been used as a positive control in western blotting experiments showing that the used antibody recognized a band of 37 kDa only in cells that stably expressed mouse adenosine A₁ receptors (data not shown).

Radioligand Binding Assay

For the determination of the maximum bound radioligand, membranes were incubated for 1 h at 25°C in 50 mM Tris/HCl (pH 7.4 at 25°C) in the presence of 4 nM [³H]-DPCPX (K_i = 1.4 nM); at this concentration one can assume that all adenosine A₁ receptors present in the preparation will be occupied, making the amount of [³H]-DPCPX binding informative on the number of adenosine A₁ receptors. Each experiment was performed in triplicate. Nonspecific binding was determined in the presence of 100 µM CPA. Incubations were stopped with 1 ml ice-cold buffer and bound radioligand was subsequently recovered by filtration through Whatman GF/B filters. Filters were washed three times and retained radioactivity was measured by liquid scintillation counting (LKB Wallac, 1219 Rackbeta).

Autoradiography

Sections were preincubated in Tris-HCl buffer, 170 mM, pH 7.4 containing 0.5 U/ml adenosine deaminase for 3 × 10 min at room temperature, washed out, and dried in a stream of cold air. For the labeling of adenosine A₁ receptors, sections were incubated with 3.0 nM [³H]N6-cyclohexyladenosine ([³H]-CHA, specific activity 20 Ci/mmol) in 170 mM Tris-HCl buffer including 0.5 IU/ml adenosine deaminase at room temperature for 90 min. To determine nonspecific binding, sections were incubated in the presence of 3 µM CPA. Following incubation, unbound radioactivity was removed by rinsing the sections three times for 2 min in ice-cold Tris-HCl buffer, 50 mM (pH 7.4), and rinsing once for 5 s in ice-cold distilled water to remove buffer salts. The sections were then dried in a stream of cold air. Dried sections were exposed to Numico hyperfilm, stored in an X-ray film cassette for 4 month at -80°C and subsequently

developed. Quantification analysis of the resulting autoradiographic images was performed using an automatic image analysis system (Quantimet 500, Leica, Cambridge). Optical density was measured in hippocampus and cortex in six sections per mouse. The values of the sections were averaged for each mouse. Optical density measurements from each brain area were converted into nCi per mg tissue, according to the calibration curve obtained from the tritium standards. Results are given as mean ± SEM (*n* = 6).

Data Analysis

Data are presented as mean ± SEM from *n* experiments. When comparing two groups of results, statistical significance was assessed using Student's *t*-test. When doing multiple comparisons, statistical significance was assessed by one-way ANOVA followed by the Bonferroni correction. Values of *P* < 0.05 were considered statistically significant.

RESULTS

Influence of IL-6 on Adenosine A₁ Receptor-Mediated Actions on Synaptic Transmission

IL-6 enhances the expression and the inhibitory actions of presynaptic adenosine A₁ receptors on glutamatergic transmission. Adenosine A₁ receptor activation is well known to decrease synaptic transmission in the hippocampus (Sebastião *et al*, 1990). Therefore, electrophysiological experiments were carried out to investigate if a sustained elevation of IL-6 levels in the hippocampus modified the neuromodulatory actions of adenosine A₁ receptors. In the first series of experiments we performed cumulative concentration–response curves for the inhibitory effects of the A₁ selective agonist, CPA (1.5–50 nM), on the slope of fEPSPs recorded from control slices and slices exposed to IL-6 (10 U/ml) for 5–8 h. As Figure 1a shows, treatment with IL-6 induced a shift to the left of the concentration–response curves of CPA, without changing its maximal effect. The EC₅₀ of CPA-induced inhibition was 5 nM (95% confidence interval 4.8–5.2 nM, *n* = 3) in the presence of IL-6, significantly different (*p* < 0.05) from the control EC₅₀, which was 7.8 nM (95% confidence interval 7.7–7.9 nM, *n* = 5) (Figure 1a and b).

Also when CPA was tested at a single concentration of 5 nM, the inhibition of fEPSPs was significantly more pronounced (*p* < 0.05) (32.2 ± 3.6%, *n* = 5) in slices exposed to IL-6 (10 U/ml for 5–8 h), as compared to slices kept in the absence of other pharmacological drugs (18.2 ± 1.8%, *n* = 6). To rule out nonspecific effects of IL-6, in a different pool of experiments we compared the effect of 5 nM CPA in control slices and in slices treated with heat-inactivated IL-6 for 5–8 h (HI-IL-6 consisted of 10 U/ml IL-6 boiled for 30 min). As expected, the inhibition caused by CPA was not significantly different (*p* > 0.05) in both conditions (control: 23.5 ± 2.5%; HI-IL-6: 23.8 ± 1.4%; *n* = 2).

Previous studies have shown that adenosine A₁ receptor activation enhances PPF, presumably by inhibiting the release of neurotransmitters induced by the first stimulus and consequently increasing the amount of neurotransmitters to be released by the second stimulus. Thus, we performed PPF experiments to investigate if IL-6-induced




Figure 1 Interleukin-6 (IL-6) potentiates the neuromodulatory actions of presynaptic adenosine A₁ receptors on hippocampal synaptic transmission. (a) Cumulative concentration–response curves for the effects of the A₁ receptor agonist, N⁶-cyclopentyladenosine (CPA, 1.5–50 nM), on the slopes of field excitatory postsynaptic potentials (fEPSPs) recorded from the CA1 area of hippocampal slices. In the ordinates, 0% corresponds to fEPSPs slopes before the addition of CPA (1.30 ± 0.14 mV/ms, *n* = 5, in control slices, ●, and 1.36 ± 0.17 mV/ms, *n* = 3, in slices exposed to IL-6 for 5–8 h, ○), and 100% represents a complete inhibition of fEPSPs. (b) Superimposed traces obtained before (pre-CPA) and after the application of CPA in a control slice (left) and in a slice previously exposed to IL-6 for 5 h (right). Each trace is the average of eight consecutive responses and is composed of the stimulus artifact, presynaptic volley, and fEPSP; the scale applies to panels (a) + (b). (c) Variation (Δ) of paired-pulse facilitation (PPF) in the absence and presence of CPA in three matched experiments performed in control slices (●) and in slices from the same hippocampi exposed to IL-6 for 5–8 h (○). Individual ΔPPF values in the three experiments were 0.07 (control) and 0.17 (IL-6); 0.09 (control) and 0.19 (IL-6); and 0.08 (control) and 0.20 (IL-6).

potentiation of A₁ receptor neuromodulatory actions occurred at presynaptic sites. In the absence of CPA, PPF corresponded to a ratio of 1.6 ± 0.08 in control conditions and 1.7 ± 0.1 in IL-6-treated slices (*n* = 3 each). As expected,

CPA (5 nM) increased PPF in both conditions; remarkably, as shown in Figure 1c, CPA-induced enhancement of PPF (ΔPPF, calculated as PPF_{CPA}–PPF_{pre-CPA} in the same slice) was always significantly greater (*p* < 0.05, *n* = 3 matched experiments) in IL-6-treated slices (0.19 ± 0.01) than in control slices (0.08 ± 0.01), supporting a presynaptic locus for IL-6-induced enhancement of adenosine A₁ receptor-mediated effects on synaptic transmission.

To investigate if IL-6-induced enhancement of adenosine A₁ receptor-mediated inhibition of synaptic transmission occurred at the single-neuron level, we performed whole-cell voltage-clamp recordings from CA1 pyramidal cells upon stimulation of the Schaffer collateral afferents. To avoid preconditioned responses of A₁ receptors, the effect of CPA (5 nM) was investigated in only one neuron per slice, control (absence of exogenous IL-6) and test (incubation with 10 U/ml IL-6 during 5–8 h) slices being from the same hippocampus. As illustrated in Figure 2 (panels A and B), CPA (5 nM) inhibited EPSCs by 27.5 ± 1.11% (*n* = 5) in control conditions. IL-6 significantly (*p* < 0.05) exacerbated CPA-induced depression of the EPSCs to 52.7 ± 5.36% (*n* = 3; Figure 2a and b). Similar effects were found with 100 U/ml IL-6 (Figure 2b). The incubation with or without IL-6 for up to 8 h did not change tissue integrity, the morphology of the cells nor the effects of CPA on EPSPs (data not shown). We also looked for a putative role of postsynaptic A₁ receptors on the modulation of synaptic transmission. Since in our conditions (high Mg²⁺ and V_h = –70 mV) synaptic transmission was mostly mediated by the activation of postsynaptic AMPA receptors, we tested the action of CPA on pharmacologically isolated postsynaptic AMPA receptor-mediated currents. The amplitude of postsynaptic AMPA-induced currents was not affected by CPA (5 nM) in control conditions (106.0 ± 7.4% of control, *n* = 3). When the concentration of CPA was increased to 100 nM, AMPA-evoked currents were still not significantly modified (*p* > 0.05), which solely confirmed the limited contribution of postsynaptic adenosine A₁ receptors to the modulation of synaptic transmission (Arrigoni *et al*, 2005); it did not preclude, however, that the upregulation of postsynaptic or somatic A₁ receptors might assume functional relevance in other experimental paradigms, a possibility which we explored and were able to demonstrate further below using a different *in vitro* model (see data on neuronal cultures).

The next group of experiments aimed to examine the participation of *de novo* protein synthesis in IL-6-induced enhancement of A₁ receptor function on hippocampal synaptic transmission. As shown in Figure 2b, treatment of hippocampal slices with the protein synthesis inhibitor CHX during the incubation period with IL-6, completely abolished the potentiation of the inhibitory effect of CPA (5 nM) on electrically evoked EPSCs. CHX *per se* did not significantly modify 5 nM CPA-induced inhibition of EPSCs (31.7 ± 1.6%, *n* = 3, *p* > 0.05 as compared to a control inhibition of 27.5 ± 1.1%, *n* = 5). To investigate whether the involvement of newly synthesized proteins was correlated with the upregulation of presynaptic adenosine A₁ receptors, we performed western blot experiments in synaptic terminals prepared from control hippocampal slices and IL-6-treated slices. As shown in Figure 2c, IL-6 induced a significant (*p* < 0.05) upregulation of A₁ receptors




Figure 2 Interleukin-6 (IL-6)-induced enhancement of adenosine A₁ receptor function on synaptic transmission requires *de novo* protein synthesis and is matched by the expression of newly synthesized A₁ receptors. (a) Representative excitatory postsynaptic potentials (EPSCs) recorded from pyramidal cells before and during superfusion of N⁶-cyclopentyladenosine (CPA, 5 nM) in control (CTR) and IL-6 (10 U/ml)-treated hippocampal slices (6 h), in the absence or in the presence of the protein synthesis cycloheximide (CHX), as indicated; superimposed recordings were from the same slice. Stimulus artifacts are truncated. (b) Histogram summary percentage decrease in EPSC amplitudes caused by CPA; the slices had been incubated in the absence (control, $n = 5$) or in the presence of the drugs indicated below each bar ($n = 3$ in all cases except for 100 U/ml IL-6 where $n = 2$); time of incubation with IL-6 and/or CHX: 6–8 h. * $p < 0.05$ vs control, ** $p < 0.05$ vs IL-6 (10 U/ml), one-way ANOVA followed by the Bonferroni correction. Zero percent in the ordinates represents the EPSCs recorded before applying CPA and were 251 ± 40 pA in the absence of any drug, 243 ± 46 pA in slices incubated with IL-6, 241 ± 59 pA in slices incubated with CHX, and 240 ± 35 pA in slices incubated with CHX plus IL-6; 100% represents a complete inhibition of currents. (c) Western blot analysis of adenosine A₁ receptor expression in nerve terminals prepared from slices hippocampal slices kept for 6–8 h in the absence or in the presence of IL-6 (10–100 U/ml). Pooled data from four individual experiments and a demonstrative experiment are shown.

in this preparation. Taken together, these data suggest that newly synthesized presynaptic A₁ receptors are involved in the enhancement of A₁ receptor-mediated effects on synaptic transmission.

While recording EPSCs, GABAergic transmission was not blocked to avoid exacerbated excitability of the slices. The next series of experiments was therefore performed to exclude putative actions of IL-6-induced upregulation of A₁ receptors on GABAergic transmission. In these trials compound synaptic currents were initially recorded and, after stabilization, the antagonists of AMPA/kainate receptors, CNQX (20 μM), and NMDA receptors, APV (10 μM), were superfused to isolate the IPSCs. The GABAergic nature of these currents was confirmed at the end of each experiment using the GABA_A receptor antagonist bicuculline (20 μM), which totally blocked the remaining current. In both control and IL-6-stimulated slices, the GABAergic current represented nearly 10% of the compound current, as assessed by the degree of inhibition by ionotropic glutamate receptor antagonists. Activation of adenosine A₁ receptors with CPA (5 nM) did not affect IPSCs ($99.7 \pm 5.9\%$ of currents amplitude before CPA, $n = 3$) recorded from control slices, which is in accordance with previous data showing the inability of A₁ receptors to modulate GABAergic transmission in mature hippocampus (Lambert and Teyler, 1991; Yoon and Rothman, 1991). In slices incubated with IL-6 (10 U/ml) for >5–8 h, CPA (5 nM) also did not induce significant modifications on IPSCs amplitudes ($95.6 \pm 9.9\%$ of currents amplitude before CPA, $n = 3$, $p > 0.05$), indicating that IL-6 was unable to unmask a putative A₁ receptor-mediated action upon GABAergic transmission in the hippocampus.

IL-6 enhances adenosine A₁ receptor-mediated depression of synaptic transmission during hypoxia. We next investigated the pathophysiological relevance of IL-6-induced enhancement of adenosine A₁ receptor function on synaptic transmission. The inhibition of synaptic transmission in the hippocampus is crucial to protect neurons from excitotoxicity induced by neuropathological events, such as hypoxia. Under such conditions, considerable amounts of adenosine are released and operate A₁ receptors to restrain synaptic glutamate release (Sebastião *et al*, 2001). Thus, we tested the effect of brief (3 min) hypoxic episodes in fEPSPs recorded from control and IL-6-treated slices. This manipulation caused little or no appreciable depression of synaptic transmission under control conditions (Figure 3a). However, the same hypoxic insult applied to hippocampal slices previously exposed to IL-6 (10 U/ml) for 6–8 h, induced a significant ($p < 0.05$) depression of fEPSPs (Figure 3a). This depression was markedly attenuated when DPCPX (100 nM) was present during the incubation period with IL-6 and throughout the experiment ($n = 2$). Yet, the presence of DPCPX during such prolonged period significantly boosted neuronal excitability, increased population spike contamination, and caused abnormal modifications of the recorded fEPSPs, which led us to perform a different pool of experiments in which DPCPX (100 nM) was applied about 30 min before the hypoxic insult. In these conditions, the depression of fEPSPs recorded from IL-6-treated slices was fully prevented (Figure 3b) and, therefore, can be attributed to the activation of A₁ receptors by adenosine. In one




Figure 3 Interleukin-6 (IL-6) enhances A₁ receptor-mediated inhibition of synaptic transmission induced by hypoxia. (a) Time course of the effects of the effect of 3 min hypoxia (N₂, applied as indicated by the horizontal bar) on field excitatory postsynaptic potentials (fEPSP) slope recorded from control slices (absence of IL-6, n = 3, ●) or from slices taken from the same hippocampi but that had been preincubated with IL-6 (10 U/ml) for 6–8 h (n = 3, ○). The inset shows sample traces of fEPSPs recorded before hypoxia (1 and 3) and after 3 min hypoxia (2 and 4) in the absence (left, 1 and 2) or in the presence of IL-6 (right, 3 and 4); superimposed recordings in each panel were from the same slice, the left and right panel recordings were from different slices from the same hippocampus (see Figure 1 legend for further details). (b) Effects of a 3 min hypoxic insult induced in the presence of a selective A₁ receptor antagonist, 1,3-Dipropyl-8-cyclopentylxanthine (DPCPX), in control conditions (n = 3, ○) and in slices preincubated with IL-6 for 6–8 h (n = 4, ●). (c) Time course of the response to a longer period of hypoxia (14 min) in the presence of DPCPX, to block A₁ receptor actions. In all panels; 100% in the ordinates corresponds to the averaged fEPSP slope recorded for 10 min before N₂; the slope values within the same panel do not differ significantly ($p > 0.05$) between test (slices incubated with IL-6) and control conditions.

receptor antagonist, DPCPX (200 nM), was not modified ($p > 0.05$) by preincubation with IL-6.

Influence of IL-6 on Adenosine A₁ Receptor Actions Upon Excitotoxicity

IL-6 increases the function and neuroprotective properties of postsynaptic adenosine A₁ receptors against glutamate-induced neurotoxicity. It is well known that the excessive activation of postsynaptic glutamate receptors by high concentrations of glutamate induces neuronal death in cultured cells. Interestingly, IL-6 was reported to minimize the consequences of high-glutamate levels by promoting neuronal rescue (Ali *et al*, 2000). To investigate a possible postsynaptic locus of IL-6-induced upregulation of A₁ receptors, we tested whether those neuroprotective actions of IL-6 could be related to the amplification of A₁ receptors function. To minimize the number of killed animals, we cultured cortical neurons, which also abundantly express A₁ receptors. The rapid induction of neuronal adenosine A₁ receptor mRNA and protein by IL-6 (1–24 h) was also observed in these neurons through reverse transcription (RT)-PCR experiments (Figure 4a) and quantitative analysis of four independent western blot experiments, respectively (Figure 4a).

In our conditions, glutamate-dependent neurotoxicity was completely abolished by MK801 treatment and therefore due to NMDA receptor stimulation (data not shown). We first compared the effect of various concentrations of glutamate (10 μ M up to 300 μ M) on the survival of cultured neurons kept in the absence or in the presence of IL-6. We found that treatment with IL-6 (10 U/ml) significantly increased neuronal rescue in comparison to control at all concentrations confirming earlier findings (Ali *et al*, 2000) whereas heat-inactivated IL-6 did not have any effect on neuronal survival (data not shown). In subsequent experiments adenosine A₁ receptor activity was blocked by a 15 min preincubation with DPCPX (100 nM). The blockade of adenosine A₁ receptor activity had no influence on glutamate-induced neuronal death but completely abolished the protective effect of IL-6 treatment as exemplified for 50, 25, and 10 μ M glutamate (Figure 4b). On the other hand a

experiment, we induced a longer hypoxic period (4 min), sufficient to cause a mild depression of synaptic transmission even in control conditions (33.3%) and the response to hypoxia was still more pronounced (67.5%) in a slice taken from the same hippocampus previously exposed to IL-6 (10 U/ml) for 6 h.

Since prolonged hypoxic periods cause an inhibition of synaptic transmission that results from both adenosine A₁ receptor-dependent and -independent mechanisms (Lucchi *et al*, 1996), we further investigated if the later were also influenced by IL-6. As shown in Figure 3c, the depression of synaptic transmission caused by 14 min of hypoxia in the presence of a supramaximal concentration of the A₁




Figure 4 Effects of interleukin-6 (IL-6) treatment on adenosine A₁ receptor expression and on excitotoxicity in primary cortical cultures. (a) Upregulation of adenosine A₁ receptor mRNA and protein expression by IL-6 treatment (10 U/ml) in cortical neurons. The inset in the figure shows a typical western blot experiment. (b) Pretreatment with IL-6 (10 U/ml 24 h) protected neurons from excitotoxicity induced by glutamate at various concentrations (50 μM, 25 μM, and 10 μM). Treatment with 100 nM DPCPX (15 min before glutamate treatment) completely abolished the protective effect of IL-6 treatment but did not influence the effect of glutamate without IL-6 pretreatment. (c) Application of N⁵-cyclopentyladenosine (CPA, 15 min before glutamate treatment) at several concentrations (1, 10, and 100 nM) significantly increased neuronal survival after glutamate treatment (50 μM) only when neurons had been pretreated with IL-6. CPA (10 nM) without IL-6 did not affect glutamate-induced neurotoxicity. Data in (a) are given as mean ± SEM from four independent western blot experiments and from three independent RT-PCR experiments. Data in (b) and (c) are given as mean ± SEM ($n = 3$). The graphs represent typical experiments. Similar results have been obtained in five independent experiments. *Significantly different ($p < 0.05$).

pretreatment with 1, 10, and 100 nM CPA, respectively). Data shown in Figure 4a–c concern a pool of experiments in which treatment with IL-6 was performed during 24 h, but similar results were obtained when treatment with IL-6 was performed during 6 h (data not shown).

Influence of IL-6 on Adenosine A₁ Receptors *In Vivo*

The next series of experiments were aimed to investigate the role of IL-6 on the regulation of adenosine A₁ receptor expression in the brain *in vivo*. It is known that the expression of A₁ receptors in WT mice is increased after neuropathological events, such as seizures (Pagonopoulou *et al*, 1993; Vanore *et al*, 2001; Angelatou *et al*, 1991, 1993). On the other hand, several reports have shown that those noxious conditions trigger the release of IL-6 (Lehtimäki *et al*, 2003; Peltola *et al*, 1998, 2002). Accordingly, we used WT animals and IL-6-deficient mice to evaluate the relationship between the expression of IL-6 and the upregulation of adenosine A₁ receptors in response to PTZ-induced seizures.

Differences in seizure scores between WT animals and IL-6-deficient mice. As shown in Figure 5, seizure scores after a first PTZ injection did not differ between WT and IL-6-deficient mice. However, whereas no change in seizure scores were observed in WT mice after a second PTZ injection round, IL-6-deficient mice displayed significant higher seizures ($p = 0.03$) on the second day (Figure 5). Moreover, none of the WT mice died after PTZ injections, whereas two of the IL-6-deficient mice did not survive the second application of PTZ.

further stimulation (15 min before glutamate treatment) of adenosine A₁ receptors by various concentrations (1, 10, and 100 nM) of CPA significantly increased the neuronal survival after glutamate treatment (50 μM) (Figure 4c). As shown in Figure 4c CPA treatment (exemplified by 10 nM) without IL-6 preincubation did not significantly protect neurons against 50 μM glutamate ($42 \pm 7\%$ surviving neurons for glutamate *per se* and 46 ± 7 , 44 ± 8 , and $49 \pm 4\%$ surviving neurons were found after 15 min

Adenosine A₁ receptor mRNA expression is downregulated after seizures in IL-6-deficient mice. Q-PCR analysis was carried out to investigate possible effects of seizure activity on A₁ receptor mRNA expression in IL-6-deficient mice and WT animals. No differences were observed in mice with lower seizure scores compared to control mice (data not shown). Similarly, no change in A₁ receptor expression was measured in WT animals with high seizure scores (Figure 6a). In contrast a significant ($p = 0.03$)




Figure 5 Differences in seizure scores between wild-type (WT) and IL-6-deficient mice. There was no difference in seizure scores between WT and IL-6-deficient mice after the first PTZ injection, both mouse strains displayed mean seizure scores around 3. Similar seizure scores were also detected in WT mice after the second PTZ injection round, whereas IL-6-deficient mice displayed significant higher seizures on the second day. *Significantly different ($p < 0.05$).

downregulation of adenosine A₁ receptor mRNA expression was detected in IL-6-deficient mice with high seizure scores compared to controls (Figure 6a).

Seizure-induced upregulation of brain adenosine A₁ receptor expression depends on IL-6. Radioligand binding experiments using [³H]-DPCPX have been performed to investigate the expression of adenosine A₁ receptors in the brain. The mice that received PTZ injections were divided in two groups according to the severity of the seizures in 'PTZ-low score' and 'PTZ-high score' groups as described in 'Materials and Methods'. No difference was obtained between control and PTZ-low score WT animals. An increase in [³H]-DPCPX binding was observed in WT of the PTZ-high score group, but this difference just failed to reach significance (p -value = 0.063). PTZ injections did not influence [³H]-DPCPX binding in IL-6-deficient mice (Figure 6b). The failure to reach significance in WT animals most likely is due to the fact that a prominent seizure-dependent induction of adenosine A₁ receptor does not occur in all brain regions (Pagonopoulou *et al*, 1993; Vanore *et al*, 2001; Angelatou *et al*, 1993). It was therefore further decided to investigate the seizure effects in autoradiographic experiments that allowed the precise analysis of adenosine A₁ receptor expression in defined brain regions.

Adenosine A₁ receptor expression in animals with high seizure scores was further determined in hippocampus (CA1, CA2, CA3, and dentate gyrus) and cortex (entorhinal cortex), brain areas with high density of adenosine A₁ receptors and in which a prominent induction of receptor expression was reported after PTZ-induced seizures (Pagonopoulou *et al*, 1993; Vanore *et al*, 2001; Angelatou *et al*, 1993). A significant seizure-dependent induction of adenosine A₁ receptor expression was observed in all these brain areas that resembled earlier findings (Pagonopoulou *et al*, 1993) (Figure 7a exemplified for CA1 and cortex), whereas no




Figure 6 Effects of seizure-induction on adenosine A₁ receptor mRNA and protein expression in brain. (a) Real-time PCR analysis of adenosine A₁ receptor mRNA level in brains of wild-type (WT) and interleukin-6 (IL-6) knockout (KO) mice (normalized to rpl32A and HPRT1). Graph shows the relative expression level of mRNA in pentylenetetrazole (PTZ)-injected animals ('PTZ-high score') compared to controls. The expression level of controls is set to 1. *Significant reduction of mRNA expression compared to controls ($p < 0.05$), determined by *t*-test. (b) Specific radioligand [³H]-1,3-dipropyl-8-cyclopentylxanthine (DPCPX) binding to adenosine A₁ receptors in brain membranes of WT and IL-6 KO mice with and without PTZ injection. The observed induction of adenosine A₁ receptor expression in WT animals just failed to reach significance ($p = 0.063$ as determined by one-way ANOVA). Results are given as mean ± SEM ($n = 5$, for mRNA expression; $n = 3$ for radioligand binding).

effect was observed in IL-6-deficient animals. Pooling together the relative changes obtained in all brain areas seizures significantly ($p = 0.02$) increased the A₁ receptor expression in WT animals by $28 \pm 8\%$ as compared to the basal situation. IL-6-deficient mice displayed significant ($p = 0.02$) lower expression of adenosine A₁ receptors ($85 \pm 2\%$ without seizure) when compared to WT controls and seizures did not change the expression of adenosine A₁ receptors in these animals ($80 \pm 3\%$ with seizures).

To investigate whether the seizure score after the first PTZ injection correlated with the upregulation of adenosine A₁ receptors we plotted day 1 seizure scores of PTZ-injected animals against the [³H]CHA-binding results of the autoradiography experiments of individual animals (control and PTZ-injected animals). Statistical analysis revealed that the Pearson's correlation coefficient was significant in all hippocampal areas (exemplified for CA1, $p = 0.003$)

(Figure 7b) and cortex $p=0.0019$ (data not shown), indicating a positive correlation between seizure scores observed on day 1 and adenosine A₁ receptor expression in WT animals.

DISCUSSION

In the present work we identified IL-6 as an important factor for regulating the expression of functional hippocampal and cortical adenosine A₁ receptors, thereby providing enhanced protection against noxious conditions both *in vitro* and *in vivo*. In our conditions, the exposure to IL-6 significantly enhanced the ability of adenosine A₁ receptors to inhibit excitatory synaptic transmission in acute hippocampal slices, an effect that was particularly evident during hypoxic insults. Since the efficient inhibition of synaptic transmission by adenosine during hypoxia facilitates the subsequent recovery to basal conditions (Sebastião *et al*, 2001), it is likely that the amplification of A₁ receptor-mediated function by IL-6 minimizes the consequences of a hypoxic period in the brain. Our data furthermore show that IL-6-induced enhancement of A₁ receptor expression has a great relevance to the survival of endangered neurons exposed to high concentrations of glutamate. However, the mechanisms by which IL-6 potentiates A₁ receptor function on synaptic transmission an on neuronal survival might not necessarily be connected. While the prevention of glutamate-induced neuronal death probably occurs at the postsynaptic or somatic level, the enhanced inhibition of synaptic transmission is mostly a presynaptic phenomenon. *In vivo* experiments furthermore suggested a crucial role for IL-6 in regulating the expression of adenosine A₁ receptor in the brain. Using autoradiography we observed a significant increase of adenosine A₁ receptor expression in the hippocampus and cortex in WT animals after PTZ-induced seizures, as it was already described by others (Pagonopoulou *et al*, 1993; Vanore *et al*, 2001; Angelatou *et al*, 1991). A remarkable novelty in the present work was that the PTZ-induced increased of adenosine A₁ receptor expression was completely lacking in the brains of IL-6-deficient mice. It was previously suggested that the upregulation of adenosine A₁ receptor expression occurring in response to seizures would protect neurons against subsequent convulsive episodes (Vanore *et al*, 2001, Gouder *et al*, 2003). Our data corroborates that hypothesis, since animals that failed to increase the expression of adenosine A₁ receptors in the brain (IL-6-deficient mice) experienced higher seizure scores after the second injection of PTZ and, in some cases, even death. Interestingly, the lack of adenosine A₁ receptor upregulation was observed in cortex and hippocampus of IL-6-deficient mice, indicating that with respect to the relationship between IL-6 and adenosine A₁ receptor expression, both brain regions behaved similar. This *in vivo* observation is corroborated by our *in vitro* findings, since IL-6 also increased neuronal adenosine A₁ receptor expression and function in hippocampal slices and in cortical neurons.

We previously showed that IL-6 induces the expression of adenosine A₁ receptors in glia cells (Biber *et al*, 2001), where even modest changes in A₁ receptor expression cause marked changes in adenosine A₁ receptor-mediated




Figure 7 Autoradiographic analysis of adenosine A₁ receptor expression in hippocampus and cortex of wild-type (WT) and interleukin-6 (IL-6)-deficient mice. Adenosine A₁ receptor expression was determined in hippocampus (CA1) and cortex (entorhinal cortex). (a) Note that seizures significantly induced the A₁ receptor expression compared to the basal situation. Untreated IL-6-deficient mice (IL-6 KO) displayed lower expression of adenosine A₁ receptor when compared to WT, which was unaffected by the induction of seizures (IL-6 KO seizures). (b) Adenosine A₁ receptor expression in WT mice is correlated to seizure scores after the first PTZ injection. Correlation analysis of adenosine A₁ receptor expression and seizure scores were carried out by plotting the expression data against the seizure scores compared to untreated control animals. Each data point in the graphs represents an individual animal (seven control animals; 3, 1, and 4 animals with seizure scores 2, 3, and 4, respectively). Positive correlations were found between adenosine A₁ receptor expression in hippocampus (b) and seizure scores induced by the first PTZ injection in WT animals. Data in (a) are represented as mean \pm SEM per group (4–8 animals per group). *Significantly different from untreated controls; $p < 0.05$ by one-way ANOVA. Statistical analysis revealed a significant Pearson's correlation coefficient of $p = 0.003$ for hippocampus (b).

signaling (Biber *et al*, 1997). Now, we extended these findings to neurons, the brain cells where the predominant expression of adenosine A₁ receptors occurs (Onodera and Kogure, 1988; Ulas *et al*, 1993; Swanson *et al*, 1995) and where adenosine acts to refrain excitatory transmission and to protect from uncoordinated firing (Dragunow, 1988; Dunwiddie and Masino, 2001). Indeed, the neuronal adenosine A₁ receptor represents a major endogenous neuroprotective system and it has long been suggested that the upregulation of A₁ receptor expression increases its

neuroprotective capacities (for review, see Rudolphi and Schubert, 1996; Dunwiddie and Masino, 2001). Accordingly, upregulation of adenosine A₁ receptor expression has been observed in mice and rats after chemically induced seizures and in post-mortem tissue of humans suffering from epilepsy (Pagonopoulou *et al*, 1993; Vanore *et al*, 2001; Angelatou *et al*, 1991). An upregulation of adenosine A₁ receptors in the presence of extracellular adenosine at high concentrations (as it occurs during seizures) seems a rather unusual feature, since adenosine A₁ receptors are normally desensitized in the presence of high concentrations of adenosine or after a prolonged exposure to selective agonists (Abbracchio *et al*, 1992; Ruiz *et al*, 1996; Hettinger *et al*, 1998; Latini and Pedata, 2001). The data presented here might provide an explanation for this paradoxical situation, since IL-6-induced synthesis of A₁ receptors could counteract or overrule A₁ receptor desensitization caused by prolonged exposure to high levels of adenosine released during seizures.

Our data showing that IL-6-deficient mice have increased susceptibility to PTZ-induced seizures are in accordance with two recent studies (de Luca *et al*, 2004; de Sarro *et al*, 2004), which showed that seizures in IL-6-deficient mice could be induced by much lower doses of several substances. By analyzing the effects of these different substances, the authors (de Sarro *et al*, 2004) concluded that IL-6 knockout mice might have impaired GABA inhibitory inputs and enhanced neurotransmission through glutamate receptors. In light of the present results, this enhancement of excitatory transmission is probably due to a reduced expression of adenosine A₁ receptors in the brain of IL-6-deficient mice. IL-6-dependent increase in neuronal A₁ receptor function might also underlie IL-6-induced inhibition of glutamate release and neuronal excitability in rat cortex (D'Arcangelo *et al*, 2000).

Inhibition of the expression of inflammatory genes in brain significantly aggravates the severity of PTZ-induced seizures (Asanuma *et al*, 1995). It is noteworthy that such inhibition only affected seizures after the second PTZ injection, similarly to what we observed in IL-6-deficient mice. Asanuma *et al* (1995) concluded that neuroinflammation occurring in response to the first seizures would initiate processes that protect the brain from upcoming seizure events. We now advanced a step in this hypothesis, showing that inflammatory events (release of IL-6) upregulate the expression and function of protective adenosine A₁ receptors.

A reduced inflammatory reaction in conjunction with an increase in mortality of IL-6-deficient mice after seizure induction was described by Penkowa *et al* (2001). Two of our IL-6-deficient mice (but none of the controls) also died as consequence of the second PTZ induction. Therefore, although IL-6 might have modest neuroprotective effects *in vitro* (present study, Ali *et al*, 2000; Pizzi *et al*, 2004) its function *in vivo* can be crucial for animal survival. Since an increased expression of IL-6 (see below) and of adenosine A₁ receptors also occurs in humans (Angelatou *et al*, 1993), it is likely that rapid IL-6 release in response to seizures is protective in upcoming seizure events by upregulating neuronal adenosine A₁ receptors. This might be different when IL-6 is chronically elevated. Increased neuronal death has been described after chronic treatment with IL-6 *in vitro*




Figure 8 Scheme that illustrates the proposed relationship between interleukin-6 (IL-6) and the adenosinergic system in neuroprotection. (I–III) Classical pathway in adenosine A₁ receptor-mediated neuroprotection. Any neuropathological condition induces the release of adenosine (I) that stimulates neuronal adenosine A₁ receptors (II) and causes neuroprotection by membrane hyperpolarization and inhibition of neurotransmitter release (III). (IV–VI) Adenosine at high concentration is sufficient enough to stimulate low-affinity adenosine A_{2B} receptors in astrocytes (IV) that induce the synthesis of IL-6 in these cells (V). IL-6 upregulates the expression of neuronal adenosine A₁ receptors (VI) and thus guarantees the presence of protective receptors that without IL-6 would undergo desensitization due to high ligand concentrations.

and animals with chronic IL-6 expression in astrocytes show enhanced neurodegeneration and increased sensitivity to kainite-induced seizures (Qiu *et al*, 1998; Nelson *et al*, 2004; Conroy *et al*, 2004; Campbell *et al*, 1993; Samland *et al*, 2003). It is not yet clear whether adenosine A₁ receptor expression is affected by chronic elevated IL-6 levels. Experiments to address this question are currently underway in our laboratory.

Rapid induction of IL-6 expression in the brain is known to occur in most (if not all) neuropathological conditions, including seizures in humans (Lehtimäki *et al*, 2003, 2004; Peltola *et al*, 1998, 2002; Jankowsky and Patterson, 1999) where IL-6 levels in the cerebrospinal fluid up to 100 pg/ml have been found (Peltola *et al*, 2000). Astrocytes are a major source of IL-6 in the brain (van Wagoner and Benveniste, 1999). We and others have shown that activation of adenosine A_{2B} receptors in astrocytes leads to the synthesis and release of IL-6 (Fiebich *et al*, 1996; Schwaninger *et al*, 1997, 2000). Since adenosine A_{2B} receptors have a low affinity for adenosine, high concentrations of the ligand (which are found during seizures or hypoxia) are required to activate this receptor. Thus, we propose that the following series of events are involved in the cross talk between IL-6 and the adenosinergic system during neuropathological conditions: (1) readily released adenosine activates low-affinity adenosine A_{2B} receptors in astrocytes that in response (2) secrete IL-6, which in turn (3) increases the expression adenosine A₁ receptors, and consequently (4) amplifies their neuroprotective properties (Figure 8).

According to the above-mentioned assumptions, high levels of extracellular adenosine are required to initiate the

process. It is worth to note that a significant correlation was found between the seizure scores in the first day and the induction of adenosine A₁ receptor expression in the hippocampus and cortex of WT mice, suggesting that only in animals with high-seizure scores a sufficient amount of adenosine was released to stimulate the synthesis of IL-6. Thus, the feed-forward loop now proposed (Figure 8) leading to amplification of adenosine A₁ receptor signaling might be particularly relevant *in vivo* to counteract pathological conditions associated with intense neuronal firing.

In conclusion, the data presented here demonstrate that IL-6 induces the expression of neuronal adenosine A₁ receptors *in vitro* and *in vivo*. This mechanism results in improved adenosinergic signaling and increased neuronal survival, and thus constitutes a molecular foundation of IL-6-induced neuromodulation and neuroprotection.

ACKNOWLEDGEMENTS

Work in Lisbon was supported by Fundação para a Ciência e Tecnologia (FCT). CC Fernandes and A Pinto-Duarte are in receipt of FCT fellowships (SFRH/BD/18046/2004/ and SFRH/BD/21589/2005). Work of K Biber and MC Witten-dorp was supported by school of Behavioral Cognitive Neurosciences (BCN, Groningen). We greatly acknowledge the gift of anti-adenosine A₁ receptor antibody by Dr Rafael Franco and support for western blot experiments by MJ Diógenes and the group of Dr João Barata at the IMM. We thank Prof B Hamprecht and Dr Alexandre de Mendonça for critical reading of the manuscript and valuable comments.

DISCLOSURE/CONFLICT OF INTEREST

The authors declare that they have no competing financial interests.

REFERENCES

- Abbracchio MP, Fogliatto G, Paoletti AM, Rovati GE, Cattabeni F (1992). Prolonged *in vitro* exposure of rat brain slices to adenosine analogues: selective desensitization of adenosine A₁ but not A₂ receptors. *Eur J Pharmacol* 227: 317–324.
- Ali C, Nicole O, Docagne F, Lesne S, MacKenzie ET, Nouvelot A et al (2000). Ischemia-induced interleukin-6 as a potential endogenous neuroprotective cytokine against NMDA receptor-mediated excitotoxicity in the brain. *J Cereb Blood Flow Metab* 20: 956–966.
- Anderson WW, Collingridge GL (2001). The LTP program: a data acquisition program for on-line analysis of long-term potentiation and other synaptic events. *J Neurosci Methods* 108: 71–83.
- Anderson WW, Collingridge GL (2007). Capabilities of the WinLTP data acquisition program extending beyond basic LTP experimental functions. *J Neurosci Methods* 162: 346–356.
- Angelatou F, Pagonopoulou O, Kostopoulos G (1991). Changes in seizure latency correlate with alterations in A₁ adenosine receptor binding during daily repeated pentylentetrazol-induced convulsions in different mouse brain areas. *Neurosci Lett* 132: 203–206.
- Angelatou F, Pagonopoulou O, Maraziotis T, Olivier A, Villemeure JG, Avoli M et al (1993). Upregulation of A₁ adenosine receptors in human temporal lobe epilepsy: a quantitative autoradiographic study. *Neurosci Lett* 163: 11–14.
- Arrigoni E, Crocker AJ, Saper CB, Greene RW, Scammell TE (2005). Deletion of presynaptic adenosine A₁ receptors impairs the recovery of synaptic transmission after hypoxia. *Neuroscience* 132: 575–580.
- Asanuma M, Ogawa N, Nishibayashi S, Kondo Y, Mori A (1995). Effects of repeated injection of cyclosporin A on pentylentetrazol-induced convulsion and cyclophilin mRNA levels in rat brain. *Neurochem Res* 20: 101–105.
- Barrie AP, Nicholls DG (1993). Adenosine A₁ receptor inhibition of glutamate exocytosis and protein kinase C-mediated decoupling. *J Neurochem* 60: 1081–1086.
- Bensadoun JC, Pereira de Almeida L, Dréano M, Aebischer P, Déglon N (2001). Neuroprotective effect of interleukin-6 and IL6/IL6R chimera in the quinolinic acid rat model of Huntington's syndrome. *Eur J Neurosci* 14: 1753–1761.
- Biber K, Klotz K-N, Berger M, Gebicke-Haerter PJ, van Calker D (1997). Adenosine A₁ receptor-mediated activation of phospholipase C in cultured astrocytes depends on the level of receptor expression. *J Neurosci* 17: 4956–4964.
- Biber K, Lubrich B, Fiebich BL, Boddeke HW, van Calker D (2001). Interleukin-6 enhances expression of adenosine A(1) receptor mRNA and signaling in cultured rat cortical astrocytes and brain slices. *Neuropsychopharmacology* 24: 86–96.
- Bolin LM, Strycharska-Orczyk I, Murray R, Langston JW, Di Monte D (2002). Increases vulnerability of dopaminergic neurons in MPTP-lesioned interleukin-6 deficient mice. *J Neurochem* 83: 167–175.
- Campbell IL, Abraham CR, Masliah E, Kemper P, Inglis JD, Oldstone MBA et al (1993). Neurologic diseases induced in transgenic mice by cerebral overexpression of interleukin 6. *Proc Natl Acad Sci* 90: 10061–10065.
- Carlson NG, Wieggl WA, Chen J, Bacchi A, Rogers SW, Gahring LC (1999). Inflammatory cytokines IL-1 α , IL-1 β , IL-6, and TNF- α impart neuroprotection to an excitotoxin through distinct pathways. *J Immunol* 163: 3963–3968.
- Chomczynski P, Sacchi N (1987). Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162: 156–159.
- Citri A, Malenka RC (2007). Synaptic plasticity: multiple forms, functions, and mechanisms. *Neuropsychopharmacology*, advance online publication, August 29, 2007 (DOI 10.1038/sj.npp.1301559).
- Conroy SM, Nguyen V, Quina LA, Blakely-Gonzales P, Ur C, Netzeband JG et al (2004). Interleukin-6 produces neuronal loss in developing cerebellar granule neuron cultures. *J Neuroimmunol* 155: 43–54.
- Dalpiaz A, Manfredini S (2002). Adenosine A₁ receptor: analysis of the potential therapeutic effects obtained by its activation in the central nervous system. *Curr Med Chem* 9: 1923–1937.
- d'Arcangelo G, Tancredi V, Onofri F, D'Antuono M, Giovedi S, Benfenati F (2000). Interleukin-6 inhibits neurotransmitter release and the spread of excitation in the rat cerebral cortex. *Eur J Neurosci* 12: 1241–1252.
- de Jong EK, Dijkstra IM, Hensens M, Brouwer N, van Amerongen M, Liem RSB et al (2005). Vesicle-mediated transport and release of CCL21 in endangered neurons: a possible explanation for microglia activation remote from a primary lesion. *J Neurosci* 25: 7548–7557.
- de Luca G, Di Giorgio R M, Macaione S, Calpona PR, Costantino S, Di Paola ED et al (2004). Susceptibility to audiogenic seizure and neurotransmitter amino acid levels in different brain areas of IL-6-deficient mice. *Pharmacol Biochem Behav* 78: 75–81.
- de Sarro G, Russo E, Ferreri G, Giuseppe B, Flocco MA, Di Paola E D et al (2004). Seizure susceptibility to various convulsant stimuli of knockout interleukin-6 mice. *Pharmacol Biochem Behav* 77: 761–766.

- de Mendonca A, Sebastiao AM, Ribeiro JA (1995). Inhibition of NMDA receptor-mediated currents in isolated rat hippocampal neurones by adenosine A₁ receptor activation. *Neuroreport* **6**: 1097–1100.
- Diógenes MJ, Fernandes CC, Sebastião A, Ribeiro JA (2004). Activation of adenosine A_{2A} receptor facilitates brain-derived neurotrophic factor modulation of synaptic transmission in hippocampal slices. *J Neurosci* **24**: 2905–2913.
- Dragunow M (1988). Purinergic mechanisms in epilepsy. *Prog Neurobiol* **31**: 85–108.
- Dunwiddie TV, Masino SA (2001). The role and regulation of adenosine in the central nervous system. *Annu Rev Neurosci* **24**: 31–55.
- Fiebich BL, Biber K, Gyufko K, Berger M, Bauer J, van Calcar D (1996). Adenosine A_{2B} receptors mediate an increase in interleukin (IL)-6 mRNA and IL-6 protein synthesis in human astroglia cells. *J Neurochem* **66**: 1426–1431.
- Gadient RA, Otten UH (1997). Interleukin-6 (IL-6)-a molecule with both beneficial and destructive potentials. *Prog Neurobiol* **52**: 379–390.
- Gerber U, Gahwiler BH (1994). GABA_B and adenosine receptors mediate enhancement of the K⁺ current, IAHP, by reducing adenylyl cyclase activity in rat CA3 hippocampal neurons. *J Neurophysiol* **72**: 2360–2367.
- Gouder N, Fritschy JM, Boison D (2003). Seizure suppression by adenosine A₁ receptor activation in a mouse model of pharmacoresistant epilepsy. *Epilepsia* **44**: 877–885.
- Gruol DL, Nelson TE (1997). Physiological and pathological roles of interleukin-6 in the central nervous system. *Mol Neurobiol* **15**: 307–339.
- Herrmann O, Tarabin V, Suzuki S, Attigah N, Coserea I, Schneider A et al (2003). Regulation of body temperature and neuroprotection by endogenous interleukin-6 in cerebral ischemia. *J Cereb Blood Flow and Metab* **23**: 406–415.
- Hettinger BD, Leid M, Murray T F (1998). Cyclopentyladenosine-induced homologous down-regulation of A₁ adenosine receptors (A₁AR) in intact neurons is accompanied by receptor sequestration but not a reduction in A₁AR mRNA expression or G protein alpha-subunit content. *J Neurochem* **71**: 221–230.
- Jankowsky JL, Patterson PH (1999). Differential regulation of cytokine expression following pilocarpine-induced seizure. *Exp Neurol* **159**: 333–346.
- Kondziella D, Bidar A, Urfjell B, Sletvold O, Sonnewald U (2002). The pentylentetrazole-kindling model of epilepsy in SAMP8 mice: behavior and metabolism. *Neurochem Int* **40**: 413–418.
- Lambert NA, Teyler TJ (1991). Adenosine depresses excitatory but not fast inhibitory synaptic transmission in area CA1 of the rat hippocampus. *Neurosci Lett* **122**: 50–52.
- Latini S, Pedata F (2001). Adenosine in the central nervous system: release mechanisms and extracellular concentrations. *J Neurochem* **79**: 463–484.
- Lehtimäki KA, Peltola J, Koskikallio E, Keranen T, Honkaniemi J (2003). Expression of cytokines and cytokine receptors in the rat brain after kainic acid-induced seizures. *Brain Res Mol Brain Res* **110**: 253–260.
- Lehtimäki KA, Keränen T, Huhtala H, Hurme M, Ollikainen J, Honkaniemi J et al (2004). Regulation of IL-6 system in cerebrospinal fluid and serum compartments by seizures: the effect of seizure type and duration. *J Neuroimmunol* **152**: 121–125.
- Livak KJ, Schmittgen TD (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔC(T)}. *Method Methods* **25**: 402–408.
- Lucchi R, Latini S, de Mendonça A, Sebastião AM, Ribeiro JA (1996). Adenosine by activating A₁ receptors prevents GABA_A-mediated actions during hypoxia in the rat hippocampus. *Brain Res* **732**: 261–266.
- Loddick SA, Turnbull AV, Rothwell NJ (1998). Cerebral interleukin-6 is neuroprotective during permanent focal cerebral ischemia in the rat. *J Cereb Blood Flow Metab* **18**: 176–179.
- Lohse MJ, Lenschow V, Schwabe U (1984). Interaction of barbiturates with adenosine receptors in rat brain. *Naunyn Schmiedebergs Arch Pharmacol* **326**: 69–74.
- Marchetti L, Klein M, Schlett K, Pfizenmaier K, Eisel UL (2004). Tumor necrosis factor (TNF)-mediated neuroprotection against glutamate-induced excitotoxicity is enhanced by N-methyl-D-aspartate receptor activation. Essential role of a TNF receptor 2-mediated phosphatidylinositol 3-kinase-dependent NF-kappa B pathway. *J Biol Chem* **279**: 32869–32881.
- Nelson TE, Netzeband JG, Gruol DL (2004). Chronic interleukin-6 exposure alters metabotropic glutamate receptor-activated calcium signalling in cerebellar Purkinje neurons. *Eur J Neurosci* **20**: 2387–2400.
- Onodera H, Kogure K (1988). Differential localization of adenosine A₁ receptors in the rat hippocampus: a quantitative autoradiographic study. *Brain Res* **458**: 212–217.
- Pagonopoulou O, Angelatou F, Kostopoulos G (1993). Effect of pentylentetrazol-induced seizures on A₁ adenosine receptor regional density in the mouse brain: a quantitative autoradiographic study. *Neuroscience* **56**: 711–716.
- Peltola J, Hurme M, Miettinen A, Keranen T (1998). Elevated levels of interleukin-6 may occur in cerebrospinal fluid from patients with recent epileptic seizures. *Epilepsy Res* **31**: 129–133.
- Peltola J, Laaksonen J, Haapala AM, Hurme M, Rainesalo S, Keranen T (2002). Indicators of inflammation after recent tonic-clonic epileptic seizures correlate with plasma interleukin-6 levels. *Seizure* **11**: 44–46.
- Peltola J, Palmio J, Korhonen L, Suhonen J, Miettinen A, Hurme M et al (2000). Interleukin-6 and interleukin-1 receptor antagonist in cerebrospinal fluid from patients with recent tonic-clonic seizures. *Epilepsy Res* **41**: 205–211.
- Penkowa M, Molinero A, Carrasco J, Hidalgo J (2001). Interleukin-6 deficiency reduces the brain inflammatory response and increases oxidative stress and neurodegeneration after kainic acid-induced seizures. *Neuroscience* **102**: 805–818.
- Pinto-Duarte A, Coelho JE, Cunha RA, Ribeiro JA, Sebastiao AM (2005). Adenosine A_{2A} receptors control the extracellular levels of adenosine through modulation of nucleoside transporters activity in the rat hippocampus. *J Neurochem* **93**: 595–604.
- Pizzi M, Sarnico I, Boroni F, Benarese M, Dreno M, Garotta G et al (2004). Prevention of neuron and oligodendrocyte degeneration by interleukin-6 (IL-6) and IL-6 receptor/IL-6 fusion protein in organotypic hippocampal slices. *Mol Cell Neurosci* **25**: 301–311.
- Qiu Z, Sweeney DD, Netzeband JG, Gruol DL (1998). Chronic interleukin-6 alters NMDA receptor-mediated membrane responses and enhances neurotoxicity in developing CNS neurons. *J Neurosci* **18**: 10445–10456.
- Ribeiro JA (2005). What can adenosine neuromodulation do for neuroprotection? *Curr Drug Targets CNS Neurol Disord* **4**: 325–329.
- Rudolph KA, Schubert P (1996). Purinergic interventions in traumatic and ischemic injury. In: Peterson PL, Willis PW (eds) *Novel Therapies for CNS Injuries*. Boca Raton: New York, London, Tokyo: CRC Press Inc, pp 327–342.
- Ruiz A, Sanz JM, Gonzalez-Calero G, Fernandez M, Andres A, Cubero A et al (1996). Desensitization and internalization of adenosine A₁ receptors in rat brain by *in vivo* treatment with R-PIA: involvement of coated vesicles. *Biochim Biophys Acta* **1310**: 168–174.
- Samland H, Huitron-Resendiz S, Masliah E, Criado J, Hendriksen SJ, Campbell IL (2003). Profound increase in sensitivity to glutamatergic- but not to cholinergic agonist-induced seizures in transgenic mice with astrocyte production of IL-6. *J Neurosci Res* **73**: 176–187.

- Schurr A, Reid KH, Tseng MT, West C, Rigor BM (1986). Adaptation of adult brain tissue to anoxia and hypoxia *in vitro*. *Brain Res* 374: 244–248.
- Schwaninger M, Neher M, Viegas E, Schneider A, Spranger M (1997). Stimulation of interleukin-6 secretion and gene transcription in primary astrocytes by adenosine. *J Neurochem* 69: 1145–1150.
- Schwaninger M, Petersen N, Prinz S, Sallmann S, Neher M, Spranger M (2000). Adenosine-induced expression of interleukin-6 in astrocytes through protein kinase A and NF-IL-6. *Glia* 31: 51–58.
- Sebastião AM, Cunha RA, de Mendonça A, Ribeiro JA (2000). Modification of adenosine modulation of synaptic transmission in the hippocampus of aged rats. *Br J Pharmacol* 131: 1629–1634.
- Sebastião AM, de Mendonça A, Moreira T, Ribeiro JA (2001). Activation of synaptic NMDA receptors by action potential-dependent release of transmitter during hypoxia impairs recovery of synaptic transmission on reoxygenation. *J Neurosci* 21: 8564–8571.
- Sebastião AM, Stone TW, Ribeiro JA (1990). The inhibitory adenosine receptor at the neuromuscular junction and hippocampus of the rat: antagonism by 1,3,8-substituted xanthines. *Br J Pharmacol* 101: 453–459.
- Staff NP, Jung HY, Thiagarajan T, Yao M, Spruston N (2000). Resting and active properties of pyramidal neurons in subiculum and CA1 of rat hippocampus. *J Neurophysiol* 84: 2398–2408.
- Stone TW (2002). Purines and neuroprotection. *Adv Exp Med Biol* 513: 249–280.
- Swanson TH, Drazba JA, Rivkees SA (1995). Adenosine A₁ receptors are located predominantly on axons in the rat hippocampal formation. *J Comp Neurol* 363: 517–531.
- Swartz KR, Liu F, Sewell D, Schochet T, Campbell I, Sandor M et al (2001). Interleukin-6 promotes post-traumatic healing in the central nervous system. *Brain Res* 896: 86–95.
- Trussell LO, Jackson MB (1985). Adenosine-activated potassium conductance in cultured striatal neurons. *Proc Natl Acad Sci USA* 82: 4857–4861.
- Ulas J, Brunner LC, Nguyen L, Cotman CW (1993). Reduced density of adenosine A₁ receptors and preserved coupling of adenosine A₁ receptors to G proteins in Alzheimer hippocampus: a quantitative autoradiographic study. *Neuroscience* 52: 843–854.
- Vanore G, Giraldez L, Rodriguez de Lores Arnaiz G, Girardi E (2001). Seizure activity produces differential changes in adenosine A₁ receptors within rat hippocampus. *Neurochem Res* 26: 225–230.
- van Wagoner NJ, Benveniste EN (1999). Interleukin-6 expression and regulation in astrocytes. *J Neuroimmunol* 100: 124–139.
- Wittendorp MC, van Frijtag Drabbe Kunzel J, Ijzerman AP, Boddeke HWGM, Biber K (2004). The mouse brain adenosine A₁ receptor: functional expression and pharmacology. *Eur J Pharmacol* 487: 73–79.
- Yan L, Burbiel JC, Maass A, Muller CE (2003). Adenosine receptor agonists: from basic medicinal chemistry to clinical development. *Expert Opin Emerg Drugs* 8: 537–576.
- Yoon KW, Rothman SM (1991). Adenosine inhibits excitatory but not inhibitory synaptic transmission in the hippocampus. *J Neurosci* 11: 1375–1380.