Iodine*

By Prof. Irvine Masson

OF the two aspects of iodine and its chemistry which are here surveyed, the first is that of the part played by the element in terrestrial economy, whether inorganic or organic or both. In the total make-up of the earth, iodine is not one of the fourteen elements which compose 99.95 per cent of the mass, but is one of the rarest of the non-metallic elements. Nevertheless, it is scattered almost everywhere, in minute traces; a ton of average rock contains some 300 milligrams, a ton of sea water about 17 milligrams.

The chemical reactions of geology have very rarely segregated any of these widely dispersed traces; there are scarcely any minerals containing iodine as an essential constituent, and the few that have been found are out-of-the-way mineralogical curiosities, small specimens rather than deposits or ores. Only in one place in the world have solid iodine compounds come together in quantity, and that is in the nitrate deposits of Chile, high up against the western slopes of the Andes. There the element occurs as inorganic iodate, and there are uncounted thousands of tons of it. Yet, even so, the iodate is not by itself in veins or beds, but is simply one of the minor impurities in the great accumulations of sodium nitrate. We might be able to explain how the iodate got there if we knew how the nitrate had got there; but, in spite of much interesting work, all that seems really safe to say about the origin of Chile nitrate is that no single hypothesis has yet met all the known facts. What is indubitably true is that the world's commercial supply of iodine (with trifling exceptions) comes from the Chile nitrate factories, from which it is exported at the rate of nearly 1,000 tons of the element yearly.

It seems from these facts that during the 2,000 million years which span geochemistry, iodine has played a part that was practically nothing, and that had the element never been created, the inorganic earth would not be appreciably different from what it is. But with the origination of organic life (800 million years ago, or earlier), iodine began to matter. First it became not wholly dispersed, but partly to be segregated from solution in the sea into the colloids of living organisms floating there. Then, as the forms of living things developed, the iodine was not only taken up into them, but

also became a chemical part of some of them. They developed still further and more variously; and now, the vertebrates, and we ourselves, could not continue without iodine, either as individuals from month to month or as species from generation to generation.

Looking more closely into this from a biochemical point of view, we learn from many recorded analyses of vegetable and animal materials that three living creatures stand out far beyond the rest as being rich in iodine. These three are seaweeds (brown and red), bath sponges, and certain classes of corals. One of these is vegetable, two are animal, but all three are marine; and this is so notwithstanding the fact that the sea water is excessively poor in iodine compared, for example, with many soils and most rocks. The iodine content of these organisms is thousands or even millions of times more than in the sea water in which they live; it may reach one half per cent of the weight of the dry tissue in seaweed, one per cent or more in the sponges, and in a coral as much as eight per cent. The chemical form of this iodine is undecided in the case of seaweed; but as to the horny sponges and horny corals, the work of Drechsel (1896) and of Wheeler and his associates (1905-7), confirmed by that of Mörner (1907-13), showed, surprisingly, that the skeletal protein releases its iodine as the definite aromatic amino-acid, 3:5 di-iodo-tyrosine:

3:5 DI-iodo-TYROSINE

It is improbable that this substance is supplied to the animal cells ready-made in their plant food; such food no doubt brings them the common protein ingredient, tyrosine, but the iodination would seem to go on in the cell. For this purpose, the very minute traces of dissolved inorganic iodine salts in sea water are drawn upon; and if these are iodides, the co-operation of an oxidizing agent is obviously necessary (an oxidase naturally suggests itself). The direct iodination of tyrosine in the laboratory, with elementary iodine as the

* Substance of a Friday evening discourse at the Royal Institution, delivered on December 3, 1937.

reagent, was easily accomplished by Oswald (1909) and has lately been studied in more detail, as a part of protein chemistry, by Bauer and Strauss (1935).

It is particularly to be noted that organic halogen compounds of any kind are extraordinarily rare among natural products. The only other cases known at present are: di-bromo-tyrosine, obtained by Möerner (1913) from horny corals, and evidently formed in the same way as the di-iodo-compound; di-bromo-indigo, the classical Tyrian purple dye from the glands of the shellfish *Murex* or *Purpura* (Friedlander, 1909, 1922); certain aromatic chloro-compounds lately isolated from lichens (Nolan, 1934; Pfau, 1934); and other chloro-compounds which, as Raistrick has shown (1936), are formed by certain moulds growing in solutions of inorganic chloride.

No experiments seem to have been made in order to learn whether the addition of iodine compounds to sea water, or their removal, affects the growth of those marine organisms in which iodine is found organically combined. It is worth noticing, however, that the iodine compound is accumulated not in the fleshy cell-tissue of the sponge or the coral, but in the horny skeleton on which these living cells sit; and this skeleton, though mechanically useful to the cell colony, may almost be called an excretory product of that colony. It is for the animal physiologist to decide whether in these cases the skeletal protein is serving as a mere dump for di-iodo-tyrosine that has been discarded by the cells as useless, or whether, on the contrary, it acts as a reserve store upon which the cells can draw for part of some vital process. This question, unsettled though it is, is not without significance in the light of the biochemistry of animals higher up in the evolutionary scale.

Sponges and corals are lowly creatures, with small differentiation of tissues and no co-ordination of parts, and they existed even in Cambrian times, long before forms of land life were evolved. Making, then, a very wide biological jump to the vertebrates, including man, we again find the faculty for the organic fixation of iodine, and it is localized now in the thyroid gland. That it here determines the metamorphosis of amphibian larvæ, such as tadpoles and axolotls, was first shown in 1912; and the well-established correlation of inadequate traces of iodine in food with irregularity in the function of the gland, and consequently with the disturbances of basal metabolism which cause goitre, imperfect growth, myxœdema, or cretinism, was given a rational basis first by Kendall's isolation (1915) of the thyroid hormone as a crystalline organic iodine compound which he called thyroxine, and then by the brilliant work of C. R. Harington and his

associates (1926 *et seq.*), to whom we owe our knowledge of what thyroxine really is.

THYROXINE

It appears that the iodine atoms in this molecule are essential to the physiological potency of the substance; and a teleologist might well find in this hormone molecule the sole *raison d'être* of the element iodine.

Harington has shown that thyroid iodine is present partly as thyroxine but also as di-iodo-tyrosine, both amino-acids being chemical components of the complex protein thyroglobulin. The direct physiological activity of the di-iodo-tyrosine appears, so far as published evidence goes, to be relatively slight; but there cannot be much doubt that it is the precursor of thyroxine in the chemical synthesis for which the thyroid gland is responsible. The conversion into thyroxine, indicated in a purely formal way by

has not been accomplished in the laboratory (the pioneering synthesis of thyroxine by Harington and Barger necessarily used materials and methods not natural to living tissues), but it would involve steps of a kind not unknown in other branches of protein chemistry.

To sum up this part of our subject: the opportunity for the natural fixation of inorganic iodine into an organic form has in all ages been extremely slight; not because the process is intrinsically difficult (for it is not), but because the supply of inorganic iodine has always been extremely scanty. Accordingly, iodine is not one of those elements which, like sulphur or phosphorus, became normal ingredients of common proteins. Living things had next to no iodine to draw upon, so most of

them managed and evolved without it. But very early in the history of multicellular animals in the sea—if we may make an inference from those simple forms which have survived into this age—the organic fixation and accumulation of iodine was accomplished by some of them, and what they did with it was to make it into di-iodo-tyrosine. This same faculty of making di-iodo-tyrosine, we find, has been preserved, or at any rate repeated, by the vertebrate animals, localized in their thyroid glands. Besides that faculty, we have put our finger upon one further chemical step, which has apparently accompanied the evolution of the vertebrates, is characteristic of them, and is indispensable to their being: namely, the final conversion of di-iodo-tyrosine into thyroxine.

That iodine is a biochemical element is a fact which lends importance to its organic chemistry in general; and in what follows, one part of this will be discussed, namely, that in which organic iodine is multivalent. In this field, some is not new but is perhaps undeveloped, while the rest is new; and to illustrate it, we shall confine ourselves here to a few connected examples drawn from work done at Durham (*J. Chem. Soc.*, 1669; 1935. 1718; 1937. *NATURE*, 139, 150; 1937). Throughout, the chemist has to bear in mind that iodine is not merely the heaviest halogen, but—especially when multivalent—is also the terminal member of its series in the Periodic System, and its behaviour is to be regarded accordingly.

Victor Meyer and Wachter's discovery of iodosobenzoic acid (1892) prompted Willgerodt to study iodobenzene dichloride, which he had isolated seven years before, and so to discover iodosobenzene and, ultimately, many other aromatic iodoso-compounds. Besides behaving as bases towards acids, these derivatives of tervalent iodine exhibit self-oxidation-reduction, as do nearly all inorganic compounds in which a halogen atom is in an intermediate state of valency; so iodosobenzene spontaneously turns into iodo- and iodoxybenzene, which latter compound has reactions of general chemical interest.

Iodoxybenzene, $C_6H_5IO_2$, is not only formally analogous to nitrobenzene, $C_6H_5NO_2$, but also by its chemical behaviour shows the internal electrical polarity which is known, on a variety of evidence, to exist in nitrobenzene. The respective donor atoms iodine and nitrogen are positive, the acceptor oxygen atoms are negative. Hence an iodoxy-compound is attacked either by negatively charged ions or parts of other molecules, which its positive iodine attracts, or else by positively charged ions or groups, which its negative oxygen atoms attract. In nitrobenzene, the small donor nitrogen atom is screened by the phenyl and by

the two oxygen atoms, and is therefore not exposed to negatively charged reagents as is the large iodine; but its negative oxygen atoms are open. Thus, both compounds form salts with

Drawings of four successive stages of a model showing the expulsion of the iodate ion. The diagram illustrates the general nature of the other models also. The backgrounds were black. Positive ions were marked by the emission of a bright red light from the group of atoms; negative ions by that of a bright blue light (indicated by a small conventional 'explosion' sign). Polarities within molecules were shown by a red glow over the disk of a positive (donor) atom and a blue glow over the disk of a negative (acceptor) atom, as suggested in the drawing of the molecule of iodoxybenzene on the right in the first stage.

strong acids, by the co-ordinating (with resonance) of a proton with their oxygen atoms; for example, the crystalline hydrogen-sulphates:

Iodoxybenzene dissolves in alkali, owing to the attachment of a negative hydroxyl ion to the

positive iodine, yielding the unstable solute, the *phenyl-iodoxylate* of the alkali metal :

Nitrobenzene, for the reason explained above, does not combine with alkali ; but the arylphosphonic and arsonic acids are probably analogous to the iodoxyates.

When the solution of an alkaline iodoxyate is boiled, its complex negative ion extrudes the group IO_3 which, as the iodate ion, carries off the ionic charge, leaving behind benzene :

This elimination of the iodate ion is characteristic of such substances, especially in alkali. Thus, in a cold iodoxyate solution, the complex ion spontaneously co-operates with dissolved iodoxybenzene to do so, and the residues couple to form *diphenyliodyl hydroxide* (or its alkali salt), with the structure of which one may compare the diaryl phosphinic and arsonic acids :

Again, the same dominant tendency is seen in the

formation of iodonium compounds by Victor Meyer and Hartmann's second method, where iodoso- and iodoxy-compound react together in an alkaline medium :

The converse of these processes, that is, the return of iodine from inorganic iodate into the organically combined state, proves easy to realize in the laboratory, under conditions of high acidity, organic derivatives of multivalent iodine being formed.

It will be understood that in these examples of reactions (all of which, in the discourse of which this is a condensation, were illustrated by working scale-models showing the atomic and electrical exchanges), the detailed intermediate mechanisms have here been slightly simplified, especially as regards the organic parts of the groups concerned. In them, however, the primary point is rather the formation and existence of hitherto unknown types of iodine compounds. These lead to a new field of chemical inquiry, and help to show, as do the biochemical facts reviewed, that we must think of iodine as an element which, like nitrogen or arsenic, is not inorganic alone, but is emphatically an organic element also.

Scientific Research in India

From a Correspondent

THE activities of the scientific delegation in India were by no means confined to the proceedings of the Congress in Calcutta and to sight-seeing. Every opportunity was taken by individual members or groups of members to inform themselves of the progress of science in the various institutions which occasion could be made to visit. The notes which follow cannot be exhaustive, as they are based upon information only from members of the party who have just returned to England in the S.S. *Strathaird* ; a substantial number of the delegates are not yet home.

The discovery of the pre-Aryan civilization of the Indus basin has stimulated studies in Indian archaeology, and the delegation was profoundly impressed by the care bestowed upon the famous frescoes and sculptures in the rock-hewn temples at Ajanta and Ellora, while excavation is in progress elsewhere in the dominions of H.E.H. the Nizam of Hyderabad. The museums at Delhi, Bombay, Sarnath (near Benares), Calcutta and Madras all possess notable treasures, and that at

Calcutta especially needs much greater space for its adequate exposition in order to guide future excavation. Turning to work within the scope of present-day anthropology, the Brahmachariya Sangha founded by the late Maharajah of Kazimbazar illustrates a type of Indian organization designed to help especially aboriginal peoples to overcome handicaps due to lack of knowledge of Hindu or Dravidian languages and to general backwardness. The aborigines around Ranchi on the Chota Nagpur plateau are being taught on the basis of a modernist interpretation of the Vedas, and schools, clinics and legal aid come within the province of this organization.

Elsewhere, the great medical institutions naturally attracted attention. At the famous Haffkine Institute in Bombay special note was taken of the work in progress towards a theory as to outbreaks of plague after periods of immunity ; it was learned incidentally that mice are found more suitable than rats for plague experimentation. The work on rabies was also remarked ; and it was