tion of both the reflected and the transmitted beams. The reflecting power is, in effect, confined to a crystal skin of thickness about 0.001 mm., and the remainder of the crystal is quite ineffective. The exact parallelism of the crystal planes also ensures that only a very small range of incident angles is effectively employed. Both these factors contribute to what is known as the *primary extinction* of perfect crystals.

But most 'single crystals' are mosaics of small crystallites, orientated more or less similarly but with small discontinuities of spacing or direction at frequent intervals. If the crystallites are large and disorientated, or if there is, for example, a regular change of orientation within the single crystal, then Laue photographs are useful. They have shown, for example, that slight spiralling is not uncommon as the result of natural growth or mechanical treatment⁸. If the crystallites are extremely small, diffuse scattering will sometimes reveal their size and shape (as in the case of the micelles of fibres, or the segregation of new and incipient crystalline phases from certain alloys during heat-treatment or age-hardening). Crystallites of medium size, say 0.001–0.00001 mm. linear dimensions, are not so easy to study. They do not show primary extinction, because there are no phase relationships between the rays scattered by one crystallite and another; they are 'all skin'. But the total reflecting power of the composite crystal is less than the sum of the reflecting powers of the individual crystallites would be if each were separately bathed in the incident X-ray beam appropriate to its position. This is because, in the case of the composite crystal, the incident beam is reduced by reflexion as well as by ordinary absorption before it reaches the lower layers of crystallites. *Secondary extinction*, as this diminution of total reflecting power of a mosaic crystal is called, is therefore complementary to the effective increase in absorbing power of the crystal for rays which meet the atomic planes at the selective reflexion angles, and it varies for different planes according to the degree of multiple reflexion that can take place, that is, according to the degree of parallelism of individual crystallites.

By studying the absorbing power of the crystal at the Bragg angles, therefore, it is possible to obtain useful information about the crystallite arrangement. The white lines on a grey background observed when a divergent X-ray beam is transmitted through a single crystal⁹ are, in fact, a record of this selective absorption at the Bragg angles, and crystals of different texture are found to give patterns which, though geometrically equivalent, are of very different quality. Really perfect crystals give no pattern, since their reflecting power is so very small. It has been rather surprisingly found that even organic compounds sometimes grow as perfect crystals, which give no divergent beam pattern until they are broken up into crystallites by sudden immersion in liquid air. Ice also forms very regular crystals even at temperatures only a little below 0° C., when the thermal vibrations are exceptionally strong. Using coarse-grained film, divergent beam patterns can be obtained with exposures of only a few seconds at, say, 5 cm. crystal to film distance; but the crystals must not be so thin that their reflecting power is small, or so thick that complete absorption occurs at all angles. The optimum thickness for diamond, ice or organic compounds, using radiation from a copper target, is about 1 or 2 mm. Patterns can only be obtained from crystals of high atomic weight if short-wave radiation is used.

Some of the difference in texture between different diamonds may be due to the presence of foreign atoms. Silicon, aluminium, calcium and various other atoms of relatively high atomic weight are known to occur in diamond, and it is quite likely that hydrogen, boron and nitrogen may also be present. It was of interest to find out whether these would produce any change in the lattice constant of individual diamonds. The divergent beam method provided an excellent method of investigation, for photographs could be taken at long distances without impossibly long exposures, and the positions of certain points of intersection of the absorption lines on the films were extremely sensitive to small changes in λ/a . The wave-length, λ , being accurately known for copper $K\alpha_1$ radiation, the lattice constant a could be determined with an accuracy of about 1 in 70,000. It was in fact found that four different diamonds studied had values of a varying between the limits 3.56000 and 3.55942 kX., corresponding to a C–C distance of 1.54153–1.54127 kX. (or 1.54465–1.54440 Å.). All precision measurements made in other laboratories, whether for single diamonds or for diamond powder, fall within these limits.

¹ Lonsdale, K., Reports on Progress in Physics (1942–43), **9**, 256, for collected references to experimental work.

² Born, M., Reports on Progress in Physics (1942–43), **9**, 294, for collected references to theoretical work.

³ Müller, A., *Proc. Roy. Soc., A*, **174**, 137 (1940); *A*, **178**, 227 (1941).

⁴ Lonsdale, K., Robertson, J. M., and Woodward, I., *Proc. Roy. Soc., A*, **178**, 43 (1941). Knaggs, J. E., and Lonsdale, K., *J. Chem. Soc.*, 417 (1942).

⁵ Robertson, J. M., and Woodward, I., *J. Chem. Soc.*, 36 (1940). Ubbelohde, A. R., and Woodward, I., *Proc. Roy. Soc., A*, **181**, 415 (1943).

⁶ Ubbelohde, A. R., and Woodward, I., *Nature*, **155**, 170 (1945) *Proc. Roy. Soc.*, in the press.

⁷ Ubbelohde, A. R., and Woodward, I., *Nature*, **156**, 20 (1945).

⁸ Lonsdale, K., *Mineral. Mag.*, **27**, 112 (1945).

⁹ Lonsdale, K., *Nature*, **151**, 52 (1943); **153**, 22, 433 (1944); *Phil. Trans. Roy. Soc.*, in the press.

INDUSTRIAL DERMATITIS

INDUSTRIAL dermatitis is a special branch of a specialized subject. It receives little attention in the systematic teaching of medicine, and is considered with diffidence by the general practitioner because of its medico-legal implications. Its proper study is confined largely to the dermatologists and to those physicians employed in industrial medicine.

The clinical manifestations of industrial dermatitis are often indistinguishable from those of eczema, and accurate diagnosis is a matter involving special experience as well as dermatological skill. Occupational dermatitis is not confined to artisans. It can be found in all walks of life, as for example in dentists, who may produce a dermatitis due to local anaesthetic solutions, or in medical practitioners, who may be sensitive to certain antiseptics.

Investigation of any case of dermatitis must be very thorough and include inquiry into all the patient's activities at work and in leisure, as the cause of dermatitis may lie not only in the substances of industry but also in things encountered in the home or garden. Psychological factors in the production of skin diseases are gradually being brought to light, and when this subject is better understood it will probably be possible further to classify the types of dermatitis.

The number of substances known to cause dermatitis grows in proportion to the rate of introduction of new processes in industry, and the war years

have added much to our knowledge. Already cases of dermatitis caused by penicillin solution have been reported as occurring in medical men and others handling it. Unfortunately the growth of our knowledge is mainly of theoretical value, and progress in prevention and in therapeutics is not so marked.

The large number of cases of industrial dermatitis and the inadequacy of dermatological facilities at the present time make it necessary for the long-suffering general practitioner to undertake, in most cases, if not the diagnostic procedures, at least the after-care of such patients.

Dr. Archibald Gray's survey* of the social aspects of industrial dermatitis is of the greatest value in illustrating some points which are not apparent to the average general practitioner who may see only a few cases each year, and which are not to be found in the standard text-books on skin diseases. He points out that the terms, 'industrial dermatitis' and 'occupational dermatitis', are not synonymous, and that industrial dermatitis defines those cases of occupational dermatitis occurring in persons coming under the Workmen's Compensation Act and described officially as 'dermatitis produced by dust or liquids'. Thus the domestic servant with dermatitis due to excessive use of soap and water can claim compensation and is a case of industrial dermatitis, while her mistress, who might equally suffer from a similar condition, would rank as a case of occupational dermatitis.

Industrial diseases of the skin, apart from 'dermatitis produced by dust or liquids', include (1) ulceration due to dust or liquids; (2) epitheliomatous cancer or ulceration due to tar, pitch, bitumen, mineral oil or any compound, product or residue of any of these substances; (3) chrome ulceration or its sequelae; (4) scrotal epithelioma (chimney-sweeps' cancer); (5) inflammation, ulceration or malignant disease of the skin or subcutaneous tissues due to exposure to X-rays or radioactive substances.

Industrial dermatitis differs from other industrial diseases in that it can be produced by a great variety of agents; but the eruption so produced gives no clue to the actual agent involved, save that a follicular inflammation suggests tar, oil or chlorinated naphthalene. The incidence of industrial dermatitis is not accurately known; but Dr. Gray suggests, not without reason, that if the manpower loss could be computed it would be a startling figure.

Most cases of industrial dermatitis are associated with damage to the horn layer of the skin. The irritating substances penetrate this layer, damage the cells from which it is formed, and by thus interfering with its production, further increase the liability to trauma. The protective power of the horn layer is naturally lower in individuals whose oil glands are underactive. The more actual irritants are mostly met with in special trades, and factory surgeons and H.M. inspectors of factories are usually quick to recognize irritant properties in new substances used in industry, and to find ways of eliminating or reducing the risks incurred. A large group of cases caused by common irritants, such as soap, soda and water, lime and cement, etc., are impossible of control. The fact that the majority of handlers of potentially irritating substances escape dermatitis requires explanation and raises the subject of relative

sensitivity. Hebra showed long ago that certain irritants, if left long enough in contact with normal skin, would produce eczematous dermatitis. The determining factors are the intensity of the irritant and the time of exposure. Dermatitis may arise in the normal skin from mild irritants if cleansing is imperfect or if impregnated clothes are constantly worn. Cleansing substances themselves (for example, turpentine used to remove grease from the skin) may cause dermatitis. In other cases the skin is not normally resistant. Rarely there is sensitivity to one specific substance, and the best example is of the gardener who is sensitive to *Primula obconica*. A common factor in those two groups is that removal of the irritant is followed by healing of the disease. In the first group, the individual can return to work, usually without danger, provided he washes properly and changes his clothes after work. Those in the second group, with an individual sensitivity, must avoid contact with the known irritant, but can otherwise do ordinary work.

This is not so in most cases of eczematous dermatitis where removal of the man from work is not at once followed by healing and where, indeed, spread or new lesions may occur after all exposure to the irritant has ceased. Most of these cases are in men over forty, many of whom have been in the same trade for years. Idiopathic eczema often occurs at this age, and MacCormac's opinion is that "many cases of alleged occupational eczema about this age are probably idiopathic eczema or alternatively examples of idiopathic eczema prematurely induced by the occupation, but maintained by the diathesis which determines eczema". These people show a loss of skin resistance due to age and unknown factors, possibly vascular or metabolic. This is borne out by the fact that after the original attack has cleared, return to work which occasions exposure to any irritant, not necessarily the one which caused the first attack, will almost certainly cause a recurrence.

Compensation under the Workman's Compensation Act is payable if the area examining surgeon, appointed by the Ministry of Labour, decides that dermatitis is due to a man's work; but employers have a right of appeal to a medical referee, whose decision is final. There is usually no delay in settlement, save when the workman is ignorant of procedure or when legal difficulties arise in deciding who shall pay compensation when a man has had more than one employer within the previous year.

Failure to respond to treatment has in some cases a profound psychological effect on the sufferer, as, for example, when a skilled worker is not sure whether he can return to his own trade. If this is to be closed to him it may mean a reduction in wages should he have to return to the ranks of unskilled labour. The fate of those certified as suffering from industrial dermatitis is thus classified by Dr. Gray: (1) some cases respond readily to treatment and can return to work if they take reasonable precautions; (2) others respond to treatment, and, though they cannot return to their original employment, can find work in other trades; (3) a last group includes cases who do not readily respond to treatment and whose condition is complicated by an anxiety neurosis which keeps up the skin condition and makes them unfit for work of any kind for a long period.

Preventive measures in industry are comprehensive and include proper ventilation and lighting of works, mechanical removal of dusts, protective clothing, provision of adequate washing facilities, the use of

* "Some Social Aspects of Industrial Dermatitis." The Malcolm Morris Memorial Lecture, delivered at the Westminster Hospital Medical School on November 29, 1945, by Dr. Archibald M. H. Gray, under the auspices of the Chadwick Trust.

barrier creams, etc. Some people, especially liable to dermatitis (for example, sufferers from eczema, people with very dry skins, etc.), can be weeded out in advance and prevented from undertaking employment which, for them, is hazardous. Skin tests can, in rare cases, be used to find susceptible individuals.

The treatment of established dermatitis falls usually to the general practitioner, and Dr. Gray regrets the fact that the teaching of dermatology to medical students at the present time is inadequate and that often attendances at skin clinics are perfunctory. Should the general practitioner wish to send his patient to a consultant, this is often difficult and involves delay. Mental and physical rest are the first essentials of treatment, and these are only attainable in hospital. The number of hospital beds at present available for cases of skin disease is grossly inadequate. Much could be done to reduce time lost by sufferers from dermatitis if there were good facilities for out-patient and in-patient hospital treatment and for the rehabilitation of patients in the period after leaving hospital and before they are fit to resume work. Part of the problem of anxiety would be solved if the workman's financial worries could be allayed by an assurance that compensation would not be delayed and would continue as long as he remained incapable of full employment.

Dr. Gray ends on a plea for increased facilities for dermatological research and outlines his conception of the problems to be studied. J. MARSHALL.

GEOLOGY OF THE SALT RANGE OF THE PUNJAB

THE Salt Range of the Punjab runs with arcuate plan for some hundred miles eastward from the Indus. Along its southern front it overlooks a sector of the great Indo-Gangetic alluvial plain, here standing about 800 ft. above sea-level. Its mountain summits often rise well above 2,000 ft., and one of them only just fails to reach 5,000 ft. The following geological formations at one place or another enter into its constitution:—

Pliocene to Miocene. Siwaliks above Murees: a non-marine assemblage.

Middle and Lower Eocene. ? Punjab Saline Series ? above Lakis and Ranikots: frequently foraminiferal. (unconformity).

Cretaceous and Upper Jurassic. (unconformity).

Lower Jurassic, Trias, Permian and Upper Carboniferous. With Talchir glacial boulder bed at base. (unconformity).

Cambrian. Salt-pseudomorph beds above magnesian sandstones, Neobolus shales and purple sandstone; the last-named, being unfossiliferous, may perhaps be pre-Cambrian.

(contact variously interpreted).

Pre-Purple-Sandstone. ? Punjab Saline Series ?

The geological structure of the Salt Range is complicated, and before 1943 such few fossils as had been obtained from the Saline Series had all been challenged as introduced material. Accordingly, it was possible for a controversy to linger on from 1903 as to whether the Saline Series was of pre-purple-sandstone or of Eocene date, a truly amazing difference of opinion fraught with big consequences. Readers of the correspondence columns of *Nature* during 1944-45 will have

realized that the debate has recently flared up with renewed vigour¹. The cause has been an unexpected discovery by B. Sahni, whose reputation as a botanist is world-wide, that the Punjab Saline Series contains myriads of minute plant fragments of types that cannot be older than mid-Mesozoic. The matter has been handled admirably. A party of experienced field geologists has visited critical exposures, and a symposium has been called and its deliberations published². Unfortunately, the two antagonistic theses indicated above are still maintained, and an onlooker finds himself in the predicament so well expressed in "The Beggar's Opera": "How happy could I be with either, if t'other dear charmer were away".

In a short review it is only possible to outline the new evidence, which Sahni accepts at face value, and the nature of the response which it has evoked from E. R. Gee and others, who are not prepared to follow suit. References inserted below are to pages in the published account of the symposium.

Sahni claims that his fossil specks, of which the largest shown in his present illustrations only measures 0.5 mm., should be ascribed to wind-borne dust contemporaneous with the precipitation of the salt (p. xxv). He has established a wide distribution in specimens collected in mines sixty miles apart along the outcrop (p. ii). One location illustrated in a photograph is recorded as 730 ft. below surface and 4,150 ft. from mine entrance (p. xxxii). The organic matter is found in material which seems to have retained unblurred bedding. It is present alike in rock salt, saline earth, dolomite and oil shale (p. ii and photos p. xxxii). It occurs not only in specimens collected by Sahni and his party, but also in others collected by Gee and his party (p. xii). It shows in thin sections as well as on surfaces exposed by solution (p. ii). Great care has been taken to avoid laboratory contamination (p. viii). At the symposium a 'block' of dolomite "under .5 cc. in size was cut out of the interior of the hand specimen and polished on all faces to show that it was not cracked or pitted. It was rotated in a gas flame before it was placed in a jar of dilute HCl. After a few minutes a small piece of carbonized wood was released from the rock" (p. 313). All except one of the specimens which Sahni has investigated have yielded organic micro-debris, chiefly wood of angiosperms or gymnosperms, grass-like cuticles, pollen grains and chitinous parts of insects (p. viii). Attempts to discredit this material on the ground that it is more recent than Eocene are rebutted with the answer that there is nothing in its nature to show that it is later than mid-Mesozoic (pp. xxvii, 325); the fresh condition of much of it may be ascribed to embalment (p. x).

A. Lahiri, in a contribution (p. 329) which serves as a valuable supplement to Sahni's own, publishes photomicrographs of organic micro-debris still embedded in oil shales, some of the latter universally admitted to belong to the Punjab Saline Series. He says of certain samples that "the organic matter is found to be mostly in the form of minute fragments (perfectly laminated with the plane of bedding of the rock) of carbonized wood, fossil microspores, cuticles and pollen grains, only recognizable under high power and after the separation of the mineral matter".

E. R. Gee has given a wide and altogether admirable account of the subject, furnished with geological maps which can usefully be treated as illustrations of the report as a whole. Gee some years ago accepted the Eocene dating of the Punjab Saline Series, but by 1939 he had seen reason to change his mind