earliest economic objective in most tropical dependencies before maturity brought a broadening of their commerce and industry. There can be little doubt that, as in Northern Australia, for example, geographical discouragement to the search for easily accessible resources of high value and light weight has been a main cause of present economic stagnation. Moreover, in the case of each Caribbean Colony now under review, the likely return on capital outlay in forestry and agricultural development would certainly be slow, even if the development areas were—as, in fact, they are not—accessible to both labour and markets. Given the complete absence of satisfactory communications in the hinterlands of both Colonies, the prospects for commercial agriculture are necessarily gloomy, as the Commissioners, from abundant evidence, tactfully insist. In their modest words: "If little else results from our recommendations, at least the construction of certain good roads into the interior would lift the veil and open up the substantial possibilities which, we believe, exist. Until that is done we fear that periodic inquiries into the industrial and other possibilities of the interior of the Colony will be impossible to follow up and consequently unfruitful." Although this was written with reference to British Guiana, it applies equally to British Honduras. From their own meagre resources alone, the two Colonies cannot support the expense and organisation of the preliminary surveys and preparation of the ground necessary for the new agricultural and industrial enterprises proposed. In consequence, the report is urgent on the need for the establishment of a local development corporation in each Colony. These bodies would necessarily be dependent upon the Colonial Development Corporation for the greater part of their finances, as well as for the services of expert consultants.

In view of the inadequacy of the data on geographical conditions within each Colony, it was inevitable that the Commission would prepare a statement on the characteristics of the little-known regions which make up the hinterland and form the greater part of each Colony. These geographical statements, which have been woven into the fabric of the report, have been prepared by officers well acquainted with the regional concept of geography; each Colony is analysed on a basis, not of administrative subdivisions which have little relation to geographical realities, but of approximations to 'natural regions'. Indeed, so successful is the geographical presentation that the report should become an essential text for the study of the West Indies.

Space is not available for reference to more than a small proportion of the recommendations intended for the rehabilitation of agriculture which the report presents. Only one further observation is possible here; it refers to sugar production, the industry which easily dominates the life of British Guiana. The expansion expected is indicated by a proposed annual target of 230,000 tons of sugar, or about one-third more than the output for 1946. Although it would seem reasonable to hope, by reason both of the extent of the land capable of sugar-cane cultivation and of the long tradition—from the middle of the seventeenth century onwards—of successful sugar cultivation based largely on East Indian immigrant labour, that the sugar industry would offer a big contribution to the further settlement of British Guiana, the report briefly negatives the probability of this. The reasons do not, however, seem to be adequately explained in the report; and comparison between

British Guiana and, for example, Natal, the natural endowment of which for sugar-growing is certainly no greater, makes the official view difficult to accept. The comparison here is with a much smaller territory than British Guiana and one, moreover, with strictly limited possibilities of increased sugar acreage; yet its already planted sugar area is five times greater than that of British Guiana and is much more effectively settled by East Indian immigrants of the type that the Caribbean authorities might well consider further as suitable for settlement.

WALTER FITZGERALD

¹ West India Royal Commission Report. (Cmd. 6607.) (London: H.M. Stationery Office, 1945.) See *Nature*, 157, 254 (1946).

² Colonial Office, Report of the British Guiana and British Honduras Settlement Commission. (Cmd. 7533.) Pp. viii+360. (London: H.M. Stationery Office, 1948.) 7s. 6d. net.

ROTATION AND TERRESTRIAL MAGNETISM

IN 1923, H. A. Wilson put forward a conjecture whereby an electrically neutral, gravitating mass was supposed to be endowed with some of the properties of a charge; the phenomena of electromagnetism and gravitation, hitherto surprisingly independent, were to be related. The suggested relation was that the motion of the gravitational field of a mass should result in magnetism, just like the motion of the electric field of a charge. If the gravitational field of a mass is as strong as the electric field of a charge, then the one should be as potent in producing magnetism as the other. (Equality of strength of the fields can be defined by means of two equal masses and two equal charges.) For a rotating object this theory would predict proportionality between angular momentum and magnetic moment.

Wilson himself pointed out that this conjecture was not tenable. The rectilinear motion of a mass did not result in any measurable magnetic field, and any such effect would conflict with the well-tried principles of special relativity. It was just a surprising fact that the magnetic moment which this theory would predict for the earth was close to the measured value.

More recently, when evidence was forthcoming regarding the magnetism of some stars and of the sun, Blackett revived great interest in such considerations by suggesting that the same relation would fit remarkably well also in the case of these objects.

At a Geophysical Discussion of the Royal Astronomical Society held on February 25, the present state of such theories and their experimental and observational justification were under discussion. Prof. L. Rosenfeld was in the chair, and opened the discussion with a brief statement regarding the present state of the subject.

Dr. S. Runcorn presented much observational material. He pointed out that recent measurements of stellar magnetic fields, carried out by Babcock in the United States, are in good agreement with the suggested formula, if one accepts the statistical evidence for the rotation speed of such stars. The sun's field is now regarded as variable, but this does not necessarily point to a variable mechanism for the origin of a primary field.

The history of the earth's field has recently been investigated by the examination of ferromagnetic

clay deposited up to one million years ago, and it appears that in that period no major change has occurred. The mean direction of the clay magnetization is the geographical north, and the mean inclination is the one which a central dipole field would have at the latitude of the deposits.

Experiments on the change of the terrestrial field with depth in mines were then discussed by Dr. Runcorn. Such measurements should enable a distinction to be made between fields originating from the core of the earth and those which are of distributed origin, such as the effect suggested by Wilson and Blackett. Such experiments have revealed an unexpected preponderance of local anomalies, even in regions which were thought to be magnetically undisturbed. It appears that such anomalies may be several times as great as the effect sought. Further measurements are in progress, and the degree of disturbance of districts was estimated independently.

Dr. Runcorn also mentioned that a measurable distinction between 'core' and 'distributed' theories of geomagnetism should arise in the neighbourhood of a deep valley or gorge.

Mr. F. Hoyle explained that he had looked for specific processes which would result in stellar magnetism. Indeed, he had found a mechanism which may be expected to generate fields of the observed orders of magnitude. If a cloud of ionized gas is subjected to bombardment by a stream of particles, then the manner in which the momentum of the incoming stream will come to be shared by positive and negative ions is such as to imply a net current. Such a process must occur in the atmosphere of a star which is acquiring material from without, which does not originally possess the angular motion of the star.

On the question of terrestrial magnetism, Mr. Hoyle remarked that there are a number of possibilities of specific processes. The long decay-time of currents in the earth left open the possibility of magnetization in the past by processes which do not occur with similar intensity now. Streams of ionized gas are known to reach us from places in the vicinity of sunspots, and such streams will transport a magnetic field. It would be legitimate, though perhaps somewhat strained, to suppose that much more violent solar outbursts in the past were responsible for the transport of magnetism to the earth. Another point of view which he had recently come to consider, together with Mr. T. Gold, was that the objections which had been raised against Gilbert's original theory were inadequate, and that permanent magnetism may, in fact, cause the earth's field.

The present writer's contribution was concerned with the theoretical aspects of a fundamental law connecting rotation of neutral masses with magnetism. Blackett's empirical formula, in which the angular momentum is proportional to the magnetic moment, could not, in itself, be a statement of a physical law. The amount of angular momentum which is ascribed to a system depends entirely on the position and state of motion of the reference point chosen; but magnetic moment does not have a similar dependence on the co-ordinate frame. The suggestion is that the real law is such as to reduce to the formula under the conditions satisfied by the earth and the stars, when their centre is taken as the reference point for the computation of the angular momentum. The angular momentum in orbits cannot have any such magnetic effects, as can be seen from direct experiment, from considerations of special relativity, and

from the divergence of such an effect for rectilinear motion. If orbital motion is to have no such effect, then a self-gravitating disk would have to show no effect at that particular rotation speed which made the path of each constituent particle a free orbit.

There are four conditions which one might expect such a new, fundamental law to satisfy:

(1) *Differential Law.* It must be possible to consider separately the contribution to the total field which arises from any one mass element. The definition of any such contribution must require no more than a specification (however detailed) of physical conditions which would be measurable at that element.

(2) *Special Relativity.* The law must not give preference to one particular inertial frame over all others.

(3) *Equality of Inert and Gravitating Mass.* There must be no distinction between the effects on a mass of a gravitational field or of an acceleration.

(4) *Magnitude and Symmetry.* The law should result in a field of the right order of magnitude and symmetry for the earth; but it must not result in effects which could not have been overlooked in the laboratory.

The proposal of a 'fundamental law' implies that only dynamical quantities should be relevant to the magnetic contribution of each mass element. Conditions like temperature or conductivity are to play no part, if any significance is to be attached to the correlation between terrestrial and stellar fields.

It appears to be impossible to meet this set of conditions. Any combination of dynamical quantities which affect an element of mass partaking in the earth's motion can be made incomparably greater in the laboratory; any law which can be constructed from those quantities must always imply an obvious laboratory effect, if it is to account for the earth's field.

The Blackett formula avoids this difficulty. There a mass-element partaking in the earth's motion should have a magnetic effect proportional to the square of its distance from the earth's axis; but no such favourable factor would enter in the case of a spinning object in the laboratory. But the distance from the earth's axis is not a physical condition of a mass-element, as it is not a locally measurable quantity; and for this reason the formula offends against condition (1).

It is concluded that we lack any suggestion of a physical quantity to be related to magnetism in a novel way; the alternative view would be that one of the conditions could be dropped, but it is unlikely that such a view would come to be held.

Prof. P. M. S. Blackett described the reasons which had caused Wilson to reject the original hypothesis; but Wilson had not been aware of the kind of evidence which Dr. Runcorn had presented, and which appeared to be very compelling. His own view was that a solution of the difficulties could be found in a re-interpretation of the quantities appearing in the formula. The radius appearing in the formula for the angular momentum requires revision, and it might be interpreted as the radius of curvature of the modified gravitational equipotentials at each contributing mass-element. This definition should be acceptable, as such a radius of curvature is a quantity which can, in principle, be detected everywhere. Two plumb-lines, a certain distance apart, will point in

clay deposited up to one million years ago, and it appears that in that period no major change has occurred. The mean direction of the clay magnetization is the geographical north, and the mean inclination is the one which a central dipole field would have at the latitude of the deposits.

Experiments on the change of the terrestrial field with depth in mines were then discussed by Dr. Runcorn. Such measurements should enable a distinction to be made between fields originating from the core of the earth and those which are of distributed origin, such as the effect suggested by Wilson and Blackett. Such experiments have revealed an unexpected preponderance of local anomalies, even in regions which were thought to be magnetically undisturbed. It appears that such anomalies may be several times as great as the effect sought. Further measurements are in progress, and the degree of disturbance of districts was estimated independently.

Dr. Runcorn also mentioned that a measurable distinction between 'core' and 'distributed' theories of geomagnetism should arise in the neighbourhood of a deep valley or gorge.

Mr. F. Hoyle explained that he had looked for specific processes which would result in stellar magnetism. Indeed, he had found a mechanism which may be expected to generate fields of the observed orders of magnitude. If a cloud of ionized gas is subjected to bombardment by a stream of particles, then the manner in which the momentum of the incoming stream will come to be shared by positive and negative ions is such as to imply a net current. Such a process must occur in the atmosphere of a star which is acquiring material from without, which does not originally possess the angular motion of the star.

On the question of terrestrial magnetism, Mr. Hoyle remarked that there are a number of possibilities of specific processes. The long decay-time of currents in the earth left open the possibility of magnetization in the past by processes which do not occur with similar intensity now. Streams of ionized gas are known to reach us from places in the vicinity of sunspots, and such streams will transport a magnetic field. It would be legitimate, though perhaps somewhat strained, to suppose that much more violent solar outbursts in the past were responsible for the transport of magnetism to the earth. Another point of view which he had recently come to consider, together with Mr. T. Gold, was that the objections which had been raised against Gilbert's original theory were inadequate, and that permanent magnetism may, in fact, cause the earth's field.

The present writer's contribution was concerned with the theoretical aspects of a fundamental law connecting rotation of neutral masses with magnetism. Blackett's empirical formula, in which the angular momentum is proportional to the magnetic moment, could not, in itself, be a statement of a physical law. The amount of angular momentum which is ascribed to a system depends entirely on the position and state of motion of the reference point chosen; but magnetic moment does not have a similar dependence on the co-ordinate frame. The suggestion is that the real law is such as to reduce to the formula under the conditions satisfied by the earth and the stars, when their centre is taken as the reference point for the computation of the angular momentum. The angular momentum in orbits cannot have any such magnetic effects, as can be seen from direct experiment, from considerations of special relativity, and

from the divergence of such an effect for rectilinear motion. If orbital motion is to have no such effect, then a self-gravitating disk would have to show no effect at that particular rotation speed which made the path of each constituent particle a free orbit.

There are four conditions which one might expect such a new, fundamental law to satisfy:

(1) *Differential Law*. It must be possible to consider separately the contribution to the total field which arises from any one mass element. The definition of any such contribution must require no more than a specification (however detailed) of physical conditions which would be measurable at that element.

(2) *Special Relativity*. The law must not give preference to one particular inertial frame over all others.

(3) *Equality of Inert and Gravitating Mass*. There must be no distinction between the effects on a mass of a gravitational field or of an acceleration.

(4) *Magnitude and Symmetry*. The law should result in a field of the right order of magnitude and symmetry for the earth; but it must not result in effects which could not have been overlooked in the laboratory.

The proposal of a 'fundamental law' implies that only dynamical quantities should be relevant to the magnetic contribution of each mass element. Conditions like temperature or conductivity are to play no part, if any significance is to be attached to the correlation between terrestrial and stellar fields.

It appears to be impossible to meet this set of conditions. Any combination of dynamical quantities which affect an element of mass partaking in the earth's motion can be made incomparably greater in the laboratory; any law which can be constructed from those quantities must always imply an obvious laboratory effect, if it is to account for the earth's field.

The Blackett formula avoids this difficulty. There a mass-element partaking in the earth's motion should have a magnetic effect proportional to the square of its distance from the earth's axis; but no such favourable factor would enter in the case of a spinning object in the laboratory. But the distance from the earth's axis is not a physical condition of a mass-element, as it is not a locally measurable quantity; and for this reason the formula offends against condition (1).

It is concluded that we lack any suggestion of a physical quantity to be related to magnetism in a novel way; the alternative view would be that one of the conditions could be dropped, but it is unlikely that such a view would come to be held.

Prof. P. M. S. Blackett described the reasons which had caused Wilson to reject the original hypothesis; but Wilson had not been aware of the kind of evidence which Dr. Runcorn had presented, and which appeared to be very compelling. His own view was that a solution of the difficulties could be found in a re-interpretation of the quantities appearing in the formula. The radius appearing in the formula for the angular momentum requires revision, and it might be interpreted as the radius of curvature of the modified gravitational equipotentials at each contributing mass-element. This definition should be acceptable, as such a radius of curvature is a quantity which can, in principle, be detected everywhere. Two plumb-lines, a certain distance apart, will point in

directions converging at the centre of the earth. Their separation and the angle between them are hence a measure of their distance from the centre.

The magnetic contribution of one mass-element can then be taken as being proportional to the product of its mass, angular velocity, and the square of this distance which can be defined there, as the plumb-line example shows. All the quantities entering now are of local significance as required. On the basis of such an interpretation no difficulty would arise in the case of an orbital motion. Plumb-lines at neighbouring points on a planet would converge on the centre of the planet, and hence only the spin would count.

Prof. Blackett then mentioned a suggestion for a laboratory experiment, which would be more direct than the measurements of the earth's field in mines or across canyons. A small massive object at rest in the laboratory should make a contribution to the earth's field, which would result in a local effect of the order of 10^{-8} gauss. Though small, this would be measurable.

The general discussion which followed touched upon many topics, but only those dealing with the theoretical approach suggested by Prof. Blackett will be presented here; although, indeed, many other stimulating points of view were advanced.

The writer said he considers the radius of curvature of the geopotentials an unsuitable length to enter into the Blackett formula. It is true that a local determination of this length is possible in principle; but locally it may easily give values which are arbitrarily large. Two plumb-lines placed closely in between two small gravitating objects may well hang parallel. The proposed interpretation would then imply an arbitrarily large magnetic contribution of matter in such localities. A slowly rotating, horizontal wheel would be another example of a locality where plumb-lines may all be parallel. The use of a geometrical property of a field, without regard to its strength, for determining the strength of an effect, must lead to the anomaly that an arbitrarily small change of circumstances may have an arbitrarily large effect.

The suggestion of a new law of magnetism is hence still quite vague. No actual physical quantity has been suggested which might be related, by way of a new law, to the magnetism of large rotating bodies. In these circumstances, even the experimental approach is severely handicapped, for it is much harder to search for a new relation than to prove an asserted law.

T. GOLD

LEAF FORM AND FUNCTION*

By SIR EDWARD SALISBURY, C.B.E., F.R.S.

THE green leaf of the flowering plant is perhaps the most important structure in the world, since upon it in large measure depends the maintenance of life upon the earth's surface. While the law of entropy applies to all our surroundings, the green leaf is a part of Nature that perhaps more than any other is doing a little winding up in a cosmos that is mostly running down.

The chemical processes which are carried on in the leaf are complex and involve a chain of reactions

and enzyme activities the details of which are still in many respects obscure; but as we are concerned more with the factory itself than the details of the processes that go on within it, we can take for granted the elaborate series of chemical changes accelerated by enzyme action. For though these may well demand an organisation of the living protoplasm of great importance, the tools for exploring the sub-microscopic structure of the cell are only now being developed.

The older botanists fully recognized that the spiral arrangements which leaves very commonly exhibit in their attachment to the stem tend to ensure the maximum display of the leaf surfaces with the minimum of overlap, and that the changes in growth of leaf stalks which accompany any changes in the direction of incident light bring about the so-called leaf mosaics which minimize the competition for light between the leaves of the same plant or branch. What they did not appreciate was the complexity of the phenomenon and of the interaction between light and growth. For not only is the direction of the leaf stalk and orientation of the blade involved but also the amount and effect of the growth-promoting substances, produced by the apical buds and by the leaves themselves. These growth hormones stimulate cell enlargement in varying degrees, according to the intensity of the light (maximum leaf area usually being attained in partial shade) and also inhibit in more or less degree the development of axillary buds.

The interrelations of growth effect that fine adjustment that brings about the display of the photosynthetic pigments to the best advantage. The various types of spacing in a spiral, such that there are always three leaves in any single circle of the stem, suggest a growing point that behaves as the multicellular equivalent of a three-sided apical cell found in many pteridophyte and gymnosperm embryos. This primary tendency is, however, modified by the competition for food and by space relations imposed by the combined effects of the rate of production of leaf rudiments on one hand and the rate of stem elongation on the other.

In general, intense light inhibits the effect of growth-promoting substances upon stem elongations so that, in bright light, the internodes are short and the foliage crowded, whereas in dull light the internodes lengthen and the leaves are widely spaced. Since bright light causes decomposition of the pigments, these reactions are obviously beneficial; but for leaf blades the optimum light intensity for permitting the growth-promoting substances to operate at a maximum is apparently higher than for stems, though this is not true of monocotyledonous leaves which are comparable to leaf stalks in their behaviour. In woodland plants there is often a marked increase in the size of the leaf blades with the diminution of the light consequent upon the canopy development of the foliage above in spring.

The span of life of leaves is usually short. Deciduous trees, such as most of the native species of Britain, bear leaves that only function for about six months, though their loss and annual renewal permit a large leaf surface with evasion of an adverse season. Evergreen leaves such as those of the holly and ivy may live for more than two years, while the needles of coniferous trees often persist for 2-12 years, according to the species. In general, it would appear that the length of life-span of leaves is related to their degree of activity. The giant leaves of *Victoria regia* with their rapid metabolism have a short life; they only function

* Substance of a Friday Evening Discourse entitled "Leaves" delivered at the Royal Institution on February 4.