Tatensturm", quoted by Carlyle in Chapter 8 of "Sartor Resartus"—indicate an anticipation of the law of the conservation of energy! Magnus suggests that Goethe's botanical work influenced Jussieu, de Candolle and Robert Brown, a suggestion which I must leave to those more learned than myself in such matters, and, rather half-heartedly, that the foundation of the study of comparative anatomy owes much to Goethe. In some such way it has been suggested that Francis Bacon had a great influence on the work of the early Fellows of the Royal Society, although Bacon's views of the way to set about scientific work were anything but those adopted. It is true, as I have pointed out, that the great physiologists who dealt with colour vision in Goethe's time owed something to him. On the whole, however, it seems to be accepted that, if Goethe had never published anything in science, things would have developed just as they, in fact, did. If Goethe is to be compared to any great figure in the history of science, it must not be to any great modern but rather to Aristotle, whose powers of observation and generalization led him to important advances in the biological sciences while his physics, dominated by philosophical speculation, was not significant. Goethe was a subjective natural philosopher rather than a man of science in the modern sense.

Must we, then, regret that the great man spent so much of his time in fields where his genius produced relatively little of worth? Regrets are, in any case, fruitless, but perhaps they are not even called for. The spectacle of this comprehensive genius attacking such varied problems with such a zest, attempting with such spirit and such confidence to bring a new order into vast fields of knowledge, is an inspiring one. It is well that there shall be seekers after a general light: a stirring up—dust, light and movement—may not be progress, but nevertheless it is occasionally not a bad thing. Be that as it may, in Goethe's activities we see a last attempt—a glorious attempt—to take all culture and all knowledge for one man's province, and we salute it.

RADIOACTIVE TRACERS IN CHEMISTRY

THE scientific programme of this year's anniversary meeting of the Chemical Society was arranged in collaboration with the Chemistry Division of the Government's Atomic Energy Research Establishment. During March 28–30 a symposium on "The Chemistry of the Heavy Elements" and on "Radioactive Tracers" took place in Oxford, and on April 1 a discussion was held in London on "The Use of Radioactive Tracers in Chemistry". The Oxford meeting has already been reported in *Nature* of June 18 and 25.

The London discussions were divided in two parts, the morning session being devoted mainly to tracer applications in inorganic and physical chemistry, the afternoon session to organic-chemical and biological problems. After opening remarks by the president of the Chemical Society, Sir Ian Heilbron, Prof. F. A. Paneth took the chair and introduced the subject by a few general remarks on the steadily growing importance of tracer methods in chemistry since the discovery of artificial radioelements and their production on a large scale. The significance of this new

research tool for biological investigations was emphasized by Prof. G. v. Hevesy when he took the chair at the afternoon session.

The question foremost in the minds of many who intend to make use of this technique is, naturally, where and how radioactive tracers can be obtained. It was answered by H. Seligman in his report on "The Availability of Radio-Isotopes". He was able to assure the audience that the Atomic Energy Research Establishment at Harwell is now in a position to provide all the tracers of long life which a year or so ago could only be bought from the United States, and those of shorter life as well, which were previously not obtainable by purchase. A novel feature of the Harwell organisation is a kind of information bureau which tries to help those with little experience in the subject by giving advice as to the tracers likely to be most satisfactory in particular research problems and industrial uses. Equally comforting was the news about the prices, which are kept very low—a point which may be sometimes decisive. It is, therefore, unlikely that any research laboratories within reasonable distance from Harwell will ever be hampered in the use of radioactive tracers by lack of the necessary radioactive material; they might, however, for some while find it more difficult to obtain the equipment of special rooms and high-class modern apparatus, and the staff trained in the new technique.

The other papers dealt with a variety of researches which had only the one feature in common that in the course of the investigation use had been made of a radioactive tracer; for this very reason they do not lend themselves easily to a summarizing review, all the more so as several of the subjects discussed were of a highly specialized nature. All the papers presented at the meeting will be printed in full in the *Journal of the Chemical Society*, and those interested in related problems will find in the complete texts a wealth of helpful information; but in this brief report we can scarcely mention more than the subjects of the contributions and give an indication of the main results.

Since the early days of indicator methods, research workers have found them most useful in the study of diffusion problems. K. E. Zimen, M. Hillert and G. Johansson, in Göteborg, have now employed silver-111 and mercury-203 for a determination of the self-diffusion coefficients of silver and mercury; their paper was read by E. Berne. A second paper from Göteborg, by R. Lindner, reported on work done in Prof. Hedvall's laboratory on the self-diffusion in silver sulphate, lead and zinc oxides, $\text{ZnO} \cdot \text{Fe}_2\text{O}_3$ and $\text{CaO} \cdot \text{SiO}_2$. That such a study of diffusion coefficients by radioactive tracers can be used as a new means for the investigation of condensation processes in solutions was shown by J. S. Anderson and K. Saddington, of the Atomic Energy Research Establishment, Harwell. From the diffusion coefficients of radioactive sodium tungstate, containing tungsten-185 and -187, it was concluded that with decreasing pH there is a progressive aggregation of the tungstate ions; above $\text{pH} = 9$, the normal tungstate ion exists, whereas over the range $\text{pH} = 2.3$ – 5 a condensed ion is predominant.

Prof. J. W. Irvine, jun., reported on researches carried out by him at the Massachusetts Institute of Technology in collaboration with Prof. C. D. Coryell and W. B. Lewis on the "Electron-transfer (Exchange) between Cobaltous- and Cobaltic-amino Complexes". Cobalt-60 was used as tracer in the cobaltous com-

plex; at suitable time intervals the complex was destroyed by acid, the cobaltous ion extracted quantitatively and its specific activity determined from its gamma-radiation; the specific activity decreases, naturally, if part of the active cobalt enters the cobaltic complex. Complexes containing either ammonia or ethylenediamine were studied, and exchange-rate constants determined.

While in the above papers radioactive tracers were employed in order to study exchange reactions, M. Haissinsky, from the Institut de Radium, Paris, gave an example of their usefulness in examining the behaviour of chemical elements in extreme dilution. Together with A. Coche, he reported new "Experiments on the Cathodic Deposition of Radioelements". Their results indicate that, contrary to some previous statements, the Nernst electrochemical equation is valid for the bismuth isotope thorium C down to 10^{-13} N, and for the lead isotope thorium B to 10^{-10} N, if the cathode consists of gold, silver, copper or nickel; they found, however, some anomalies with other electrodes. In the case of polonium, these were attributed to the penetration of polonium atoms into the metal of the cathode.

Before organic-chemical and biological problems were discussed in the afternoon session, J. F. Duncan described studies carried out by himself, G. B. Cook and A. W. Kenny at Harwell on "Techniques in Radiochemical Assay". He issued a warning against possible errors in beta counting, such as those due to multiple counts, variation of high tension or temperature, self-absorption, back-scattering, etc.; fortunately, most of these can be avoided without great difficulty if for the purpose of the experiment only knowledge of the relative values of activities is necessary, and this is the case in the majority of biological applications.

Prof. E. D. Hughes, of University College, London, spoke on "Radioactive Halogens and the Mechanism of Substitution Reactions". He showed how the study of such old problems of classical organic chemistry as the mechanism of replacement, racemization and Walden inversion can be helped by the radioactive technique; in the cases considered, it could be shown that bond-formation and bond-fission occurred synchronously.

Biochemical problems were discussed by the last four speakers. In their "Studies with Radioactive Penicillin", S. Rowlands, D. Rowley and E. Lester Smith, of the University of Edinburgh, prepared penicillin labelled with sulphur-35 by growing the mould on a medium containing activated sulphate. The penicillin obtained was sufficiently active to make it possible to follow up its fate in experimental animals. It was found that, if injected intramuscularly, 100 per cent of it was excreted with the urine. On the other hand, if bacteria were killed with the activated penicillin, no active sulphur could be detected in them, which proved that less than ten molecules of penicillin had been taken up by a bacterium.

F. D. S. Butement, of the Atomic Energy Research Establishment, Harwell, showed in his paper on "Radioactive Tracers in Antisera" how the study of problems of immunology can be refined. He succeeded in introducing iodine-131 into the globulin fraction of antiserum without destroying its specific agglutinating properties; the amount of iodine activity carried on *Proteus vulgaris* when agglutinated by the specific antiserum could be used to measure the agglutinin.

From the University of Saskatchewan came a paper by Prof. J. W. T. Spinks and G. Dion on the "Study of Fertilizer Uptake using Radio-Phosphorus"; it was read by R. Spence. The authors made use of phosphorus-32 to measure the uptake of fertilizer phosphorus and soil phosphorus by wheat plants; as fertilizers they employed, in field and in greenhouse experiments, ammonium, sodium- and calcium-phosphates. Their results showed, among other facts, that in the early stages of growth practically all the phosphorus taken up comes from the fertilizer, while during the later stages the phosphorus uptake from this source is relatively small.

In his report on "Some Chemical Problems in the Use as a Fumigant of Methyl Bromide Labelled with ^{82}Br ", F. P. W. Winteringham, from the Pest Infestation Laboratory, Slough, demonstrated a new way in which the fate of the fumigant can be followed up in treated foodstuffs and in infesting insects. It concluded a well-attended and very successful discussion meeting. F. A. PANETH

PROGRESS IN HIGH-SPEED CALCULATING MACHINE DESIGN

A CONFERENCE on high-speed automatic digital calculating machines was held in the University Mathematical Laboratory at Cambridge during June 22-25 to mark the completion of the EDSAC (electronic delay storage automatic calculator).

The EDSAC is a serial binary machine based on the ideas put forward in an unpublished report circulated in 1945 by von Neumann on behalf of the group working at the Moore School of Electrical Engineering in Philadelphia, where the ENIAC—the first electronic calculating machine ever built—was at that time under construction. The EDSAC is believed to be the first complete machine of its kind to be put into operation, although a slightly smaller machine, without automatic input and printing devices, has been built at the University of Manchester. This machine was described to the conference by Prof. F. C. Williams and Dr. T. Kilburn. Several machines are nearly completed in the United States. Information about these was given in a paper communicated by Dr. H. D. Huskey (National Bureau of Standards).

In the EDSAC, ultrasonic delay units are used for storage, the total capacity for orders and numbers (each with sixteen binary digits—equivalent to five decimals) being 1,024. There are facilities for combining two adjacent storage locations so as to hold a number with the equivalent of ten decimal digits. The programme of orders and any numerical data required are fed into the machine from a punched tape, and the results of the calculation are printed automatically on a teleprinter. The EDSAC was demonstrated to members of the conference by the writer and Mr. W. Renwick.

The EDSAC and the machines like it are a great advance on the ENIAC in that they are very much smaller and yet have greater storage capacity for numbers. Most people will, however, agree that electronic calculating machines are still too large—the EDSAC has more than three thousand valves—and that one of the primary aims of future research must be to reduce their size. In the present writer's