giving a resolution within 3 Å., using currents as low as 10^{-10} amp. It was suggested that the technique has application, for example, to the investigations of fragmentation processes on metal catalysts.

The third session dealt with the investigation of surface phenomena using the field-emission microscope. Further discussion on adsorption and on the mobility of oxygen on tungsten was given in a paper by R. Gomer and J. K. Hulm (Chicago). Oxygen in the monolayer is not mobile below 450° K., and becomes a two-dimensional gas only at temperatures exceeding 700° K. The layers formed at low temperatures do not form at room temperatures. No changes in the chemisorbed layer are produced by heating in high vacuum to temperatures up to 600° K. Studies of the reaction of oxygen were described by E. A. Gulbransen (Westinghouse). Transmission optical methods have application in studies of chemical and physical reactions on metals as well as of the study of fine details of the substructure of the metal itself. A. T. Gwathmey (Virginia) described work on the nature of a metal surface as determined with the aid of large single crystals in the form of a surface exposing all possible faces, and also in a form of a flat surface exposing only specific faces. Study of surface processes such as etching, wetting, electrodeposition, and friction and wear, confirm the dependence of surface properties on crystal face. Oxidation of a copper crystal was shown on a coloured ciné film. P. W. Kruse (Notre Dame and Farnsworth) and E. A. Coomes (Notre Dame) described a study of the kinetics of strontium oxide on molybdenum. The pattern obtained on a heat-treated molybdenum point corresponds to that proposed for clean tungsten by Becker and by Muller.

The conference closed with a series of papers from Linfield College, Oregon. The stability of electron emission from metals in the combined presence of high surface electric field and elevated cathode temperature (T - F emission) was discussed by W. P. Dyke; and experiments using temperatures of 2,000° K. for tungsten were described, evaporation and surface migration being used to obtain a smooth clean surface. Stability of the field-emitter was discussed by E. E. Martin; instability can be due to the action of helium atoms, which diffuse through the tube envelope; the effect is removed by using an evacuated buffer space. Emission of current densities $\sim 5 \times 10^6$ amp./cm.² from a single lattice step of clean tungsten was observed by J. K. Trolan using a pulse technique. W. W. Dolan described the dissolution of individual lattice planes made visible by the technique of high surface electric field and high cathode temperature, thus permitting a new approach to the study of transport mechanism at the emitter surface.

F. LLEWELLYN JONES

CHROMATOGRAPHY

AN anonymous American Fulbright scholar, in a generous appreciation of his visit to the biochemistry department of an English university, wrote as follows: "It is certainly not entirely a misfortune that English laboratories to-day are much less lavishly equipped than those in the United States. For example, one of the most important advances of recent years has been the use of the method known as paper chromatography. To operate this method, one needs only a strip of moistened

filter paper, suitably mounted in a box, and a few substances that can be bought from any dealer in pure chemicals. Yet by this technique complicated mixtures of numerous different amino acids or other substances can be sharply separated into their components, and then identified, even when separation by the methods of classical chemistry was impossibly tedious and ineffective. The method has come rapidly into use all over the world, for countless different kinds of problems. It is almost certainly no accident that this method was developed in England in the midst of all its shortages, toward the end of the war, rather than in the far more opulent United States" (Report on Fulbright Program, 1951-52, p. 23). However this may be (and one would not like to have the argument accepted too readily by those responsible for financing the universities), it is fair to say that British workers have been pioneers in the modern development of chromatography in its various aspects, and it is appropriate that the *British Medical Bulletin*, which is sponsored by the British Council, should publish a survey of this subject (Vol. 10, No. 3, pp. 161-252. London: British Council. 15s. net).

Dr. C. E. Dent acted as scientific editor, and there are articles by nineteen authors who are all experts in their respective fields. Dr. A. J. P. Martin, who, jointly with Dr. R. L. M. Synge, received the Nobel Prize in 1952, was chairman of the planning committee and he writes the foreword. He outlines the immensely wide scope of chromatography, which has become an indispensable tool in almost all fields of biochemistry, and he ends by trailing his coat with the words, "I hope that chromatography may ultimately play some part in the process, which I believe to be wholly desirable, of the conversion of medicine from an art to a science". This is followed by a section on general principles of chromatography by Dr. R. J. P. Williams. For the benefit of newcomers to the subject, this might perhaps have been made a little more 'general' even at the risk of some overlapping with other articles. A large proportion of the limited space of this introductory section is devoted to gradient analysis, which is a rather specialized type of technique. There follows a brilliant chapter by Drs. A. T. James and A. J. P. Martin on gas-liquid chromatography, a technique which can be used to separate and estimate the members of a homologous series of substances such as hydrocarbons, alcohols, fatty acids, etc. This technique has a very wide scope and it is likely that in time it will completely replace analytical distillation. Some indication is given of the devices of these authors for estimating and automatically recording the substances after separation, devices which are elegant and ingenious in the extreme.

There are many references to adsorption chromatography and to ion exchange resins; but the main part of the volume is devoted to paper chromatography, starting with Dr. R. Consden's introduction, which is full of useful practical tips based on his wide experience as a pioneer in this technique. There seems to be a slip in the statement (p. 182) "if the R_F is too high in one solvent, a lower [sic] homologue may prove satisfactory". Other sections include the chromatography of inorganic substances, organic acids, proteins, peptides, amino-acids (dealt with in three sections), porphyrins, sterols, antibiotics and vitamins, carbohydrates and phosphoric esters. This last section, by Dr. F. A. Isherwood, is outstandingly good, as is Dr. R. Markham's review of nucleotide

work, with a mass of useful information compressed into a small space. Dr. J. Gross deals with the thyroid hormones and includes an account of the recent discoveries which he and Mrs. R. Pitt-Rivers have made on the various iodine compounds in the thyroid gland and in the blood. Much of this important work would have been impossible without the use of paper chromatography in combination with tracer technique using radioactive iodine, which permitted analyses to be made on an ultra-micro scale. This article, and the one by Drs. C. E. Dent and J. M. Walshe on amino-acid metabolism, are of direct medical appeal.

Some of the articles contain a good deal of hitherto unpublished work, and each includes a bibliography which greatly adds to the value of the publication. Incidentally, references to *Nature*, which is being regularly used as a forum for the recording and exchange of new chromatographic procedures, appear frequently in these bibliographies. It seems a pity that all the illustrations in a volume entitled "Chromatography" should be produced in black-and-white. Some private journals, notably *Endeavour*, have shown what aesthetically pleasing results can be achieved by the use of colour in illustrating chromatographic techniques, even if, from a purely scientific point of view, the gain does not justify the large extra cost. No worker who is interested in chromatography, whether beginner or expert, can fail to benefit from reading this admirable publication. It is likely to be used on the laboratory bench as well as in the library, and it deserves to be protected from accidental 'paper chromatography' by being given a permanent binding. D. C. HARRISON

THE EXPANDING UNIVERSE

A SLIGHTLY modified version of a talk given by G. J. Whitrow in February 1953 in the B.B.C. Third Programme has now been published as a paper under the title of "The Orthodox Theory of the Expanding Universe" in the October issue of *Occasional Notes of the Royal Astronomical Society* (17, 3; 1954). A fairly large portion of the talk is concerned with a work by Paul Coudere, a French astronomer, which appeared in a French edition in 1950 and was translated into English by J. B. Sidgwick under the title "The Expansion of the Universe" (1952). The first part of the talk gives a very useful résumé of the developments in our knowledge of the universe during the past thirty years, in which period it was accepted that the universe extends far beyond the limits of our galaxy, and also that the red shift strongly suggested the recession of the extra-galactic nebulae, all of which were moving away from one another. The talk then considers other interpretations that have been given to the displacement towards the red end of the spectrum, dealing in particular with various matters connected with Coudere's book.

This book is recognized as a very useful contribution to cosmology in so far as it provides us with a simplified account of the mathematical theory of the expanding universe; but in recent years some British cosmologists have advocated a new theory based on the hypothesis of the continuous creation of matter. Differences of view arise regarding the form in which this new matter emerges. C. Pascual Jordan advocates the formation of super-giant stars;

but other cosmologists who challenge this view have put forward the conception of new matter appearing in the form of hydrogen molecules or even neutrons, and the density of the universe remains constant, the decrease in density which would result from the recession of the galaxies being exactly balanced by the continuous creation of matter. As might be expected, these theories have been subjected to severe criticisms; nevertheless, they are able to claim one strong argument in their favour—that they allow more time for stellar evolution than the older orthodox view which led to a conception of the earth being older than the galaxies. This argument is no longer valid since Baade's announcement about the wrong 'calibration' of the distances of the Cepheid variables (the investigations of Thackeray and Wesselink on the distance of the Magellanic Clouds have corroborated Baade's conclusions); now it seems that it must have been about four thousand million years ago since the universe started expanding, and in addition, the distances of the extra-galactic nebulae must now be taken as twice those previously estimated.

Whitrow concludes with an expression of disagreement with Coudere's acceptance of the two fundamental hypotheses at the basis of the 'orthodox' theory of the expanding universe; these are that the red shifts are due to the Doppler effect associated with recessional motion, and also that the laws of Einstein's general theory of relativity, either with or without the idea of cosmical repulsion, govern the structure of the universe as a whole. He thinks that neither of these is on such firm foundations as Coudere believes; but in spite of this disagreement he speaks highly of the latter's book which, on p. 213, seems to express a little doubt on the inadequacy of some of his own arguments. "Everything possible has been extracted from existing observations; until the appearance of new data we would be wise to give our imaginations a rest." To this remark Whitrow makes the very relevant reply: "... if such advice had been accepted by Copernicus he would never have ventured beyond the Ptolemaic theory".

RECENT MOLLUSCA OF CANADA

THE great extent of Canada has resulted, and will probably long result, in its molluscan fauna being studied in four separate areas: the Eastern Provinces, British Columbia, the Arctic region based on Hudson Bay, and the Prairie Provinces. In the first three, marine, freshwater and land forms are involved, but in the fourth only freshwater and land. Partly for this reason and also because many of the species were first described from other countries, the literature involved is extensive and scattered.

A very useful service in rendering this literature and its results available for the whole of Canada has been performed by the publication of Bulletin No. 129 of the National Museum of Canada, entitled "Catalogue of the Recent Mollusca of Canada", by Dr. Aurèle La Rocque (Biological Series No. 44. Pp. ix+406. Ottawa: Queen's Printer, 1953; 2.50 dollars). It fills a very obvious gap, for it provides a list of the accepted species and forms that have so far been recorded, numbering about 2,250. The author has wisely included forms that have actually