

Table 3 Comparison of innominate measurements normalised to acetabulum height

Ratio to acetabulum height of:	Libben mean	σ	Distance from Libben mean in s.d.			
			STS 14	SK 3155	SK 50	OH 28
Ilium height	2.33	0.14	+2.8	> +3.4*		
Greater sciatic to anterior notch	1.37	0.08	-2.3	-0.05	+0.46	-1.52
Anterior inferior to ischial spine	1.98	0.09	-1.3	-2.2		-2.9
Anterior inferior to posterior superior spine	2.57	0.14	-2.4	1.7		-0.4
Functional length of ischium	1.45	0.08	-3.8	-4.9	-0.8	-4.5

These measurements were taken by me. The functional length of the ischium⁴ is measured on both the Australopithecines and the Libben sample; the breaks on the STS 14 ischium and on the SK 3155 ischium minimise the measurement. The Libben sample size is 30. That some of these measures which cross, or almost cross the acetabulum differ considerably from the Libben mean when normalised by acetabulum size further emphasises the differences in proportion; when normalised to the acetabulum the iliac blades are relatively large and most other measures small, but normalised to the ilia everything else is extremely small.

*Addition of the unfused iliac crest would make this larger.

relative head size is, if anything, above the human mean⁵. These head, neck and shaft length relations are shown by two femora attributed to "*Homo*," and one attributed to "*Australopithecus*." Moreover, judging from the published photographs they also characterise the complete Afar hominid femur attributed to "*Australopithecus*"⁶. Former claims of relatively small head size come from comparison of head diameter with shaft diameter in incomplete specimens. Analysis of femora of known or reconstructable lengths shows that the shaft diameters were relatively large.

The demonstration that the heads were not relatively small calls to question the claim that the acetabula were relatively small¹, since of all pelvic dimensions the acetabulum has the highest correlation with femur length (0.840 for 30 Libben Amerinds). Schultz⁷ demonstrated the large relative size of the STS 14 and reconstructed SK 50 acetabula. When various comparable innominate measurements are normalised to acetabulum height (Table 3), and compared with the Libben sample, only relative ilium height is consistently above the Libben mean, as expected given the greater amount of iliac flare in the early hominids⁴. Iliac breadths are generally below the Libben mean, except for one measure of SK 3155, and contrary to "conventional knowledge", the functional length of the ischium is relatively short. The *H. erectus* innominate follows the australopithecine pattern. Only if normalised to iliac height would the early hominid acetabulum appear relatively small, and the other measurements would be relatively yet further below the Libben mean. Given the above, it is far more likely that australopithecine acetabula are human-like relative to body size, and the ilia are relatively long.

In sum, the distinguishing features of the early hominid hip complex reveal a pattern of form differing from living humans because of a combination of narrower birth canal and

markedly greater muscular activity, but not differing in locomotor capacity.

M. H. WOLPOFF

Department of Anthropology,
University of Michigan,
Ann Arbor, Michigan 48104

- 1 McHenry, H. M., and Corruccini, R. S., *Nature*, 259, 657-658 (1976).
- 2 McHenry, H. M., *Am. J. phys. Anthropol.*, 40, 329-340 (1974).
- 3 Burns, P. E., *Nature*, 232, 350 (1971).
- 4 Lovejoy, C. O., Heiple, K. G., and Burstein, A. H., *Am. J. phys. Anthropol.*, 38, 757-780 (1973).
- 5 Lovejoy, C. O., and Heiple, K. G., *Am. J. phys. Anthropol.*, 32, 33-40 (1970).
- 6 Walker, A. C., *J. human Evol.*, 2, 545-555 (1973).
- 7 Schultz, A. H., *Folia Primat.*, 11, 181-199 (1969).
- 8 Johanson, D. C., and Taieb, M., *Nature*, 260, 293-297 (1976).

McHENRY AND CORRUCINI REPLY—The discussion here¹ is about different fossils. We show that taken together the relative proportions of the proximal ends of KNM-ER 1481 and 1472 are distinguishable from those of SK 82 and 97 and KNM-ER 1504. Wolpoff argues¹ that the reconstructed femoral head diameter and neck length relative to reconstructed femoral length in Sts 14 are similar to KNM-ER 1481. We feel that a meaningful and unambiguous argument should not be based upon measurements of a fossil (Sts 14 femur) described as "a useless jumble of glued fragments surmounted by a crude plaster head and neck"³. Nor should the length of a half missing ischium (SK 3155)⁴⁻⁶ be used to bolster such an argument.

A few points need to be clarified. (1) Our analysis is based on both univariate and canonical variates analysis and the results agree. (2) The variation on the fourth canonical variate is not interpreted by us as the result of femoral head size only. In fact, greater trochanter projection has the highest correlation with that variate, followed by neck length and femoral head size. The canonical variate is complex, reflecting the complexity of interrelationships among the variables. (3) Because variance scales to the number of subjects, one would not expect a high percentage of

variance on the fourth canonical variate. The fourth variate contributes substantially to the multivariate distances between subjects. (4) KNM-ER 1481 does indeed have a relatively big head, which is just our point because it represents the *Homo* taxon. The ratio of femoral head size to length in KNM-ER 1472 is 0.10 (ref. 7), which is exactly the Libben mean. This specimen is also classified as *Homo*. The only specimen classified as *Australopithecus* to which Wolpoff refers in regards femoral head size and length is Sts 14, but this specimen is missing both its head and its distal end⁴. (5) Walker⁶ did not conclude that the "relative head size . . . is above the human mean". He showed that his composite and reconstructed australopithecine femur has a ratio of head size to length slightly below the human mean.

In our view canonical variates analysis is one more useful tool for analysing complex biological shape. The method allows one to take into account variability within taxa, it reduces the complexity of interrelated shapes and sizes to understandable dimensions, and it allows an anatomical region to be treated as a total complex (inasmuch as the available fossil material will allow).

HENRY M. McHENRY

University of California, Davis

ROBERT S. CORRUCINI

Smithsonian Institution

- 1 Wolpoff, M. H., *Nature*, 264, 812-813 (1976).
- 2 McHenry, H. M., and Corruccini, R. S., *Nature*, 259, 657-658 (1976).
- 3 Dav, M. H., *Symp. zool. Soc. Lond.*, 33, 29-51 (1973).
- 4 McHenry, H. M., thesis, Harvard Univ. (1972).
- 5 McHenry, H. M., *Am. J. phys. Anthropol.*, 43, 245-262 (1975).
- 6 McHenry, H. M., *Am. J. phys. Anthropol.*, 43, 39-46 (1975).
- 7 Dav, M. H., Leakey, R. E. F., Walker, A. C., and Wood, B. A., *Am. J. phys. Anthropol.*, 42, 461-476 (1975).
- 8 Robinson, J. T., *Early Hominid Posture and Locomotion* (University of Chicago Press, Chicago, 1972).
- 9 Walker, A., *J. human Evol.*, 2, 545-555 (1973).