

## Sequence and topology of a model intracellular membrane protein, E1 glycoprotein, from a coronavirus

John Armstrong, Heiner Niemann\*, Sjeff Smeekens††, Peter Rottier† & Graham Warren

European Molecular Biology Laboratory, Postfach 10.2209, 6900 Heidelberg, FRG

\* Institut für Virologie, Fachbereich Humanmedizin der Justus-Liebig Universität, Giessen, FRG

† Institute of Virology, Veterinary Faculty, State University of Utrecht, 3509 TD Utrecht, The Netherlands

In the eukaryotic cell, both secreted and plasma membrane proteins are synthesized at the endoplasmic reticulum, then transported, via the Golgi complex, to the cell surface<sup>1-4</sup>. Each of the compartments of this transport pathway carries out particular metabolic functions<sup>5-8</sup>, and therefore presumably contains a distinct complement of membrane proteins. Thus, mechanisms must exist for localizing such proteins to their respective destinations. However, a major obstacle to the study of such mechanisms is that the isolation and detailed analysis of such internal membrane proteins pose formidable technical problems. We have therefore used the E1 glycoprotein from coronavirus MHV-A59 as a viral model for this class of protein. Here we present the primary structure of the protein, determined by analysis of cDNA clones prepared from viral mRNA. In combination with a previous study of its assembly into the endoplasmic reticulum membrane<sup>9</sup>, the sequence reveals several unusual features of the protein which may be related to its intracellular localization.

The coronaviruses are a diverse class of enveloped RNA viruses of considerable medical and agricultural significance; they also provide a model for the study of persistent viral infections (see ref. 10 for review). In contrast to many enveloped viruses, the coronavirus mouse hepatitis virus (MHV) A59 buds inside the cell, into the lumen of the endoplasmic reticulum<sup>11-14</sup>. The assembled virion then appears to travel, via the Golgi complex, to the cell surface. Of the two viral membrane proteins, the smaller one, E1, is necessary for formation of the envelope, and is restricted to internal cell membranes; apparently it only reaches the cell surface as part of the budded virion<sup>12,13</sup>. Thus, the E1 glycoprotein is potentially a convenient model for studying those features of a membrane protein that determine its arrest at a particular destination on the membrane transport pathway.

The mRNAs of MHV-A59 form a 'nested set': the seven RNAs share the 3' region of the positive-stranded genome, but extend to different lengths towards the 5' end<sup>15-18</sup>. From each RNA, only the 5' gene is translated<sup>19,20</sup>. In addition, a non-coding 'leader' sequence of approximately 70 bases, from the 5' end of the genome, is common to the mRNAs<sup>18,21,22</sup>. The E1 gene is second from the 3' end and is therefore translated from the second smallest mRNA, RNA 6 (refs 19, 20). The sequence of the 3'-terminal gene, encoding the viral nucleocapsid protein, has been determined previously<sup>23,24</sup>.

Copy DNA clones spanning the E1 gene were prepared by two methods<sup>23-25</sup> and sequenced in the vectors M13mp8 (ref. 26) or pEMBL8 (ref. 27) by the chain-termination method<sup>28</sup> (data available on request). A sequence of 780 nucleotides (Fig. 1), containing a single long open reading frame, precedes the coding region for the viral nucleocapsid protein. A leader of 76 nucleotides, almost identical to the leader of the smallest mRNA, RNA 7 (ref. 22), lies in front of the first potential initiator codon. Thus, the sequence in Fig. 1 represents the 5' end of RNA 6, encoding the E1 protein and starting at or near the extreme 5'-terminal nucleotide.

```

CCTATAAGAGTGTGGCGTCCGTACGTACCCCTCAACTCTAAACTCTGTAGTTTAA
  10          30          50
MetSerSerThrThrGlnAlaProGluProValTyrGlnTrpTh
ATCTAATCAAACATTATGAGTAGTACTACTAGCCGCCAGGCGCTATCAATGGAC
  70          90          110
rAlaAspGluAlaValGlnPheLeuLysGluTrpAsnPheSerLeuGlyIleIleLeuLe
GGCCAGCAGGCGAGTCAATTCCTTAAGGAATGGAAGTCTCGTTGGGCATTACTACT
 130          150          170
UpheIleThrIleIleLeuGlnPheGlyTyrThrSerArgSerMetPheIleTyrValVa
CTTTATTACTATCATACTACAGTTCGGTTACAGGCGGTAGCATGTTTATTATTGTTG
 190          210          230
lLysMetIleIleLeuTrpLeuMetTrpProLeuThrIleValLeuCysIlePheAsnC
GAAATGATAAATCTTGGTTAATGTGGCTACTGATTTTGTGTTTTCATTTCAATG
 250          270          290
sValTyrAlaLeuAsnAsnValTyrLeuGlyPheSerIleValPheThrIleValSerII
CGTGATCGGCTAAATAATGTGTATCTGGATTTTCTATAGTGTATTACTATAGTCCAT
 310          330          350
eValIleTrpIleMetCtyrPheValAsnSerIleArgLeuPheIleArgThrGlySerTr
TGTAATCTGGATTATGATTTTGTAAATAGCATAAGGTTGTTTACAGGACTGGTAGCTG
 370          390          410
pTrpSerPheAsnProGluThrAsnAsnLeuMetCysIleAspMetLysGlyThrValTy
GTGGAGCTCAACCCGAAACAAACACCTATGTATAGATAGAAAGTACCGTGTA
 430          450          470
rValArgProIleIleGluAspTyrHisThrLeuThrAlaThrIleIleArgGlyHisLe
TGTAGACCCATTATTGAGGATTACCATACACTAACAGCCACTATTATTCGTGGCCACCT
 490          510          530
uTyrMetGlnGlyValLysLeuGlyThrGlyPheSerLeuSerAspLeuProAlaTyrVa
CTACATGCAAGGTGTTAAGCTAGGACCGCTTCTCTTGTCTGACTGCGCCGCTTATGT
 550          570          590
lThrValAlaLysValSerHisLeuCysThrTyrLysArgAlaPheLeuAspLysValAs
TACAGTTGCTAAGGTGCACACCTTGCCTTAAGCCGCGCATTTCTAGACAAGGTAGA
 610          630          650
pGlyValSerGlyPheAlaValTyrValLysSerLysValGlyAsnTyrArgLeuProSe
CGGTGTTAGCGGTTTTGCTGTTTATGTGAAGTCCAAGGTCGGAATTCAGCACTGCCCTC
 670          690          710
rAsnLysProSerGlyAlaAspThrAlaLeuLeuArgIle*
AAACAACCGAGTGGCGGACACCCGATTTGTGAGAATCTAATCAAACTTAAGGATG
 730          750          770

```

Fig. 1 Sequence of the E1 cDNA and protein extending to the initiator codon of the adjacent nucleocapsid gene<sup>23</sup>. Proposed membrane-spanning regions are overlined.

Two versions were found, in two different clones, for the sequence immediately upstream from the E1 initiator codon. The shorter one is shown in Fig. 1; in the second clone, an additional copy of the pentanucleotide ATCTA was found between nucleotides 65 and 66, making the sequence similar to that of the region adjacent to the nucleocapsid gene of another strain of MHV<sup>29</sup>. This difference could represent a mutation; alternatively, it may reflect heterogeneity in the normal mRNA population. Indirect support for the latter possibility comes from the observation that a RNase-T<sub>1</sub> oligonucleotide from this region of RNA 6, corresponding to the shorter sequence, was recovered in markedly lower yield than those from the rest of the molecule<sup>30</sup>. This site represents the point of fusion between the 5' leader sequence and the coding portion of the RNA. The fusion is thought to occur by 'jumping' of the viral RNA polymerase to particular sites on its genome-length, negative-stranded template; the resumption of transcription then produces each of the subgenomic mRNAs<sup>22,31,32</sup>. Thus, it seems possible that the polymerase may jump to more than one point on the template for each mRNA, generating variable numbers of the repeated pentanucleotide AUCUA in the resulting transcript.

Figure 1 shows the amino acid sequence encoded by the E1 gene. The predicted molecular weight of the protein is 26,000, slightly higher than that observed by gel electrophoresis<sup>19,33</sup> but consistent with the unusual electrophoretic behaviour of this<sup>33</sup>, and other, hydrophobic proteins. Several features of the protein, when assembled into membranes in the virus<sup>33</sup>, or *in vitro*<sup>9</sup>, are reflected in the sequence. First, in contrast to the majority of membrane proteins, E1 is known to lack a cleaved 'signal peptide'<sup>9</sup>: the N-terminal region of the sequence contains no good candidate for a cleavage site<sup>34</sup>. Second, the N-terminal region bears O-linked sugars<sup>35,36</sup>, which, uniquely among viral proteins so far studied, are the only known post-translational modification to E1. Assuming that the terminal Met is removed<sup>37</sup>, the N-terminal sequence is Ser-Ser-Thr-Thr, which is identical to the O-glycosylated amino terminus of M-type glycoprotein A (ref. 38). The O-linked sugars of E1 are them-

† Present address: Molecular Cell Biology Group, State University of Utrecht, 3509 CH Utrecht, The Netherlands.

