

HIGHLIGHTS



MODEL SYSTEMS

The dawning of the Age of Pisces

HIGHLIGHT ADVISORS

ALLAN BASBAUM

UNIVERSITY OF CALIFORNIA
SAN FRANCISCO, CA, USA

RANDY BUCKNER

WASHINGTON UNIVERSITY,
MO, USA

DAVID CLAPHAM

HARVARD MEDICAL SCHOOL,
MA, USA

PIETRO DE CAMILLI

YALE UNIVERSITY SCHOOL OF
MEDICINE, CT, USA

BARRY EVERITT

UNIVERSITY OF CAMBRIDGE,
UK

GORDON FISHELL

SKIRBALL INSTITUTE, NY, USA

MARY KENNEDY

CALIFORNIA INSTITUTE OF
TECHNOLOGY, CA, USA

LYNN NADEL

UNIVERSITY OF ARIZONA,
AZ, USA

DENNIS O'LEARY

THE SALK INSTITUTE FOR
BIOLOGICAL STUDIES, CA, USA

TERRY SEJNOWSKI

THE SALK INSTITUTE FOR
BIOLOGICAL STUDIES, CA, USA

WOLF SINGER

MAX-PLANCK-INSTITUT FÜR
HIRNFORSCHUNG, GERMANY

CLAUDIO STERN

COLUMBIA UNIVERSITY, NY,
USA

RICHARD W. TSJEN

STANFORD UNIVERSITY
SCHOOL OF MEDICINE, CA, USA

RAFAEL YUSTE

COLUMBIA UNIVERSITY, NY,
USA


It may not be a system as popular as *Drosophila* or the mouse for neuro-genetic studies, but the zebrafish is quickly becoming a prime tool in our quest for the relationship between genes and brain function. Several screens have already identified genes essential for the development of this vertebrate, and the promise that the whole sequence of the zebrafish genome will be completed in about two years guarantees that this creature will be the subject of an even more extensive genetic analysis. In preparation for the bonanza to come, two recent papers explore two experimental angles — one genetic and one psychophysical — which illustrate the wide spectrum of opportunities that the study of the zebrafish can offer us.

First, Nasevicius and Ekker tested the possibility of knocking down zebrafish genes by injecting modified oligonucleotides known as morpholinos. Morpholinos act by blocking the initiation of translation, which reduces the non-specific effects observed with conventional antisense oligonucleotides. The authors injected diverse morpholinos into the yolk of embryos as young as the one-cell stage and found that the oligonucleotides reduced the expression of targeted proteins effectively. More importantly, morpholino injections phenocopied previously described mutations in some of the target genes and, in other cases, led to the appearance of new phenotypes, which could help to determine the role of genes of unknown

function. Although this technology still has some limitations, the use of morpholino oligonucleotides can become a faster, efficient alternative to the production of knockout animals and it could allow the use of classic reverse-genetic approaches in the zebrafish.

Most of the genetic screens done in the zebrafish so far have focused on mutations that affect embryonic development. However, the behavioural repertoire of this species could also be exploited if we develop the appropriate tools. By studying the optomotor behaviour of larval zebrafish, Orger *et al.* have provided us with a good lead to tackle the problem. When a grating is moved under normal fish, they will swim in the direction of the perceived motion (see movies of the stimuli online). The authors asked what features of movement perceived by the fish are responsible of eliciting the swimming behaviour. They found that both first-order, luminance-modulated features and second-order, contrast-modulated motion were equally capable of triggering optomotor responses, a surprising finding considering that only mammals have been thought to detect second-order motion.

In addition to challenging the idea that cortical processing is required for the detection of second-order motion, the basic characterization of the optomotor reflex provided by this work should increase the usefulness of this task for the behavioural screening of mutant fish. The Orger *et al.* study,



Images of mutant zebrafish taken, with permission, from *Nature Genet.* **26**, 216–220 (2000) © Macmillan Magazines Ltd.

together with the findings of Nasevicius and Ekker would seem to indicate the delayed arrival of the Age of Aquarius.

Juan Carlos López

References and links

ORIGINAL RESEARCH PAPER Nasevicius, A. & Ekker, S. C. Effective targeted gene 'knockdown' in zebrafish. *Nature Genet.* **26**, 216–220 (2000) | Orger, M. B. *et al.* Perception of Fourier and non-Fourier motion by larval zebrafish. *Nature Neurosci.* **3**, 1128–1133 (2000)

FURTHER READING Summerton, J. Morpholino antisense oligomers: the case for an RNase H-independent structural type. *Biochim. Biophys. Acta* **1489**, 141–158 (1999) | Haffter, P. *et al.* The identification of genes with unique and essential functions in the development of the zebrafish, *Danio rerio*. *Development* **123**, 1–36 (1996) | Driever, W. *et al.* A genetic screen for mutations affecting embryogenesis in zebrafish. *Development* **123**, 37–46 (1996)

ENCYCLOPEDIA OF LIFE SCIENCES Transgenic zebrafish production | Zebrafish embryo as a developmental system
WEB SITE Morphant database