sequence has a G3:C70 pair (Fig. 3b), it is probably a tRNA<sup>Ile</sup>.

Although it follows most of the rules for tRNA structure, this structure has one unusual feature (Fig. 3b). A very highly conserved purine-pyrimidine tertiary base pair, between position 15 and the last base of the extra arm, is a pyrimidine-purine pair in this sequence, a feature shared by only a few other tRNAs10.

All three introns identified so far in eubacteria are in tRNA genes. This may be due to constraints of our detection method (which favours small, abundantly transcribed introns that selfsplice efficiently in vitro, for example), or introns in tRNA may have been selectively retained during evolution. Although unrelated to group 1, introns are frequently present in archaebacterial and eukaryotic tRNAs, within or next to the anticodon. Deletion of the intron results in failure to modify an anticodon uridine to pseudouridine in eukaryotic tRNA<sup>Tyr</sup> (refs 18-20). It is possible that introns also have a role in post-transcriptional modification of bacterial tRNAs.

Are group I introns ancient? If introns were present in the common ancestor of all cells, some should have been retained in the eubacterial lineage despite genome 'streamlining'2. Closely related introns have now been found in the anticodon loops of tRNA genes from widely divergent bacterial phyla. But as they are in different tRNAs, we cannot determine when these introns were acquired. One intriguing possiblity is suggested by the recent demonstration that the aminoacyl tRNA synthases for Leu, Ile and (unexpectedly) Arg are related<sup>21</sup>, which raises the interesting corollary that their respective tRNAs may also have had a common ancestor. We must therefore consider the possibility that tRNAs were interrupted by introns before completion of the genetic code. Alternatively, the introns may have arrived recently at their present locations, for example by reversal of splicing<sup>22</sup> followed by reverse transcription.

Southern hybridization with the Azoarcus intron probe (Fig. 4) suggests that related introns are widespread among Proteobacteria. As more bacterial introns are sequenced, a comparison of the sequences of introns in homologous genes (for example, for the same tRNA) will provide conclusive evidence concerning the antiquity of group I introns. 

- Darnell, J. E. & Doolittle, W. F. Proc. natn. Acad. Sci. U.S.A. 83, 1271-1275 (1986).
- Palmer, J. D. & Logsdon, J. M. Curr. Opin. Genet. Dev. 1, 470-477 (1991). 3
- Kuhsel, M. G., Strickland, R. & Palmer, J. D. Science 250, 1570-1573 (1990) 4
- 5. Xu, M.-Q., Kathe, S. D., Goodrich-Blair, H., Nierzwicki-Bauer, S. A. & Shub, D. A. Science 250, 1566-1570 (1990).
- Yang, D., Oyaizu, Y., Oyaizu, H., Olsen, G. J. & Woese, C. R. Proc. natn. Acad. Sci. U.S.A. 82, 6. 4443-4447 (1985)
- Stackebrandt, E., Murray, R. G. E. & Trüper, H. G. *Int. J. syst. Bact.* **38**, 321–325 (1988). Cech, T. R., Zaug, A. J. & Grabowski, P. J. *Cell* **27**, 487–496 (1981).
- 8
- Winans, S. C., Kerstetter, R. A. & Nester, E. W. J. Bact. 170, 4047-4054 (1988).
- 10. Kruger, K. et al. Cell **31**, 147–157 (1982). 11. Cech, T. R. Gene **73**, 259–271 (1988).
- 12. Michel, F. & Westhof, E. J. molec. Biol. 216, 585-610 (1990).
- Antao, V. P., Lai, S. Y. & Tinoco, I. Jr Nucleic Acids Res. 19, 5901–5905 (1991).
  Muramatsu, T. et al. Nature 336, 179–181 (1988).
- Sprinzl, M., Dank, N., Nock, S. & Schön, A. Nucleic Acids Res. 19, (suppl.) 2127-2171 (1991). 15
- Schulman, L. H. & Pelka, H. Science 242, 765–768 (1988).
  Lee, C. P., Seong, B. L. & RajBhandary, U. L. J. biol. Chem. 266, 18012–18017 (1991).
- Johnson, P. F. & Abelson, J. Nature 302, 681-687 (1983). 19. van Tol, H. & Beier, H. Nucleic Acids Res. 16, 1951-1966 (1988).
- 20. Choffat, Y., Suter, B., Behra, R. & Kubli, E. Molec. cell. Biol. 8, 3332-3337 (1988).
- Nagel, G. M. & Doolittle, R. F. Proc. natn. Acad. Sci. U.S.A. 88, 8121–8125 (1991).
  Woodson, S. A. & Cech, T. R. Cell 57, 335–345 (1989).
- 23. Woese, C. R. Microbiol. Rev. 51, 221-271 (1987).
- Young, J. P. W. in Biological Nitrogen Fixation (eds Stacey, G., Burris, R. H. & Evans, H. J.) 43-86 24. (Chapman & Hall, New York, 1992).
- 25. Gott, J. M., Shub, D. A. & Belfort, M. Cell 47, 81-87 (1986).

ACKNOWLEDGEMENTS. We thank F. Ausubel, D. Arp and E. Nester for bacterial strains; M. Gillis and M. Shavegani for DNA; E. Shub, S. Sousa and M. Xu for help with experiments, H. Goodrich-Blair for critically reading the manuscript, and J. P. W. Young for Fig. 1c. This work was supported by a research grant from the N.I.H. (to D.A.S.) and by a grant from the Deutsche Forschungsgemeinschaft (to B.R.-H.).

### CORRECTIONS

# **Evolution of ecological** differences in the Old World leaf warblers

### Adam D. Richman & Trevor Price

Nature 355, 817-821 (1992)

IN Fig. 2a of the above letter the second and third rows were inadvertently interchanged. In addition, the sequence at position 15176 (ref. 1) for both Sylvia and P. coronatus should read 'A' and not 'T'. 

1. Anderson, S. et al. Nature 290, 457-465 (1981).

## Structure and expression of a human oxytocin receptor

### Tadashi Kimura, Osamu Tanizawa, Kensaku Mori, Michael J. Brownstein & Hiroto Okayama

Nature 356, 526-529 (1992)

WE have discovered an error in the DNA sequence of our pOTR gene in Fig. 1b. The nucleotides 714-717, CGGC should be GGCGGCG. The amino acids underneath should be Ala Ala Ala instead of Ala Gly. Accordingly, the complementary DNA encodes a 389 amino-acid protein with a relative molecular mass of 42,819. This error does not affect the conclusions of the  $\square$ paper.

## GUIDE TO AUTHORS

follow these guidelines so that your manuscript may be handled expeditiously

Pit a so follow these guidelines so that your manuscript may be handled expeditionsly. Mantles is an intermational journal covering all the sciences. Contri-butors simulate therefore bear in mill those readers who work in other fields and those for whom lengths is a science language, and write clearly and simple, avoiding unnecessary technical terminology. Space in the journal is finited, making compatibility of *Nature*, without interruptions of the jest, concarins about 1/393 works. Manuscripts are selected for penitochappe of *Nature*, without interruptions of the jest, concarins about 1/393 works. Manuscripts are selected for publication according or deducid Manuscripts are selected for publication according to endersid the safety of the selection of the pathetic strate according to the selec-ter provide the selection of the selection of the selection of the Manuscripts and the selection of the selection of the selection the selection of the selection of the selection of the selection of the selection manuscripts are been to the adverted on receipt but (event that and fare sent for review). These that are not treiveness, limited requires for the evolution of specific reviewers are usually hexeded. Manuscripts are usually sent to too or three regregation location. Manuscripts are usually sent to too or three regregation location. Manuscripts are usually sent to too or three regregation location. Manuscripts are usually sent to too. The regregation and location. Manuscripts are usually sent to too or three regregation and location. Manuscripts are usually sent to too or three regregation and location. Manuscripts are usually sent to too or three regregation and location. Manuscripts are usually sent to too or three regregation and location. Manuscripts are usually sent to too or three regregation and location. Manuscripts are usually sent to too or three regregation and location. Manuscripts are usually sent to too or three regregation and location. Manuscripts are usually sent to be and the sent tor both t

Inclum: Monisciple inserpret in provide the provided of the

data in the databases that exis for the purpose, and crystalistightly availability of these data. Once any second of the purpose, and to mention of these and an another the second of the purpose, and the mention receive galley proofs in about 4 weeks. *Nature's staff* will deft manuscripts with a werk to hereity and clearly, so constitutors should check there provide arefully. Manuscripts are generally published 2–3 weeks after receive a cerpinal order. *Nature* does not exact page charges. Contributors receive a reprint order form with their proofs: reprint orders are processed after the manuscript is published and payment received.

#### Categories of paper

<text><text><text><text><text><text>

News and Views editor Scientific Correspond

constants travecular: Scientific Gorespondence is for discussion of topical scientific matters, including those published in *Nature*, and for miscellaneous contributions. Priority is given to letters of rever than 500 words and 5 references.

#### Preparation of manuscripts

Proparation of manuscripts All manuscripts shull be typed, double-spaced, on one side of the paper only. An orginal and three copies are required, each accom-pointed by atrovic, II photographic are included. Ion sets of orginal-are required, for line drawings, one set of originals and three good-quality photocopies are acceptable. Reference ishs, figure legends and tables should all the in-separate sheets, all of which should be double-spiced and manipered. Reference isha such accept ory of a systemic annascript, and dearly marked as such end labeled swith their manuscript numbers. Tubes say which they are many sources of the set of a strate-terin long and in the paper is about which the minum or the charing terminology and in fewer than 80 characters in the case of Arrides and Letters. Acrive every, numerical vulses, abbreastions and parctu-tion are to be avoided. These should contain one or two key works for indexing parses.

tion are to be avoided. This should contain one or two key words for indexing purposes. Artwork should be marked individually and clearly, with the addition standard when knows, the manuscript manifer likelity, no figure should be larger than 20 kg. 20 em. Figures with several pairs are to be avoided and are permitted only if the parts are closely related, either experimentally or logically. Unlettened onlyinals of phos-graphs should be provided. Organizations is a technical when a maniference of the provided of the parts are to closely related. The provided of the provided of the parts are to closely related and not the single-kiter code for animo acids. Or collours with an *Nature* can accommidate 20 animo acids. Or coll histo pairs. Single of the or ever illustrations, with a single space between rows.

acc recover rows. Suggestions for cover illustrations, with captions and labelled with

spectra and the second and a second secon first and last page numbers are included, reference to books should include publisher, place and date. **Abbreviations**, symbols, units and greek letters should be identified the first time they are used. Acronyms should be avoided whenever possible and, if used, defined. Footnotes are not used except for changed addresses.

changed addresses. Acknowledgements are brief, grant and cuntribution numbers are not allowed. Submission. Manuscripts can be sent to the Editor at 4 Little Essex Siteret, London WC2R N.F. UK at 1137 National Press Bailding. Washington, DC 20045, USA. Manuscripts or proofs sont by air contert to London shulld be declared as 'manuscripts' and 'value S' to prevent the imposition of import daity and value-added tax (VAL).

Received 26 December 1991; accepted 17 March 1992.

<sup>1.</sup> Gilbert, W. Nature 271, 501 (1978).