

Ferromagnetism and EMFs

SIR — The question of whether weak, extremely low-frequency electromagnetic fields (EMFs) can cause cancer always generates heated debate (see, for example, refs 1–3). In addition to epidemiological studies, a substantial body of literature exists on EMF stimulation of cells grown *in vitro* (for example, refs 4, 5). Although numerous effects have been reported, many have been difficult to replicate (see refs 6, 7), and no clear biophysical mechanism has emerged. Many of the proposed mechanisms, like ion cyclotron resonance⁸, have drawn criticism for being physically unrealistic (see ref. 9).

From developments in a totally unrelated field, there may be a much simpler, as yet overlooked, mechanism for explaining many of these *in vitro* EMF cellular effects. For the past two decades, the study of the biologically precipitated ferrimagnetic mineral magnetite (Fe_3O_4) has relied heavily on the use of ultrasensitive superconducting quantum interference device (SQUID) magnetometers to quantify trace levels of magnetite in various biological and laboratory materials^{10,11}. It rapidly became clear that unique clean-laboratory techniques were required for this work because of the ubiquitous presence of ferromagnetic contamination. This contamination included ferromagnetic particulates present not only in the dust in the air, but also adsorbed onto the surfaces of laboratory equipment, present within glass and plastics, and even in reagent-grade laboratory chemicals and water.

We have encountered the same problem in our recent attempts to grow cells in tissue culture for an investigation of their magnetic properties. It is customary to use disposable, pre-sterilized plastic labware (flasks, pipettes, centrifuge tubes, and so on) and commercially prepared culture media in tissue-culture experiments because of their convenience and the assumption of a high level of quality control and cleanliness. We have found that none of these materials is free of ferromagnetic particulate contamination. Liquid-transfer manipulations, typical of cell-culture protocols, wash these particles from the surfaces of flasks and pipettes,

and concentrate them with the cells during centrifugation. As an example, in a sham experiment we used 50 ml of leukocyte culture medium to rinse ten plastic T-250 flasks, ten 10-ml pipettes and ten 50-ml centrifuge tubes. After final centrifugation, we detected the equivalent of 160 ng magnetite in the rinsate, and the magnetic data indicated that the contaminants are small particles, usually in the sub-100-nm size range. As 160 ng magnetite equates to about 32 million 100-nm³, this can be compared to the approximately 1 million cells that would have been produced in an equivalent culture volume.

Magnetite particles, 100 nm in diameter, either naked or coated with bovine serum albumin, are readily taken up by human white blood cells, including non-phagocytic lymphocytes as well as phagocytes¹². Because the ferromagnetic particles interact strongly with magnetic fields, their presence in cell cultures, at a number density far higher than that of the cells, may provide a simple mechanism to account for links between EMF exposure and *in vitro* biological effects. A simple calculation shows that the mechanical energy present in a single 100-nm magnetite crystal exposed to a 60-Hz, 0.1-mT magnetic field is many times the thermal background noise¹³. Such particles, if adsorbed on cell surfaces or ingested by the cells, could conceivably transfer this energy to contiguous cell structures such as mechanically activated ion channels (which operate with a gating force close to the thermal noise limit^{14,15}), and thereby alter cytoplasmic ion concentrations sufficiently to produce the observed biological effects.

We are not aware that the authors of any of the published studies of *in vitro* EMF effects have either controlled for, or attempted to reduce the levels of, ferromagnetic contamination. Although this is understandable, because the particles are difficult to detect and quantify except by sensitive magnetometry, their existence should not be ignored. *In vitro* studies may ultimately provide the information that will explain the connection between EMF exposure and biological effects, and as

such they constitute roughly half of the projects at present being sponsored by the 5-year, \$65 million NIEHS/DOE research programme on the biological effects of EMF. However, any effect of EMF exposure on cultured cells, if it is due to the presence of ferromagnetic contaminants, would have no relevance to *in vivo* biology. Data used to establish human exposure standards to electromagnetic fields must rely on properly controlled experiments.

Atsuko K. Kobayashi

Joseph L. Kirschvink

Division of Geological & Planetary Sciences,

California Institute of Technology,

Pasadena,

California 91125, USA

Michael H. Nesson

Department of Biochemistry & Biophysics,

Oregon State University,

Corvallis,

Oregon 97331, USA

Extant fauna of ancient carbon

SIR — We have found that organisms of the rich, freshwater biological community involving sponges, flatworms and other benthic species near a thermal vent at the bottom of Lake Baikal are built of ancient carbon lacking ¹⁴C.

The vent occurs at a depth of 420 m in Frolikha Bay (55°31' N, 109°46' E)^{1,2}. It was found earlier that the carbon of its benthic organisms was produced by methanogenic bacteria, as revealed by the very small values of $\delta^{13}\text{C}$ (–60 to –72‰)². However, the age of this carbon was not known.

Using a Tandemtron accelerator mass-spectrometer, we measured the contents of ¹⁴C in carbon of two *Bdellocephala* flatworms and a sponge collected on a bacterial mat of Frolikha vent, and found them to be equal to 0.43, 0.34 and 0.28 of that typical of modern organic matter, corresponding to apparent radiocarbon ages of 6,860 ± 260, 8,740 ± 80, and 10,200 ± 220 years before present (BP), respectively.

Hence, about 60–70 % of the carbon of the near-vent organisms has originated from ancient methane, rather than from modern atmospheric CO₂ due to photosynthesis or methanogenesis. The source of ancient carbon was also not limestone, as is sometimes the case in freshwater systems, since the uppermost layer of Baikal sediments is known to have an apparent radiocarbon age of less than 1,000 years BP at many locations (see ref. 3 and T. N. *et al.*, unpublished data).

Frolikha vent arises from meteoric water seeping through Baikal sediments⁴ that are known to contain high concentra-

1. Pool, R. *Nature* **349**, 554 (1991).

2. Florig, H.K. *Science* **257**, 468 (1992).

3. Hilemann, B. *Chem. Engng News* **71**, 15–29 (1993).

4. Goodman, R., Wei, L.-X., Bumann, J. & Shirley-Henderson, A. *Bioelectrochem. Bioenerget.* **29**, 185–192 (1992).

5. Blackman, C.F., Benane, S.G. & House, D.E. *Bioelectromagnetics* **12**, 173–182 (1991).

6. Lacy-Hulbert, A., Wilkins, R., Metcalfe, J.C. & Hesketh, R. *Bioelectromagnetics Soc. 16th a. meet.* (abstr.) 52 (Bioelectromagnetics Soc., Copenhagen, 1994).

7. Saffer, J.D. & Thurston, S.J. *Bioelectromagnetics Soc. 16th a. meet.* (abstr.) 54 (Bioelectromagnetics Soc., Copenhagen, 1994).

8. Liboff, A.R. *J. Biol. Phys.* **13**, 99–102 (1985).

9. Adair, R.K. *Phys. Rev.* **A34**, 1039–1048 (1991).

10. Walker, M.M., Kirschvink, J.L., Perry, A.S. & Dizon, A.E. in *Magnetite Biomineralization and Magnetoreception in Organisms: A New Biomagnetism* (eds Kirschvink, J.L., Jones D.S. & MacFadden, B.J.) 154–166 (Plenum, New York, 1985).

11. Kirschvink, J.L., Kobayashi-Kirschvink, A. & Woodford, B.J. *Proc. natn. Acad. Sci. U.S.A.* **89**, 7683–7687 (1992).

12. Matsunaga, T., Hashimoto, K., Nakamura, N., Nakamura, K. & Hashimoto, S. *Appl. Microbiol. Biotechnol.* **31**, 401–405 (1989).

13. Kirschvink, J.L. *Phys. Rev.* **A46**, 2178–2184 (1992).

14. Denk, W. & Webb, W.W. *Phys. Rev. Lett.* **63**, 207–210 (1989).

15. Howard, J. & Hudspeth, A.J. *Neuron* **1**, 189–199 (1988).