

*Biophysics Program, University of California, San Francisco, California 94143, USA

e-mail: cavoigt@picasso.ucsf.edu

†Center for Systems and Synthetic Biology and Institute for Cell and Molecular Biology, and ‡Department of Chemistry and Biochemistry, University of Texas, Austin, Texas 78712, USA

§Department of Synthetic Biology, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

||Department of Pharmaceutical Chemistry, University of California, San Francisco, California 94107, USA

1. Yeh, K.-C., Wu, S.-H., Murphy, J. T. & Lagarias, J. C. *Science* **277**, 1505–1508 (1997).
2. Schmitz, O., Katayama, M., Williams, S. B., Kondo, T. & Golden, S. S. *Science* **289**, 765–768 (2000).
3. Davis, S. J., Vener, A. V. & Vierstra, R. D. *Science* **286**, 2517–2520 (1999).
4. Utsumi, R. et al. *Science* **245**, 1246–1249 (1989).
5. Jin, T. & Inoué, M. *J. Mol. Biol.* **244**, 477–481 (1994).
6. Kwon, O., Georgellis, D. & Lin, E. C. C. *J. Biol. Chem.* **278**, 13192–13195 (2003).
7. Shimizu-Sato, S., Huq, E., Tepperman, J. M. & Quail, P. H. *Nature Biotechnol.* **20**, 1041–1044 (2002).
8. Gambetta, G. A. & Lagarias, J. C. *Proc. Natl Acad. Sci. USA* **98**, 10566–10571 (2001).

Supplementary information accompanies this communication on Nature's website.

Competing financial interests: declared none.
doi:10.1038/nature04405

INSECT COMMUNICATION

'No entry' signal in ant foraging

Forager ants lay attractive trail pheromones to guide nestmates to food^{1,2}, but the effectiveness of foraging networks might be improved if pheromones could also be used to repel foragers from unrewarding routes^{3,4}. Here we present empirical evidence for such a negative trail pheromone, deployed by Pharaoh's ants (*Monomorium pharaonis*) as a 'no entry' signal to mark an unrewarding foraging path. This finding constitutes another example of the sophisticated control mechanisms used in self-organized ant colonies.

To investigate whether foragers lay a negative signal on the unrewarding branch of a trail bifurcation, we removed paper substrate from immediately after the fork on the unrewarding branch (the other branch led to a sucrose feeder) after it had been used by a trail-laying colony of ants. This paper substrate was transferred to the entrance of one branch of a similar set-up, in which both branches had previously led to sucrose and had been used by a second colony of ants. The other branch of the second set-up received a neutral control paper substrate (for details, see supplementary information). Foragers walking from the nest could choose either of the test branches or make a U-turn.

We found that 69% continued to walk away from the nest and make a branch choice. Of these, most (71%) chose the branch with the control substrate ($\chi^2=22.1$, d.f.=1, $n=137$, $P<0.001$); the remainder U-turned towards the nest on reaching the trail bifurcation. U-turns were more than four times as likely if the ant had contacted the unrewarding-branch substrate (55%) as opposed to the neutral-control substrate (13%) ($\chi^2=40.9$, d.f.=1, $n=200$, $P<0.0001$). Neither substrate came from a previously rewarding trail, so this result cannot be attributed to differences in positive-trail pheromone concentrations.

We next investigated the negative signal's location by taking substrate from five locations on a bifurcating trail that had one rewarding

and one unrewarding branch. These sections, along with neutral controls, were tested on unbranched foraging trails (see supplementary information) by noting whether individual foragers walking over them did a U-turn. Compared with ants on the control substrate, almost twice as many ants U-turned when walking on substrate from the unrewarding branch near the bifurcation (N_b) (19% and 34%, respectively; $P<0.001$) (Fig. 1a). However, U-turns were as frequent on substrate from the unrewarding branch end (N_e) (27%) as on the control (27%) (NS) (Fig. 1a). Ants U-turned less often on sections from the rewarding trail (stem S, 12%; feeder branch close to the bifurcation F_b , 12%; and feeder-branch end F_e , 13%). These values are significantly lower than those for the relevant control (S, $P<0.001$; F_b , $P<0.05$; F_e , $P<0.001$) (Fig. 1a).

In the same experiment, we also determined whether foragers could detect the negative signal before reaching the substrate on which it had been laid, using walking behaviour (zigzagging versus walking straight) as a bio-assay. Our results show that significantly more ants zigzagged when approaching substrate from an unrewarding branch just after the bifurcation ($P<0.01$) or at the branch end ($P<0.05$) than did controls (Fig. 1b). Conversely, significantly fewer zigzagged when approaching substrate leading to the feeder (S, $P<0.01$; F_b , $P<0.05$; F_e , $P<0.05$) (Fig. 1b).

Our results show that Pharaoh's ants use a sophisticated trail system with a negative, repellent pheromone to mark unrewarding branches. The signal is concentrated at decision points — trail bifurcations⁵. As it is volatile, it provides advance warning — like human road signs situated before junctions. Across a trail network, the pheromone could help direct foragers to food by closing off unrewarding sections. Exactly how negative pheromones enhance foraging efficiency in trail networks is not known, but they might complement attractive trail pheromones^{6,7}




Figure 1 | Identifying the location of the negative pheromone. The ants' response is monitored by their walking behaviour, with U-turning or zigzagging on unbranched trails indicating detection. Test sections: S, 1 mm before bifurcation; F_b and N_b , 3 mm after bifurcation on feeder and non-feeder branches, respectively; F_e and N_e , 60 mm from bifurcation at the ends of feeder and non-feeder branches, respectively. (For details and chi-squared tests, see supplementary information.) **a**, Number of ants that U-turned while walking on different test sections, relative to controls. **b**, Percentage of straight-walking (left bars) or zigzagging (right bars) ants, relative to controls.

used by Pharaoh's ants in trail choice, or they could prevent strong positive feedback by attractive pheromones from locking the system into suboptimal solutions^{1,8}.

Elva J. H. Robinson*, Duncan E. Jackson†, Mike Holcombe†, Francis L. W. Ratnieks*

*Laboratory of Apiculture and Social Insects, Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK
e-mail: e.robinson@dcs.sheffield.ac.uk

†Department of Computer Science, University of Sheffield, Sheffield S1 4DP, UK

1. Sumpter, D. J. T. & Beekman, M. *Anim. Behav.* **66**, 273–280 (2003).
2. Camazine, S. et al. *Self-Organization in Biological Systems* (Princeton Univ. Press, Princeton and Oxford, 2001).
3. Stickland, T. R., Britton, N. F. & Franks, N. R. in *Information Processing in Social Insects* (eds Dethlefsen, C., Deneubourg, J. L. & Pasteels, J. M.) 83–100 (Birkhäuser, Basel, 1999).
4. Britton, N. F., Stickland, T. R. & Franks, N. R. *J. Biol. Syst.* **6**, 315–336 (1998).
5. Jackson, D. E., Holcombe, M. & Ratnieks, F. L. W. *Nature* **432**, 907–909 (2004).
6. Sudd, J. H. *Br. J. Anim. Behav.* **5**, 104–109 (1957).
7. Jeanson, R., Deneubourg, J. L. & Ratnieks, F. L. W. *Physiol. Entomol.* **28**, 192–198 (2003).
8. Beckers, R., Deneubourg, J. L., Goss, S. & Pasteels, J. M. *Insectes Soc.* **37**, 258–267 (1990).

Supplementary information accompanies this communication on Nature's website.

Competing financial interests: declared none.
doi:10.1038/438442a

BRIEF COMMUNICATIONS ARISING online
www.nature.com/bca see Nature contents.