



**CLIMATE CHANGE**  
Find all our news on  
global warming  
<http://www.nature.com/news/infocus/climatechange.html>

before the IPCC report's release, Rahmstorf and his colleagues argue that sea-level rises will be close to the worst-case predictions of climate models (S. Rahmstorf *et al. Science* doi:10.1126/science.1136843; 2007). "If anything, the IPCC has been conservative," he says.

Key sticking points include the inability of global climate models to produce the amount of sea-level rise observed over the past couple of decades and whether ice flow at the bases of glaciers is accelerating or not. How volatile Antarctic and Greenland glaciers might become in a warmer world is therefore pretty much guesswork.

For the first time, the IPCC report predicts how changing climate might affect particular regions of the world. But these forecasts are only in their infancy, modellers warn. For some areas, models predict specific and well understood effects, such as hotter summers in Spain and smaller snowpacks (the accumulation of snow each season) in the Rocky Mountains in the United States. But improved analyses that incorporate clouds, snow and ice into the models must be developed if regional predictions are to become more accurate, says Rasmus Benestad, a climate modeller at the Norwegian Meteorological Institute in Oslo.

Extreme weather is another example of the remaining uncertainties. Climate researchers believe that storms and heavy rainfall will become more frequent as the planet warms. But pinning down where and when that might happen is not so simple.

In the tropics, rising sea-surface temperatures can be linked in a relatively straightforward manner to storm formation, and the case for more intense storms seems more or less settled. But in the mid-latitudes, where atmospheric processes are more complex, some climate models predict more storms whereas others do not.

Improving the models, experts say, requires better data. Gaps and errors in observations are attributable to many causes: snowfall gauges that ice up, oceanographic floats that get lost, and changeovers in satellites that throw off carefully calibrated trends, to name but a few. Cloud and storm records urgently need to be reprocessed using uniform techniques, says Kevin Trenberth, a senior scientist at the National Center for Atmospheric Research in Boulder, Colorado, and coordinating lead author of the report's chapter on surface and atmospheric change.

"The IPCC report is a consensus report, and one that develops over nearly three years," he says. "This means that it is not the leading — or bleeding — edge of the science."

Quirin Schiermeier

## Data keep flooding in

The 2007 report of the Intergovernmental Panel on Climate Change (IPCC) represents the work of thousands of researchers, compiled and summarized by hundreds of climatologists. Nominally, the cut-off for inclusion in the assessment was the end of 2005, allowing a year for the panel to make sense of the vast tracts of data. But notable research arising after that date will not have escaped attention. Here is a round-up of some of the most prominent studies.

### Greenland ice

Greenland is losing ice at an ever-increasing rate, according to data from the GRACE gravity-measuring satellite (J. L. Chen, C. R. Wilson & B. D. Tapley *Science* **313**, 1958–1960; 2006).

### Antarctic air

Weather balloons reveal that the troposphere above Antarctica has warmed by 0.5–0.7 °C per decade over the past 30 years, although it is not clear why (J. Turner, T. A. Lachlan-Cope, S. Colwell, G. J. Marshall & W. M. Connolly *Science* **311**, 1914–1917; 2006).

### Established forests

Old forests keep soaking up atmospheric carbon long after they reach maturity, according to measurements from China. Soil carbon in a forest reserve in Guangdong increased by 68% in 25 years (G. Zhou *et al. Science* **314**, 1417; 2006).

### Atlantic currents

The Gulf Stream, which brings heat from the tropics to the North Atlantic, weakened

by 10% between 1200 and 1850, during the cold spell known as the Little Ice Age. The authors suggest that this demonstrates the link between this ocean current and temperatures in northern Europe (D. C. Lund, J. Lynch-Stieglitz & W. B. Curry *Nature* **444**, 601–604; 2006).

### Atlantic hurricanes

Rising sea-surface temperatures correlate strongly with the observed increase in the number of category 4 and 5 Atlantic hurricanes between 1970 and 2004. Other factors that affect hurricane formation, such as wind shear, do not seem to have increased in line with the upward trend (C. D. Hoyos, P. A. Agudelo, P. J. Webster & J. A. Curry *Science* **312**, 94–97; 2006).

### River runoff

More carbon dioxide in the atmosphere leads to plants losing less water by transpiration, a model suggests. This could affect the amount of fresh water available for human use.

(N. Gedney *et al. Nature* **439**, 835–838; 2006).

### Polar temperatures

Bubbles dating back 150,000 years in an Antarctic ice core show that warming events have tended to seesaw back and forth between the poles (EPICA Community Members *Nature* **444**, 195–198; 2006).

### Ocean temperatures

The upper layers of the oceans cooled, on average, between 2003 and 2005. Factoring in this downturn, the rate of warming in these layers between 1993 and 2005 was equivalent to 0.33 watts per square metre over the whole of the planet's surface (J. M. Lyman, J. K. Willis & G. C. Johnson *Geophys. Res. Lett.* **33**, L18604; 2006).

### Sea levels

If the rate of sea-level rise is proportional to the global rise in temperature since pre-industrial times, sea levels could rise by up to 1.4 metres by 2100 (S. Rahmstorf *Science* **315**, 368–370; 2007).

Michael Hopkin



Greenland ice is melting faster than before.

N. COBBING/STILL PICTURES