

that resources could be simultaneously depleted, so achieving sustainability might be more challenging than was thought.

Ralf Seppelt of the Helmholtz Centre for Environmental Research in Leipzig, Germany, and his colleagues estimated the peak rate of extraction for 27 resources. For 20 of them, mostly renewables such as meat and rice, the peak-rate years occurred between 1960 and 2010, with many clustering around 2006. Only coal, gas, oil, phosphate, farmed fish and renewable energy have yet to peak.

Humans use multiple resources to generate new ones and to meet basic needs, which could explain the synchronicity of peak usage, the authors suggest.


Ecol. Soc. 19, 50 (2014)

MATERIALS

Silicon buckles to form 3D shapes

Researchers have created a variety of small, three-dimensional structures by buckling strips of silicon and other materials.

Turning advanced two-dimensional materials into three-dimensional shapes has proved difficult. John Rogers at the University of Illinois at Urbana-Champaign and his colleagues added hydroxyl groups at specific locations along the length of silicon ribbons as narrow as 800 nanometres and as thin as 100 nm. They also added these groups in specific patterns to an elastic, stretched substrate, and allowed the silicon strips to bond to the substrate.



When the substrate returned to its original shape, the silicon buckled to form a range of structures less than a millimetre wide (pictured), including helices, boxes and flowers. These could then be assembled into larger configurations.

The method uses a variety of materials, such as metals and polymers, and has the potential to make structures for a wide range of electronic devices, the authors say.

Science 347, 154–159 (2015)

EVOLUTION

Mosquitoes gain resistance

A malaria-carrying mosquito inherited insecticide-resistance genes from a related species, around the time that bed nets treated with insecticide were increasingly used in West Africa.

Gregory Lanzaro at the University of California, Davis, and his colleagues analysed DNA from more than 1,000 specimens of *Anopheles coluzzii* and *Anopheles gambiae* in Mali from 2002 to 2012. They found that a group of genes, including one for insecticide resistance, from *A. gambiae* moved into *A. coluzzii* around 2006 when the two species mated.

Campaigns to encourage the use of insecticide-treated bed nets began in 2005 in this region, and the authors suggest that the nets favoured the selection of hybrid, insecticide-resistant mosquitoes.

Proc. Natl Acad. Sci. USA <http://doi.org/x56> (2015)

PHOTONICS

Few photons make 'ghost image'

Physicists have captured an image of a wasp's wing using less than one photon per pixel.

Peter Morris and his colleagues at the University of Glasgow, UK, used a technique called ghost imaging, which uses pairs of photons whose positions

SOCIAL SELECTION

Popular articles on social media

Unveiling secret funding decisions

Many scientists struggle to understand why some grant applications succeed and others fail, perhaps explaining the online popularity of an article calling for increased transparency in the grant peer-review process. Writing in *PLoS Biology*, Daniel Mietchen, an evolutionary biologist at Berlin's Museum of Natural History, argues that, among other things, all successful proposals and their reviews should be released to the public. "Open science is now old hat; here's a call for open research funding processes," tweeted Dorothy Bishop, a neuropsychologist at the University of Oxford, UK. But some see a downside to transparency. James Johnson, a diabetes researcher at the University of British Columbia in Vancouver, Canada, tweeted: "I try to be open, but I've also had desperate people steal my ideas."

PLoS Biol. 12, e1002027 (2014)



Based on data from altmetric.com. Altmetric is supported by Macmillan Science and Education, which owns Nature Publishing Group.

➔ **NATURE.COM**

For more on popular papers: go.nature.com/tm0bc

are inextricably correlated, or entangled. One of each photon pair was either transmitted or absorbed by the wing, while its twin was used to reconstruct the image.

Each photon was collected at a separate detector. To avoid recording stray photons that cause noise, a camera captured the 'image-creating' photon only when its partner from the wing was detected. Applying image-compression techniques, the authors further reduced the number needed to just 0.45 photons per pixel.

Such techniques could be useful in biological imaging when high levels of light could damage the sample, they say. *Nature Commun.* 6, 5913 (2015)

PLANT GENETICS

Maize's journey out of Mexico

DNA from the cobs of ancient maize (corn) shows how the crop was taken to the US southwest from Mexico.

Maize was domesticated from the wild grass teosinte, an inedible weed, more than

6,000 years ago in southern Mexico, and later spread throughout the Americas. A team led by Rute da Fonseca and Thomas Gilbert at the University of Copenhagen analysed nuclear DNA from 32 maize samples from several archaeological sites spanning Mexico and the US southwest (pictured is a 5,000-year-old specimen).

They conclude that maize arrived in the US southwest around 4,000 years ago along a highland route in central Mexico — not by a Pacific coastal route as other studies had suggested. Along the way, maize evolved into a sweeter and more drought-tolerant crop.

Nature Plants <http://doi.org/x6p> (2015)

➔ **NATURE.COM**

For the latest research published by Nature visit: www.nature.com/latestresearch

