

www.nature.com/ejhg

# ARTICLE

# Supernumerary marker chromosomes in man: parental origin, mosaicism and maternal age revisited

John A Crolla<sup>\*,1,2</sup>, Sheila A Youings<sup>1</sup>, Sarah Ennis<sup>2</sup> and Patricia A Jacobs<sup>1,2</sup>

<sup>1</sup>Wessex Regional Genetics Laboratory, Salisbury District Hospital, Salisbury, Wiltshire, UK; <sup>2</sup>Division of Human Genetics, Duthie Building, Southampton General Hospital, Southampton, Hampshire, UK

The details of all cytogenetic abnormalities diagnosed in the Wessex Regional Genetics Laboratory (WRGL) since 1967 to the present day have been recorded in the Salisbury Treasury of Interesting Chromosomes (STOIC). From this resource, we identified 137 patients with constitutional autosomal supernumerary marker chromosomes (SMC) ascertained in four principal groups: (i) 37% with abnormal phenotypes; (ii) 7% couples with reproductive difficulties; (iii) 47% antenatal diagnoses and (iv) 9% miscellaneous. Overall, 81 (59%) SMCs were mosaics and 56 (41%) nonmosaics. Of the 109 cases with known parental origins, 70% were *de novo*, 19% maternally and 11% paternally inherited. The chromosomal origins of 112/137 (82%) of the SMCs have been determined by fluorescence *in situ* hybridization (FISH). In all, 36/112 (32%) were derived from nonacrocentric autosomes, and 76/112 (68%) from the acrocentric autosomes 13/21, 14, 15 and 22. Of these acrocentric SMCs, 39 (51%) were derived from chromosome 15, so that SMC(15) constituted 39/112 (35%) of all SMCs with known chromosomal origins. The frequencies with which mosaicism was observed varied considerably according to the chromosomal origin of the SMCs and accounted for 8/39 (20%) SMC(15), 13/37 (35%) SMCs from other acrocentrics and 25/36 (69%) of nonacrocentric SMCs. The data were analysed for parental age effects, and only *de novo* SMC(15)s were found to be associated with a significantly increased maternal age.

*European Journal of Human Genetics* (2005) **13**, 154–160. doi:10.1038/sj.ejhg.5201311 Published online 27 October 2004

Keywords: supernumerary marker chromosomes; maternal age; parental origin; chromosomal origin; fluorescence *in situ* hybridization (FISH); mosaicism

### Introduction

Over the past 30 years, a large number of reports have been directed towards the identification, classification and clinical significance of human constitutional supernumerary marker chromosomes (SMC). Historically, Buckton *et al*<sup>1</sup> first described a correlation between SMCs and their possible phenotypic effects when they showed that the

ascertainment of SMCs among individuals in psychiatric institutions was 3.27/1000 compared to their frequency of only 0.24/1000 among consecutive newborns.

Population studies on antenatally ascertained cases show that SMCs occur with a frequency ranging from  $0.4/1000^2$ to  $0.8/1000.^{3,4}$  Hook and Cross<sup>3</sup> also published data suggesting that antenatally ascertained SMCs were associated with advanced maternal age. Among these early studies, the reported overall frequencies of mosaicism involving SMCs ranged from 27 to  $66\%.^{1-3}$  These early studies using conventional cytogenetic techniques also divided the SMCs based on whether they had recognizable satellites and therefore assumed to be derived from the

<sup>\*</sup>Correspondence: Dr JA Crolla, Wessex Regional Genetics Laboratory, Salisbury District Hospital, Salisbury, Wiltshire SP2 8BJ, UK.

Tel: +44 1722 429069; Fax: +44 1722 338095;

E-mail: john.crolla@salisbury.nhs.uk

Received 22 June 2004; revised 30 August 2004; accepted 1 September 2004

Supernumerary marker chromosomes and maternal age JA Crolla *et al* 

acrocentric autosomes, and whether they were without satellites and were assumed to be comprised of the nonacrocentric chromosomes. From these early data, it was clear that SMCs derived from the acrocentric autosomes comprised the majority population, for example, 64% acrocentrics *vs* 36\% nonacrocentrics.<sup>2</sup>

Studies of the cytogenetic and phenotypic effects of SMCs gained a fresh impetus when fluorescence *in situ* hybridization (FISH) was utilized to identify their chromosomal origins.<sup>5–10</sup> From these studies further insights were obtained into karyotype/phenotype correlations, particularly with respect to SMCs derived from chromosomes 15<sup>11–13</sup> and 22.<sup>14–16</sup> Delineations of possible phenotypic effects in SMC derived from the other autosomes, however, remains more problematic (reviewed by Crolla<sup>17</sup>).

From the combined conventional and molecular cytogenetic studies of autosomal SMCs, a number of trends have emerged from the literature. Principally, these include: (a) SMCs overall are associated with advanced maternal age, (b) among acrocentric SMCs, the majority are derived from chromosome 15 and (c) mosaic and nonmosaic cells lines are observed with variable frequencies depending on whether or not the SMCs have satellites.<sup>2-4</sup>

We have re-examined the above parameters using a population of autosomal SMCs diagnosed and characterized in a contiguous geographical area in the UK over a continuous period of 30 years. The results of these analyses in the context of previous studies of SMCs are presented.

#### Materials and methods

All the cases presented were referred to the WRGL for routine conventional chromosome analysis. Analyses were carried out on either PHA-stimulated peripheral blood and/ or amniotic fluid or CVS fibroblasts following standard culture and chromosome preparation protocols used to produce conventional cytogenetic preparations. Chromosome analyses were carried out following standard Gbanding to a minimum of 550 bands, and in mosaic cases, 30 metaphases were scored.

FISH was carried out using standard techniques based on a modification of Pinkel *et al*,<sup>18</sup> and the probes used have been reported previously (eg Crolla<sup>19</sup> and Crolla *et al*<sup>20</sup>).

Retrieval and analyses of the SMC data were facilitated by the Salisbury Treasury of Interesting Chromosomes (STOIC) database. For the purpose of these studies, the populations have been subdivided into four groups, *viz* nonacrocentric SMC; acrocentric SMCs excluding chromosome 15 (ie 13/21, 14 and 22); chromosome 15-derived SMCs and SMCs with unknown chromosomal origins.

### Statistical analyses

Statistical analyses were performed using SAS (SAS Institute Inc., Cary, NC, USA) for regression, logisitic regression



Figure 1 Mean maternal age in England and Wales from 1938 to 2000. Notes: (1) Mean maternal age data for England and Wales may not be the most accurate control for our sample. The available data suggest that mean maternal age is approximately 1 year earlier for mothers in Wales compared to mothers in England and this could have an anticonservative effect on our analysis. (2) Mean maternal age data for England and Wales is given irrespective of birth order. Our Wessex data may be enriched for non-firstborn individuals. (3) Despite concerns outlined in 1 and 2, all our results from our regression show the intercept as not being significantly different from zero, which suggests the control population is appropriate. It is possible that individuals in the marker chromosome database have been ascertained after reproductive difficulties and so maternal age may be advanced compared to the mean.

means tests and  $\chi^2$  statistics in frequency tables. Data analysis was carried out using date of birth as well as the maternal age at the birth of each patient. Paternal age at birth is included where given, but these data are sparse. As dates of birth for patients range from 1928 to 2003, changing trends of mean maternal age over this time period were accounted for (Figure 1). This was achieved by taking the mean maternal age for England and Wales for the relevant years and calculating the deviation from this mean for each of the mothers of individuals with marker chromosomes. This derived variable was labelled 'deviate' and used as the dependent variable for a number of the regression analyses examining the effect(s) of maternal age.

#### Results

### Ascertainment and study population

The 137 patients in this study group with autosomal SMCs were derived from four principal ascertainment groups: (1) 51(37%) with abnormal phenotypes; (2) nine (7%) presenting with reproductive difficulties, principally multiple miscarriages; (3) 64 (47%) antenatal diagnoses and (4) 13 (9%) with other referral reasons (Table 1). These patterns of ascertainment were the same irrespective of whether the

SMCs were acrocentric or nonacrocentric in structure. This study has excluded SMCs that are resolvable by conventional light microscopy and associated with a defined clinical phenotype, that is, i(12p), i(18p).

#### Chromosomal origins of the SMCs

A total of 112 of the 137 SMCs were further characterized by FISH and the distribution of chromosomal origins is shown in Table 2. In all, 36 (32%) of the SMCs with known chromosomal origins were nonacrocentrics and 76 (68%) were derived from the acrocentric autosomes. Of the 76 acrocentrics, 14 (18%) were SMC(13/21), 13 (17%) were SMC(14), 10 (13%) were SMC(22) and 39 (51%) were SMC(15). Overall therefore, where the chromosomal origin was known, SMC(15) comprised 51% of all acrocentric SMCs and 35% of the SMC population overall.

#### Parental origin

The parental origins of 109/137 (79%) cases were known, of which 76 (70%) were *de novo*, 21 (19%) maternally and 12 (11%) paternally inherited (Table 3). However, there were marked differences in parental origin among the groups: 27 of the 30 nonacrocentric SMCs (90%) were *de novo*, and three (10%) maternally transmitted, whereas of the 62 acrocentric SMCs with known parental origins, 37 (60%) were *de novo*, 15 (24%) maternally and 10 (16%) paternally inherited (Table 3).

#### Mosaicism

A total of 81 (59%) of the 137 cases were nonmosaic and 56 (41%) mosaic (Table 4). In all, 25 nonacrocentric SMCs (69%) were mosaics, with the remaining 11 (31%) nonmosaic. By comparison, 21 (28%) of all the acrocentric SMCs were mosaics so that the frequency of mosaicism

**Table 1**Ascertainment and distribution of mosaicism intotal population (n = 137)

Ascertainment	Nonmosaic	Mosaic	Total
Abnormal phenotype	33	18	51 (37%)
Reproductive problems	5	4	9 (7%)
Prenatal diagnosis	31	33	64 (47%)
Other	12	1	13 (9%)
Total	81	56	137

seen among nonacrocentric SMCs was twice that seen in SMCs derived from the acrocentric chromosomes. However, within the acrocentric SMCs, mosaicism was observed considerably less frequently among SMC(15)s, that is, (20%) compared with the SMCs derived from 13/21, 14 and 22 (38%) (see Table 5).

#### Analyses of parental age effects

For the analyses of parental ages, the marker chromosomes were also categorized into four classes according to their chromosomal composition, *viz*: (A) unknown, n = 25; (B) nonacrocentrics, n = 36; (C) acrocentrics excluding chromosome 15, n = 37 and (D) chromosome 15, n = 39.

A number of the patients in the database are known to have been ascertained through advanced maternal age. These individuals fall into the antenatal diagnosis ascertainment groups representing serum increased risk, nuchal translucency risk and maternal age, respectively. We also expected that individuals in the ascertainment group representing a familial marker chromosome may have older mothers. We tested this expectation by regressing the dependent variable 'deviate' against these groups.

**Table 3** Parental origins of SMCs with known chromosomal origins (n = 109)

	De	novo	Μ	lat	Pat		
Category	n	%	n	%	n	%	
Nonacrocentric	27	90	3	10	0	0	
All acrocentric	37	60	15	24	10	16	
Acrocentric excluding SMC(15)	19	63	7	24	4	13	
SMC(15) Overall	18 76	56 70	8 21	25 19	6 12	19 11	

**Table 4**Parental origins and distribution of mosaicism intotal population (n = 137)

Origin	Nonmosaic	Mosaic	Total
De novo	36 (47%)	40 (53%)	76 (56%)
Mat	18 (86%)	3 (14%)	21 (15%)
Pat	12 (100%)	0` ´	12 (9%)
Unknown	15 (54%)	13 (46%)	28 (20%)
Total	81 (59%)	56 (41%)	137 (100%)

Table 2 Supernumerary marker chromosomes. Chromosome assignment by FISH

Chromosome	1	2	3	4	5/19	6	7	8	9	10	11	12	13/21	14	15	16	17	18	19	20	21	22	?	Total
Nonmosaic Mosaic	1 0	1 0	0 1	0 2	1 3	0 2	0 0	2 1	0 3	0 0	1 0	0 7	7 7	9 4	31 8	2 5	0 0	2 0	1 1	0 0	0 0	8 2	15 10	81 56
Total	1	1	1	2	4	2	0	3	3	0	1	7	14	13	39	7	0	2	2	0	0	10	25	137

**European Journal of Human Genetics** 

156

Category	Mosaic (%)	Nonmosaic (%)	Total (%)
Nonacrocentric Acrocentric excl SMC(15) SMC(15) only	25 (69%) 13 (35%) 8 (20%)	11 (31%) 24 (65%) 31 (80%)	36 (32%) 37 (33%) 39 (35%)
Total	46 (41%)	66 (59%)	112 (100%)

**Table 5** Distribution of mosaicism in SMC with known chromosomal origins (n = 112)

**Table 6**Mean deviate from maternal age

	N <sub>obs</sub>	N <sub>missing</sub>	N <sub>nonmissing</sub>	Mean	SE	Min	Max	Origin
A Chromosome class								
A – Unknown	15	9	6	3.2167	1.5217	-1.8	8.3	
B – Nonacrocentrics	17	6	11	-0.9364	1.3287	-7.9	5.6	
C – Acrocentrics excluding chromosome 15	22	8	14	1.3214	1.6586	-12.0	9.3	
D – Chromosome 15	24	6	18	7.2944	1.4817	-4.6	21.6	
D group parental origin+all others	61	35	26	1.1971	0.9017	-12.0	9.3	0
D aroup unknown origin	6	3	3	4.9333	2.0169	0.9	7	1
D group <i>de novo</i>	11	11	0	9.2818	1.9709	0.4	21.6	2
	Estimate		SE		F	Р		
B Regression of deviation on X								
Intercept	1.1851		0.9198		1.66	0.2039		
D group <i>de novo</i>	4.0292		0.9391		18.41	0.0001		

As expected, mothers of patients ascertained through advanced maternal age were significantly older than the mothers of the rest of the cohort (F = 16.04, P < 0.0001). The same was true for mothers of patients ascertained through a familial marker chromosome, although the effect here (as a partial regression after allowance for maternal age ascertainment) was less striking (F = 4.44, P = 0.04). These results confirmed that individuals ascertained in these ways should be excluded from further analyses. No individuals were recorded as ascertained through the nuchal translucency test. Surprisingly, we found no significant increase in maternal age in the group ascertained through serum increased risk; however, we also excluded this group on the basis of theoretical bias. A total of 59 observations were therefore excluded from further analyses, leaving a cohort of 78 individuals for our study (Table 6).

Regression analysis, again using 'deviate' as the dependent variable, identified class D [SMC(15)] as having a significantly advanced maternal age (F = 14.57, P = 0.0004), whereas classes A, B and C showed no significant increase in maternal age (see Table 6A and B). We then included the origin of the marker chromosome into our analysis and found that the significance of the advanced maternal age effect found in class D was increased in those patients with *de novo* marker chromosomes (F = 18.4072, P < 0.0001). We also found that the increase in maternal age was linear when those with either maternally or paternally inherited chromosomes were coded as 'Origin' = 0, those whose marker chromosome origin was unknown (and so presumably a mix of inherited and *de novo*) were coded 'Origin' = 1 and those with *de novo* chromosomes were coded 'Origin' = 2. This finding was expected and confirmed the heterogeneity of those marker chromosomes of unassigned origin.

To test for any additive effect of mosaicism, we recoded the Origin variable. Assuming mosaicism arises mitotically and nonmosaicism meiotically, group D *de novo* mosaic individuals were recoded as 'Origin' = 0. We had 11 informative observations for individuals in group D of *de novo* origin, two of which are mosaics. Regressing this recoded Origin variable against the dependent variable 'deviate' did not result in any increase of significance. This would suggest that there is no evidence for any effect of mosaicism with regard to maternal age; however, the data are few and perhaps more data are needed to observe a small effect.

We have examined the distribution of the Prader–Willi/ Angelman Syndrome Critical region (PWACR) in group D with regard its origin. We find a significant deficiency of observations for SMC(15)s containing two copies of the PWACR (CR++) (Fisher's exact test;  $p_{two-tailed} = 0.018$ ) in the maternally and paternally derived markers with a relative abundance of CR– chromosomes (ie SMCs not containing the PWACR) when compared with those observations of *de novo* origin. Observations of unknown origin appear to have a mixed CR++ and CR– composition.

## Discussion

### Total population and methods of ascertainment

The data used for this study are derived from the STOIC database, which records all chromosome abnormalities reported following routine clinical ascertainment in the Wessex Regional Genetics Laboratory (WRGL) since 1967. We have therefore excluded for the purposes of the present study any cases that were recruited and analysed as part of defined SMC research protocols.<sup>11,16,19,20</sup> The population served by the WRGL, approximately 2.5 million, over the period of the study has been stable and has not witnessed significant changes in migration patterns as seen in other parts of the UK.

The WRGL reports constitutional cytogenetic chromosome results from both perinatal and postnatal patient groups. The postnatal ascertainment patterns have remained relatively unchanged consisting largely of dysmorphic perinates and infants, patients with congenital abnormalities, developmental delay and/or idiopathic mental retardation and couples with reproductive difficulties (Table 1). By contrast, over the same time period, the prenatal referral pattern has changed from predominantly maternal age referrals to ascertainment following either serum screening and/or ultrasound measurements of nuchal thickness and congenital abnormality scans (data not shown - see below). Overall, 47% of all the SMCs were ascertained during prenatal diagnosis, and among those where the chromosomal origin was determined, 29% were SMC(15)s, 33% SMC(other acrocentrics) and 37% nonacrocentric SMCs.

# The distribution of SMCs by chromosomal origin and ascertainment

Using FISH, we were able to identify the autosomal chromosomal origins of 112/137 (82%) of the SMCs in our study population, of which 76 (68%) were derived from the acrocentric autosomes with all other autosomes represented at least once except for chromosome 7, 10, 17, 20 and 21 (Table 3). SMCs derived from chromosome 15 were predominantly of the type usually referred to as inv dup(15)s and accounted for 39/112 (35%) of all the chromosomally identified SMCs and for approximately one-half of all the acrocentric SMCs observed. The results of the present study confirm that SMC(15)s are the most frequently diagnosed SMCs in man. Some previous reports have suggested that SMC(15)s account for over half of all SMCs detected.<sup>21</sup> However, more recently, FISH-based

studies have shown that the distribution of antenatally ascertained acrocentric SMCs varies from 77 to 81%,<sup>4,22,23</sup> and within these studies SMC(15)s accounted for 76, 45 and 33% of all the acrocentric markers identified. By combining the data from the present study with the three studies quoted above, a frequency overall of 36% SMC(15) encountered during prenatal diagnosis is obtained.

### Mosaicism and the chromosomal origins of SMCs

Our data confirm previous reports showing striking differences in the distribution of mosaicism when the SMCs are classified according to their chromosomal origins. Overall, 59% of SMCs were found in association with a normal cell line, but the distribution was markedly asymmetric with 28% of the acrocentric SMCs compared with 69% of nonacrocentric SMCs being mosaic. Previous larger studies have also reported differences in the levels of mosaicism depending on whether the SMC was satellited or nonsatellited (pre-FISH)<sup>1,2</sup> or acrocentric or nonacrocentric (post-FISH). In these studies, the level of mosaicism overall was reported to range from 13 to 50% (ave. 46%), but with ~30% of the acrocentric SMCs being classified as mosaics.<sup>2,4,10,22,23</sup>

Further analyses of our data show that within the acrocentric SMCs, 20% of the SMC(15) and SMC(22) were mosaic compared with 31% for SMC(14) and 50% for SMC(13/21) respectively. Previous reports on SMC(15) have shown that on average ~85% of SMC(15)s present as nonmosaics<sup>13,24,28</sup> and the majority of SMC(22) reported are also nonmosaics.<sup>25,26</sup>

The fact that the majority of SMC(15) are nonmosaics suggests a meiotic origin of these markers and this is supported by the observations that all *de novo* SMC(15)s characterized molecularly have been shown to be maternal in origin,<sup>11,24,27–29</sup> and in a proportion of those containing additional copies of the PWACR, molecular studies show that they are divided equally between being inter- and intrachromosomal in origin.<sup>30</sup> As far as we are aware, no comparable and detailed molecular studies have been carried out on SMCs derived from chromosome 13/21, 14 and 22, and so it is not clear if the mechanisms responsible for the formation of SMC(15) are unique to this chromosomal class.

### Parental age effects in SMCs

Hook and Cross<sup>3</sup> reported a significant maternal age effect in association with *de novo* SMCs in a study population of 75 000 prenatal cytogenetic diagnoses. This study was carried out before the advent of FISH so that the chromosomal subclasses of SMC were not analysed independently as in the present study. As far as we are aware, this original observation with respect to parental age effects associated with SMCs has not been re-examined, so it is generally assumed that all SMCs, irrespective of

158

maternal age effect. Our results have shown that although there is no overall maternal age effect associated with SMCs, de novo SMCs derived from chromosome 15 are associated with a significantly increased maternal age. It has previously been shown that there is an exponential increase with maternal age associated with maternal meiosis I nondisjunction involving chromosome 15<sup>31</sup> and with uniparental disomy (UPD) for chromosome 15.32 There have also been a number of hypotheses to account for the relatively high frequency of SMC(15)s in humans, and it is true that all of the SMC(15)s studied molecularly have been found to have arisen following female meiotic errors. The maternal age effect associated with de novo SMC(15)s would suggest therefore a common origin for both trisomy 15 following nondisjunction at meiosis I with subsequent complete trisomy rescue leading to UPD(15) or, incomplete trisomy rescue resulting in the formation of a SMC(15). In either event, the common origin appears to be a maternal agerelated disruption of normal disjunction of the chromosome 15 bivalent during female meiosis. As discussed above, lack of comparable data concerning SMCs from the other acrocentric chromosomes does not allow a conclusion to be drawn on the uniqueness or otherwise of the observations concerning chromosome 15.

Our previous studies classified the larger SMC(15)s, which predominantly contained two copies of the PWACR into two broad groups based on their most distal breakpoint. The larger group ( $\sim$ 70%) were generally asymmetrical with two distal breakpoints, both located distal to the common PWACR distal breakpoint (BP3). The remaining  $\sim$  30% of SMC(15)s by comparison reveal a more heterogeneous distribution of breakpoints most of which had occurred within BP3.<sup>33</sup> Both groups of SMC(15)s were also shown to have asymmetrical breakpoints and, furthermore, the two main breakpoint regions occur at sites of duplicated genomic segments,<sup>34</sup> suggesting that these duplicons are responsible, at least in part, for the increased instability in this region, and the most popular theory for the formation of SMC(15)s is a U-type exchange between homologous chromosomes during meiosis I followed by illegitimate fusion of the chromatids and nondisjunction.<sup>35,36</sup> A combination of these events, with maternal age related nondisjunction as the primary mechanism, may go some way to explain the relatively high frequency with which SMC(15)s are ascertained in the human population.

In conclusion, this study has shown that autosomal SMCs derived from acrocentric and nonacrocentric chromosomes differ in the proportion, which have arisen de novo, and also in the frequency with which the SMC is seen associated with a normal cell line. Furthermore, we have shown that a significant maternal age effect is restricted to de novo SMCs derived from chromosome 15.

#### Acknowledgements

We thank the many clinical cytogeneticists and molecular cytogeneticists who carried out the karyotyping and FISH analyses. This work was partly funded by the Medical Research Council and the Wellcome Trust.

#### References

- 1 Buckton KE, Spowart G, Newton MS, Evans HJ: Forty four probands with an additional 'marker' chromosome. Hum Genet 1985; 69: 353-370.
- 2 Warburton D: De novo balanced chromosome rearrangements and extra marker chromosomes identified at prenatal diagnosis: clinical significance and distribution of breakpoints. Am J Hum Genet 1991; 49: 995-1013.
- 3 Hook EB, Cross PK: Extra structurally abnormal chromosomes (ESAC) detected at amniocentesis: frequency in approximately 75000 prenatal cytogenetic diagnoses and associations with maternal and paternal age. Am J Hum Genet 1987; 40: 83-101.
- 4 Blennow E, Bui TH, Kristoffersson U et al: Swedish survey on extra structurally abnormal chromosomes in 39105 consecutive prenatal diagnoses: prevalence and characterization by fluorescence in situ hybridization. Prenat Diagn 1994; 14: 1019-1028.
- 5 Blennow E, Anneren G, Bui TH, Berggren E, Asadi E, Nordenskjold M: Characterization of supernumerary ring marker chromosomes by fluorescence in situ hybridization (FISH). Am J Hum Genet 1993; 53: 433-442.
- 6 Callen DF, Ringenbergs ML, Fowler J, Freementle J, Haan EA: Small marker chromosomes in man: origin from pericentric heterochromatin of chromosomes 1, 9 and 16. J Med Genet 1990; 27: 155-159.
- 7 Callen DF, Eyre H, Yip MY, Freemantle J, Haan EA: Molecular cytogenetic and clinical studies of 42 patients with marker chromosomes. Am J Med Genet 1992; 43: 709-715.
- Crolla JA, Dennis NR, Jacobs PA: A non-isotopic in situ hybridisation study of the chromosomal origin of 15 supernumerary marker chromosomes in man. J Med Genet 1992; 29: 699-703.
- 9 Friedrich U, Nielsen J: Bisatellited extra small metacentric chromosome in newborns. Clin Genet 1974; 6: 23-31.
- Gravholt CH, Friedrich U: Molecular cytogenetic study of supernumerary marker chromosomes in an unselected group of children. Am J Med Genet 1995; 56: 106-111.
- 11 Crolla IA. Harvey IF. Sitch FL. Dennis NR: Supernumerary marker 15 chromosomes: a clinical, molecular and FISH approach to diagnosis and prognosis. Hum Genet 1995; 95: 161-170.
- 12 Leana-Cox J, Jenkins L, Palmer CG et al: Molecular cytogenetic analysis of inv dup(15) chromosomes, using probes specific for Prader–Willi/Angelman syndrome the region: clinical implication. Am J Hum Genet 1994; 54: 748-756.
- 13 Mignon C, Malzac P, Moncla A et al: Clinical heterogeneity in 16 patients with inv dup 15 chromosome: cytogenetic and molecular studies, search for an imprinting effect. Eur J Hum Genet 1996; 4: 88-100.
- 14 McDermid HE, Duncan AMV, Brasch KR et al: Characterization of the supernumerary chromosome in cat eye syndrome. Science 1986: 232: 646-648.
- 15 Mears AJ, ElShanti H, Murray JC, McDermid HE, Patil SR: Minute supernumerary ring chromosome 22 associated with cat eye syndrome: Further delineation of the critical region. Am J Hum Genet 1995; 57: 667-674.
- 16 Crolla JA, Howard P, Mitchell C, Long FL, Dennis NR: A molecular and FISH approach to determining karyotype and phenotype correlations in six patients with supernumerary marker(22) chromosomes. Am J Med Genet 1997; 72: 440-447.
- 17 Crolla JA: FISH and molecular studies of autosomal supernumerary marker chromosomes excluding those derived

from chromosome 15. II. A review of the literature. *Am J Med Genet* 1998; **75**: 367–381.

- 18 Pinkel D, Landegent J, Collins C *et al*: Fluorescence *in situ* hybridization with human chromosome-specific libraries: detection of trisomy 21 and translocations of chromosome 4. *Proc Natl Acad Sci USA* 1988; **85**: 9138–9142.
- 19 Crolla JA: FISH and molecular studies of autosomal supernumerary marker chromosomes excluding those derived from chromosome 15: II. Review of the literature. *Am J Med Genet* 1998; **75**: 367–381.
- 20 Crolla JA, Long F, Rivera H, Dennis NR: FISH and molecular study of autosomal supernumerary marker chromosomes excluding those derived from chromosomes 15 and 22: I. Results of 26 new cases. *Am J Med Genet* 1998; **75**: 355–366.
- 21 Webb T: Inv dup(15) supernumerary marker chromosomes. *J Med Genet* 1994; **31**: 585–594.
- 22 Hastings RJ, Nisbet DL, Waters K, Spencer T, Chitty LS: Prenatal detection of extra structurally abnormal chromosomes (ESACs): new cases and a review of the literature. *Prenat Diagn* 1999; 19: 436–446.
- 23 Brondum-Nielsen K, Mikkelsen M: A 10-year survey, 1980–1990, of prenatally diagnosed small supernumerary marker chromosomes, identified by FISH analysis. Outcome and follow-up of 14 cases diagnosed in a series of 12 699 prenatal samples. *Prenat Diag* 1995; 15: 615–621.
- 24 Leanna-Cox J, Jenkins L, Palmer CG *et al*: Molecular cytogenetic analysis of inv dup(15) chromosomes, using probes specific for the Prader–Willi/Angelman syndrome region: clinical implications. *Am J Hum Genet* 1994; **54**: 748–756.
- 25 Schinzel A, Schmid W, Fraccaro M *et al*: The 'cat eye syndrome': Dicentric small marker chromosome probably derived from a no.
  22 (tetrasomy 22pter→q11) associated with a characteristic phenotype. *Hum Genet* 1981; 57: 148–158.
- 26 Crolla JA, Howard P, Mitchell C, Long FL, Dennis NR: A molecular and FISH approach to determining karyotype and phenotype correlations in six patients with supernumerary marker(22) chromosomes. *Am J Med Genet* 1997; **72**: 440–447.

- 27 Robinson WP, Binkert F, Gine R *et al*: Clinical and molecular analysis of five inv dup(15) patients. *Eur J Human Genet* 1993; 1: 37–50.
- 28 Roberts SE, Maggouta F, Thomas NS, Jacobs PA, Crolla JA: Molecular and fluorescence *in situ* hybridization characterization of the breakpoints in 46 large supernumerary marker 15 chromosomes reveals an unexpected level of complexity. *Am J Hum Genet* 2003; **73**: 1061–1072.
- 29 Huang B, Crolla JA, Christian SL *et al*: Refined molecular characterization of the breakpoints in small inv dup(15) chromosomes. *Hum Genet* 1997; **99**: 11–17.
- 30 Wandstrat AE, Leanacox J, Jenkins L, Schwartz S: Molecular cytogenetic evidence for a common breakpoint in the largest inverted duplications of chromosome 15. *Am J Hum Genet* 1998; 62: 925–937.
- 31 Robinson WP, Kuchinka BD, Bernasconi F *et al*: Maternal meiosis I non-disjunction of chromosome 15: dependence of the maternal age effect on level of recombination. *Hum Mol Genet* 1998; 7: 1011–1021.
- 32 Robinson WP, Langlois S, Schuffenhauer S *et al*: Cytogenetic and age-dependent risk factors associated with uniparental disomy 15. *Prenat Diag* 1996; **16**: 837–845.
- 33 Christian SL, Bhatt NK, Martin SZ *et al*: Integrated YAC contig map of the Prader-Willi/Angelman region on chromosome 15q11-q13 with average STS spacing of 35 kb. *Gen Res* 1998; 8: 146-157.
- 34 Christian SL, Fantes JA, Mewborn SK, Huang B, Ledbetter DH: Large genomic duplicons map to sites of instability in the Prader-Willi/Angelman syndrome chromosome region (15q11-q13). *Hum Mol Genet* 1999; **8**: 1025-1039.
- 35 Schreck RR, Berg WR, Erlanger BF, Miller OJ: Preferential derivation of abnormal human G-group-like chromosomes from chromosome 15. *Hum Genet* 1997; **36**: 1–12.
- 36 Martinsson T, Johannesson T, Vujic M *et al*: Maternal origin of inv dup(15) chromosomes in infantile autism. *Eur Child Adolesc Psychiatry* 1996; 5: 185–192.