

Erratum

Effect of hysterectomy on incidence trends of endometrial and cervical cancer in Finland 1953–2010

R Luoto, J Raitanen, E Pukkala and A Anttila

British Journal of Cancer (2004) 91, 1979. doi:10.1038/sj.bjc.6602288 www.bjcancer.com
 © 2004 Cancer Research UK

Correction to: *British Journal of Cancer* (2004) 90, 1756–1759. doi: 10.1038/sj.bjc.6601763

Due to an error, Table 1 and some text on page 1756 of the above paper were shown incorrectly. The correct Table 1 and text are given below:

Table 1 Number (N) and incidence of total and partial hysterectomies in Finland 1991–99, and incidence of endometrial and cervical cancer with and without correction for hysterectomies

Year	Overall incidence of hysterectomy ^a		Incidence of endometrial cancer/10 ⁶		Incidence of total hysterectomy ^b		Incidence of cervical cancer/10 ⁶	
	N	/10 ⁶	Uncorrected	Corrected	N	/10 ⁶	Uncorrected	Corrected
1991	10333	3162	134	158	8237	2500	28	30
1992	10619	3185	137	162	8398	2497	33	35
1993	10571	3089	142	171	8733	2535	37	39
1994	10554	3022	136	165	8890	2531	37	40
1995	10169	2855	135	166	8900	2487	44	48
1996	10759	2981	153	190	9923	2741	44	47
1997	11816	3247	142	179	10935	3001	39	42
1998	12227	3315	159	203	11351	3071	45	49
1999	11956	3208	146	188	11246	3017	38	42

^aAll incidence rates are adjusted for age to the world standard population. ^bNumber of women with hysterectomy, cervix removed.

On page 1756 of this article, the second sentence in the Results section should read as follows:

The incidence of hysterectomies with cervix removed increased from 2500 in 1991 to 1717 per million women in 1999.