

Side effects during chemotherapy predict tumour response in advanced colorectal cancer

B Schuell^{*,1}, T Gruenberger², GV Kornek¹, N Dworan¹, D Depisch³, F Lang⁴, B Schneeweiss⁵ and W Scheithauer¹

¹Division of Clinical Oncology, Department of Internal Medicine I, University Hospital, Waehringer Guertel 18-20, Vienna A-1090, Austria; ²Department of Surgery, University Hospital, Waehringer Guertel 18-20, Vienna A-1090, Austria; ³Department of Surgery, Wr.Neustadt General Hospital, Corvinusring 3-5, Wr.Neustadt A-2700, Austria; ⁴Department of Surgery, Neunkirchen General Hospital, Peischinger Strasse 19, Neunkirchen A-2620, Austria; ⁵Department of Internal Medicine, Kirchdorf General Hospital, Kirchdorf an der Krems A-4560, Austria

To investigate whether a relationship between chemotherapy-associated adverse events and treatment efficacy exists, we have analysed the toxicity, objective response and survival data of 303 patients with advanced colorectal cancer. Patients were divided into two groups: the first with beneficial effect (I, $n = 245$), and the second with progressive disease (II, $n = 58$). Differences in terms of incidence rates, type and severity of adverse events were analysed with univariate and multivariate models. The median number of side effects in group I was 6 vs 4 in group II (OR = 1.342; $P = 0.0001$). An inverse correlation between disease control and treatment tolerance was confirmed when side effects were analysed according to severity and type of treatment-associated toxicities (haematological: $P = 0.0005$ vs nonhaematological $P = 0.0001$). When median survival was analysed according to the number of adverse events, it was 10 (95% CI, 3–7), 16 (14–18), and 18 (16–20) months in case of 0–1, 2–5, and ≥ 6 adverse events, respectively ($P = 0.01$). In conclusion, the results of this analysis suggest that occurrence of side effects during chemotherapy in advanced colorectal cancer is an independent and reliable prognostic indicator for response and survival.

British Journal of Cancer (2005) **93**, 744–748. doi:10.1038/sj.bjc.6602783 www.bjcancer.com

Published online 13 September 2005

© 2005 Cancer Research UK

Keywords: colorectal cancer; side effects; chemotherapy; treatment efficacy

Colorectal cancer is one of the most common malignancies in the western world (Landis *et al*, 1999). Nearly one-third of all patients initially present with locally advanced, inoperable tumours and/or distant metastases (Williams *et al*, 1995; Midgley and Kerr, 1999), and approximately 50% of those having undergone potential curative surgery will ultimately develop recurrent disease. Median survival of these patients treated with best supportive care alone is approximately 6 months (Scheithauer *et al*, 1993); if treated with modern combination chemotherapy, median survival ranges from 17 to 21 months (Grothey *et al*, 2004). The effectiveness of conventional intravenous 5-fluorouracil (FU) given with or without biochemical modulators such as leucovorin (LV) or methotrexate, in fact, has been considerably improved with the recent development of a number of new, promising anticancer agents. These include the third generation 1,2-diaminocyclohexane-platinum derivate oxaliplatin, and the topoisomerase I inhibitor irinotecan. When combined with FU/LV or the oral FU prodrug capecitabine, they seem to exert a synergistic effect which results in major improvement in overall response rate and progression-free survival, occasionally in overall survival (Grothey *et al*, 2004).

Potential prognostic markers determining response and/or survival have been described in numerous studies, and include a variety of clinical and laboratory parameters. In the largest series, a multivariate analysis of 3825 patients treated with FU-based

chemotherapy, performance status, the number of metastatic sites, alkaline phosphatase, and the leukocyte count were identified as the most relevant parameters (Köhne *et al*, 2002).

As it concerns the relationship between toxicity incidence and treatment efficacy, only historical data are available. As indicated by Moertel in 1969, mild to moderate toxicity as objectively calibrated by leukocytopenia seems to result in optimal response to FU as well as its deoxyriboside FUDR (Moertel and Reiemeier, 1969; Ansfield, 1975). Apart from another small patient series with rectal cancer undergoing preoperative radiochemotherapy (Dahl *et al*, 1994), to our knowledge, no other conclusive data are available about treatment tolerance as a potential prognostic marker for response despite the fact that in >80% of all patients receiving chemotherapy some kind of side effects (of course largely depending on which chemotherapy regimen is administered) will occur. Apart from use of inappropriate drug dosages in patients with impaired renal or hepatic function, presence of well-defined anticancer drug-specific risk factors such as DPD deficiency in case of FU-based chemotherapy (Diasio *et al*, 1988, 1998; Johnson *et al*, 1999; Milano *et al*, 1999), and to some extent gender (Stein *et al*, 1995; Sloan *et al*, 2002) and age >65 (Stein *et al*, 1995; Cascinu *et al*, 1996; Chiara *et al*, 1998; Popescu *et al*, 1999), the reason/mechanism is still unclear which predisposes patients to experience chemotherapy-related side effects.

As a result of uncertainties of a correlation between incidence rates and severity of side effects with the effectiveness of palliative chemotherapy in colorectal cancer, particularly as it concerns the novel more effective combination regimens, the present study was

*Correspondence: Dr B Schuell; E-mail: birgit.schuell@meduniwien.ac.at
Revised 1 August 2005; accepted 12 August 2005; published online 13 September 2005

performed. In an attempt to answer this potentially important basic question, we have analysed the toxicity, objective response and survival data of 303 patients with advanced colorectal cancer who were consecutively treated in four prospective oxaliplatin- or irinotecan-based combination chemotherapy studies.

PATIENTS AND METHODS

Patients

A total of 303 patients with advanced colorectal cancer, who were entered in four different multicentre first- or second-line chemotherapy protocols between 1998 and 2001 were analysed (Scheithauer *et al*, 2001a, b, c, 2003). All patients had histologically confirmed metastatic or locally advanced/recurrent adenocarcinomas with bidimensionally measurable disease (defined as presence of at least one index lesion capable of two-dimensional measurement by computed tomography (CT) scan. Additional eligibility criteria in all four trials included age between 19 and 75 years, a World Health Organisation (WHO) performance status of two or less, and adequate bone marrow reserve, as well as renal and hepatic function. Adjuvant 5-FU-based chemotherapy and/or radiation was allowed if it was completed ≥ 6 months before study entry. Within 2 weeks prior to initiating chemotherapy, all patients were assessed by physical examination, routine haematology and biochemistry analyses and CT-scans (or MRI) to define the extent of disease. Complete blood cell counts with platelet and differential counts were repeated weekly, and serum chemistries were determined at least once every course. All side effects, performance status, body weight, physical examination and subjective symptoms were recorded before each treatment course. Target lesions were reassessed by CT-scan or MRI every 8–12 weeks; objective response was evaluated according to WHO standard criteria.

Chemotherapy

The following six treatment regimens were administered to the 303 patients, who actively participated in four different trials: The first trial was a phase II study of front-line combination chemotherapy with raltitrexed 3 mg m^{-2} and oxaliplatin 130 mg m^{-2} both administered on day 1 every 3 weeks (Scheithauer *et al*, 2001a). In the second trial, the same dose regimen was administered to patients failing prior fluoropyrimidine/LV-based chemotherapy (Scheithauer *et al*, 2001b). The third trial was a randomised multicentre phase II trial of oxaliplatin plus irinotecan *vs* raltitrexed as first-line treatment with a crossover design upon progression (Scheithauer *et al*, 2001c): patients randomised to the combination arm, were treated with oxaliplatin 85 mg m^{-2} and irinotecan 175 mg m^{-2} every 2 weeks. Patients randomised to the raltitrexed-arm received a dose of 3 mg m^{-2} given on day 1 every 3 weeks. The last trial was also a randomised multicentre phase II study of two different schedules of capecitabine plus oxaliplatin as first-line treatment in advanced colorectal cancer (Scheithauer *et al*, 2003). Half of the patients received oxaliplatin 130 mg m^{-2} on day 1 plus capecitabine 2000 mg m^{-2} on days 1–14 every 3 weeks, in the other treatment arm patients received oxaliplatin 85 mg m^{-2} on days 1 and 14 combined with capecitabine 3500 mg m^{-2} day on days 1–7 and 14–21 every 4 weeks.

Side effects

Only chemotherapy-associated adverse reactions that occurred during the first 3 months of treatment were analysed. This was performed because (1) acute side effects generally occur within the first few courses, and (2) to avoid a possible selection bias by analysing fewer treatment courses in unresponsive patients. All except five patients received a minimum of 3 months chemotherapy

(98.3%). Out of this five patients, two had a progressive disease after 2 months, two experienced severe toxicities and one patient refused further treatment after the first cycle of chemotherapy. Oxaliplatin-associated sensory neuropathy, which is cumulative in its nature, was not included in this analysis. Adverse events were graded according to WHO standard toxicity criteria. All side effects were first analysed in total, and then subdivided into haematological and nonhaematological side effects, as well as severe (grade 3/4) and minor (grade ≤ 2) toxicities.

Statistical analyses

Continuous covariates are described with mean and s.d. in the case of normally distributed data and with median, minimum and maximum for skewed covariates. Categorical variables are described with frequencies and percentages. Differences between the two interesting groups (complete remission (CR), partial remission (PR) and stable disease (SD) *vs* progressive disease (PD)) in terms of number of side effects, number of tumour sites, gender, age and chemotherapy regimen were examined with univariate and multiple logistic regression models always modelling the probability of experiencing CR; PR or SD *vs* PD. Number of side effects and all other categorisations of side effects (haematological and nonhaematological, WHO grade > 2 and WHO grade ≤ 2) as well as number of tumour sites are modelled as linear factors in logistic regression analyses, implying a constant odds ratio between consecutive variable values. All other scrutinised factors are taken into account as categorical variables, whereas categories male in case of gender, regimen, irinotecan plus oxaliplatin for chemotherapy, and age up to 65 years served as reference categories. All analyses were carried out using the statistical software package SAS (version 8.02, SAS Institute, Cary, NC, USA), *P*-values are two-sided and *P* < 0.05 was considered statistically significant.

RESULTS

A total of 303 patients (190 men and 113 women) who received first- or second-line chemotherapy for advanced colorectal cancer in four different clinical trials between 1998 and 2001 were included in this analysis. The pretreatment characteristics of the study population(s) which were essentially similar across the four trials are summarised in Table 1. Patients were divided into two groups: the first group (group I) had a positive treatment effect, which included complete response, partial response and SD; the second group (group II) was rated progressive during chemotherapy. As shown in Table 2, out of the 303 patients, 245 were categorised in group I, and 58 patients in group II.

Out of the entire study population only six patients had no side effects, and 297 patients had experienced at least one adverse event. The median number of side effects (haematological and nonhaematological, all WHO grades) in group I was 6 (range, 0–12) compared to 4 (range, 0–13) in case of treatment failure.

The chance to have CR, PR or SD, in fact, was noted to be increasing with an increasing number of side effects (OR = 1.342, *P* < 0.0001). Specifically, one additional side effect increases the odds of no PD by 34% (Table 3). The increasing incidence of side effects from PD to complete response showed a highly significant correlation (*P* < 0.0001) without overlapping 95% confidence intervals from PD to SD onwards.

The observation of an inverse correlation of disease control with treatment tolerance was confirmed when side effects were analysed according to severity: in patients experiencing minor toxicities (WHO grade ≤ 2 ; *n* = 284), the median number of side effects was 3 (range, 0–10) in case of treatment efficacy, and 2 (range, 0–7) in nonresponders. The corresponding odds ratio in a univariate logistic regression model was statistically significant (OR = 1.319,

Table 1 Selected pretreatment characteristics of the analyzed study population

	No of patients, n (%)	
Gender		
Male	190	63%
Female	113	37%
Age		
Median	66.5	
Range	38–79	
WHO performance status		
0	138	46%
1	123	40%
2	42	14%
Primary site		
Colon	193	64%
Rectum	110	36%
Sites of metastases		
Liver	218	
Lung	75	
Abdominopelvic mass	99	
Other	118	
Number of metastatic sites		
Single	116	38%
Multiple	187	62%

$P=0.002$). In patients with severe toxicities (WHO grade >2 ; $n=45$), responders had a median number of 2 side effects (range, 0–9), whereas nonresponder only suffered from 1 (range, 0–7; OR = 1.372, $P=0.0006$).

We then divided treatment-associated side effects into haematological and nonhaematological. In all, 252 patients (83%) had haematological side effects and 289 patients (95%) had at least one nonhaematological side effect. Patients with at least SD had significantly more ($P=0.0005$) haematological side effects (median 3, range 0–4) than patients with tumour progression (median 1, range 0–4). One haematological side effect more implies a 1.472 times higher odds to experience at least SD ($P=0.0005$). The most common nonhaematological side effects (apart from the cumulative oxaliplatin-associated peripheral sensory neuropathy, which was excluded from this analysis), were diarrhoea, transient elevation of liver functional parameters and nausea/emesis. The median number of nonhaematological side effects of a responder were 4 (range, 0–9) vs 2 (range, 0–9) in case of treatment failure, a difference that was again statistically significant ($P<0.0001$, OR: 1.40).

We subsequently analysed the correlation of the response status and side effects according to the various treatment regimens (Table 2). Of the patients (87%) who received chemotherapy with oxaliplatin plus capecitabine were in group I (80 patients), and 13% (12 patients) had PD (group II). The median number of side effects in both groups was 6 (group I: range, 0–11; group II: range, 2–8); no difference was noted between the two groups for this particular combination regimen ($P=0.578$, OR: 1.07). In all, 82 out

Table 2 Descriptive statistics

	Response to Treatment		Total (n = 303)
	CR+PR+SD vs (n = 245)	PD (n = 58)	
Gender			
Female	88 (77.9%)	25 (22.1%)	113
Male	157 (82.6%)	33 (17.4%)	190
Age			
Up to 65 years	127 (79.4%)	33 (20.6%)	160
Older than 65 years	118 (82.5%)	25 (17.5%)	143
Mean (\pm s.d.)	63.12 (\pm 9.70)	62.05 (\pm 9.93)	62.92 (\pm 9.74)
Min–max	37–79	38–77	37–79
Number of side effects			
Median (min–max)	6 (0–12)	4 (0–13)	6 (0–13)
Number of side effects WHO-grade <2 (n = 284)			
Median (min–max)	3 (0–10)	2 (0–7)	3 (0–10)
Number of side effects WHO-grade ≥ 2 (n = 45)			
Median (min–max)	2 (0–9)	1 (0–7)	2 (0–9)
Number of haematological side effects (n = 252)			
Median (min–max)	3 (0–4)	1 (0–4)	2 (0–4)
Number of non-haematological side effects (n = 289)			
Median (min–max)	4 (0–9)	2 (0–9)	3 (0–9)
Number of tumour sites			
Median (min–max)	2 (1–5)	2 (1–5)	2 (1–5)
Chemotherapy regimen			
Capecitabine plus oxaliplatin	80 (87.0%)	12 (13.0%)	92
Raltitrexed plus oxaliplatin	82 (91.1%)	8 (8.9%)	90
Irinotecan plus oxaliplatin	53 (82.8%)	11 (17.2%)	64
Raltitrexed	30 (52.6%)	27 (47.4%)	57

CR = complete response, PR = partial response, SD = stable disease, PD = progressive disease according to the World Health Organisation standard criteria.

of 90 patients (91%) who received raltitrexed plus oxaliplatin had an objective response or SD, and eight patients had PD. The median number of side effects in group I was 5 compared with 3.5 in group II, again a difference that did not reach the level of statistical significance ($P=0.0623$, OR: 1.46). A total of 64 patients with metastatic colorectal cancer were treated with irinotecan plus oxaliplatin as first-line therapy. Of these patients, 83% were categorised in group I, and 11 patients in group II. In group I the median number of side effects was 8 as opposed to group II, where patients experienced a median number of 6 side effects ($P=0.028$, OR: 1.4). In all, 57 patients received raltitrexed as first-line chemotherapy. The median number of side effects was 5 in group I ($n=30$), and in group II ($n=27$) the median number of side effects was 3 ($P=0.0038$, OR: 1.46).

The median survival of all patients who had at least a stable disease was 19 months compared to 5.5 months in those with PD. When median survival was analysed according to the number of adverse events, it was 10 (95%CI, 3–7), 16 (14–18), and 18 (16–20) months in case of 0–1, 2–5 and ≥ 6 AEs, respectively (log-rank test: $P=0.01$; Figure 1).

As indicated in Table 4, a multiple logistic regression model was used to assess the partial effects of number of side effects, number of tumour sites, gender, age and chemotherapy regimen in terms of treatment benefit. The number of side effects ($P<0.0001$), the number of tumour sites ($P=0.003$), and treatment with raltitrexed plus oxaliplatin ($P=0.0081$) turned out to be significant prognostic factors. As it concerns gender and age, no statistical

difference was seen neither in the univariate nor in the multivariate analysis.

DISCUSSION

Administration of the same dose regimen of a specific anticancer drug or drug combination to a population of patients is likely to result in considerable variations of toxicities ranging from no side effect to potentially lethal events (Kuilenburg *et al*, 2001; Rothenberg *et al*, 2001). In our univariate and multivariable analyses the number of side effects associated with modern combination chemotherapy in advanced colorectal cancer turned out to be a significant independent prognostic factor; one additional adverse reaction, in fact, seems to increase the odds of response or SD by 34%. Similarly, occurrence of side effects was found to correspond with an improved overall survival. The correlation between toxicity and therapeutic benefit was noted irrespective of the degree and type of toxicity, that is, nonhaematological vs haematological. The latter observation seems important because haematological side effects are objectively measured toxicity parameters, which are not subject to a potential reporting bias. Variations of the correlation between toxicity and therapeutic benefit between the four treatment regimens are most likely to be related to the rather small study patient populations analysed, and especially the low rate of treatment failures in case of combination chemotherapy; this hypothesis is supported by the largest correlation of adverse events and response noted in the (least effective) raltitrexed control group in one of the studies.

It seems noteworthy that the correlation between toxicity and therapeutic benefit was not affected by gender and age. Differences in reporting subjective symptoms between men and women have been described in an analysis of 2448 patients who received FU-based chemotherapy for colorectal cancer (Sloan *et al*, 2002). Women experience toxicity more frequently and with more severity than men, whereby the gender variation could be related to the methylenetetrahydrofolate reductase C677T polymorphism (Van Rijnsoever *et al*, 2002), and/or the different levels of dihydropyrimidine dehydrogenase (Diasio, 1998; Milano *et al*, 1999). The current analysis also shows no difference between sexes in objective response rate or survival. The same is true for various otherwise rather divergent data about age and toxicity (Stein *et al*, 1995; Cascinu *et al*, 1996; Chiara *et al*, 1998; Popescu *et al*, 1999). Response and survival were not affected in these studies as they were in the current study, regardless of a difference in tolerance of 5-FU-based therapy between younger and older patients.

The results of our findings with the use of modern combination chemotherapy in advanced colorectal cancer are in agreement with an analysis of the relationship between the toxicity of FU ($n=176$) and FUDR monotherapy ($n=203$) to objective response rates in a comparable study population, which was published in 1969 (Moertel and Reiemeier, 1969). Fluoropyrimidine treatment to the point of mild or moderate leukocytopenia (1500–4500) resulted in a response rate of 22.7 vs 8.7 and 15% in case of no

Table 3 Univariate logistic regression analyses

	Odds ratio	95% CI	P
Number of side effects	1.342	1.186–1.519	<0.0001
Number of side effects WHO grade 2	1.319	1.107–1.572	0.0020
Number of side effects WHO grade >2	1.372	1.146–1.642	0.0006
Number of haematological side effects	1.472	1.183–1.832	0.0005
Number of nonhaematological side effects	1.398	1.182–1.654	<0.0001
Gender (reference group: male)	0.740	0.414–1.323	0.3099
Age (reference group: up to 65 years)	1.226	0.689–2.184	0.4880
Number of tumour sites	0.595	0.428–0.828	0.0021




Figure 1 Median survival according to number of adverse events.

Table 4 Multiple logistic regression analysis

	Odds Ratio	95% CI	P
Number of side effects	1.349	1.167–1.559	<0.0001
Gender (reference group: male)	0.715	0.357–1.433	0.3439
Age (reference group: up to 65 years)	1.556	0.799–3.033	0.1939
Number of tumour sites	0.570	0.394–0.826	0.0030
<i>Chemotherapy regimen</i>			
Irinotecan plus oxaliplatin	1.000		
Raltitrexed plus oxaliplatin	4.271	1.457–12.521	0.0081
Raltitrexed	0.526	0.198–1.395	0.1966
Capecitabine plus oxaliplatin	1.697	0.654–4.406	0.2770

(>4500) or more severe (<1500) haematotoxicity. A randomised evaluation of a planned subtoxic FU dosage conducted by the Central Oncology Group has also demonstrated a significant inferiority in response rate when compared to a dose-producing mild to moderate toxicity (Ansfield, 1975). A third analysis we could identify in terms of correlating side effects with tumour response was published by Dahl *et al* (1994). The investigators described 159 patients who were treated with preoperative radiotherapy (31.5 Gy in 18 fractions) for potentially resectable rectal adenocarcinoma. Patients were examined for a possible relationship between bowel toxicity manifested as diarrhoea and tumour size in the operative specimen as well as recurrence rate. Patients who required drugs for diarrhoea had significantly smaller tumours at surgery (2.5 vs 3.5 cm). Furthermore, patients without significant radiation-induced diarrhoea had more recurrences (37.5 vs 14.3%). The disease-specific survival rate was also significantly better ($P=0.02$) at 1.5 and 10 years in patients with diarrhoea WHO grades 3 and 4. It was therefore concluded that a correlation between bowel sensitivity and tumour sensitivity to radiation might exist.

REFERENCES

- Ansfield FJ (1975) A randomized phase III study of four dosage regimens of 5-fluorouracil. *Proc Am Assoc Cancer Res* 16: 224 (abstract)
- Blum J, Jones S, Buzdar A (2001) Capecitabine (Xeloda) in 162 patients with paclitaxel-pretreated MBC: updated results and analyses of dose modification (abstract 693). *Eur J Cancer* 37(Suppl 6): S190 (abstract)
- Cascinu S, Del Ferro E, Catalano G (1996) Toxicity and therapeutic response to chemotherapy in patients aged 70 years or older with advanced cancer. *Am J Clin Oncol* 19: 371–374
- Cassidy J, Twelves C, Van Cutsem E, Hoff P, Bajetta E, Boyer M, Bugat R, Burger U, Garin A, Graeven U, McKendric J, Maroun J, Marshall J, Osterwalder B, Perez-Manga G, Rosso R, Rougier P, Schilsky RL (2002) First-line oral capecitabine therapy in metastatic colorectal cancer: a favorable safety profile compared with iv 5-fluorouracil/leucovorin. *Ann Oncol* 13: 566–575
- Chiara S, Nobile MT, Vincenti M, Lionetto R, Gozza A, Barzacchi MC, Sanguineti O, Repetto L, Rosso R (1998) Advanced colorectal cancer in the elderly: results of consecutive trials with 5-fluorouracil-based chemotherapy. *Cancer Chemother Pharmacol* 42: 336–340
- Dahl O, Horn A, Mella O (1994) Do acute side-effects during radiotherapy predict tumor response in rectal carcinoma? *Acta Oncol* 33: 409–413
- Diasio RB, Beavers TL, Carpenter JT (1988) Familial deficiency of dihydropyrimidine dehydrogenase. Biochemical basis for familial pyrimidinemia and severe 5-fluorouracil-induced toxicity. *J Clin Invest* 81: 47–51
- Diasio RB (1998) The role of dihydropyrimidine dehydrogenase (DPD) modulation in 5-FU pharmacology. *Oncology (Huntingt)* 12: 23–27
- Grothey A, Sargent D, Goldberg R, Schmoll HJ (2004) Survival of patients with advanced colorectal cancer improves with the availability of fluorouracil, leucovorin, irinotecan, and oxaliplatin in the course of treatment. *J Clin Oncol* 22: 1209–1214
- Johnson MR, Hageboutros A, Wang L, High L, Smith JB, Diasio RB (1999) Life-threatening toxicity in a dihydropyrimidine dehydrogenase-deficient patient after treatment with topical 5-fluorouracil. *Clin Cancer Res* 5: 2006–2011
- Köhne CH, Cunningham D, Di Costanzo F, Glimelius B, Blijham G, Aranda E, Scheithauer W, Rougier P, Palmer M, Wils J, Baron B, Pignatti F, Schoffski P, Micheel S, Hecker H (2002) Clinical determinants of survival in patients with 5-fluorouracil based treatment for metastatic colorectal cancer: results of a multivariate analyses of 3825 patients. *Ann Oncol* 13: 308–317
- Kuilenburg AB, Muller EW, Haasjes J, Meinsma R, Zoetekouw L, Waterham HR (2001) Lethal outcome of a patient with a complete dihydropyrimidine dehydrogenase (DPD) deficiency after administration of 5-fluorouracil: frequency of the common IVS14 + 1G>A mutation causing DPD deficiency. *Clin Cancer Res* 7: 1149–1153
- Landis SH, Murray T, Bolden S (1999) Cancer statistics 1999. *CA Cancer J Clin* 49: 8–13
- Midgley R, Kerr D (1999) Colorectal cancer. *Lancet* 353: 391–399
- Milano G, Etienne M, Pierreite V, Barberi-Heyob M, Deporte-Fety R, Renee N (1999) Dihydropyrimidine dehydrogenase deficiency and fluorouracil-related toxicity. *Br J Cancer* 79: 627–630
- Moertel CG, Reiemeier RJ (1969) *Advanced Gastrointestinal Cancer. Clinical Management and Chemotherapy*. New York: Harper & Row Pubs Inc.
- Popescu RA, Norman A, Ross PJ, Parikh B, Cunninham D (1999) Adjuvant or palliative chemotherapy for colorectal cancer in patients 70 years or older. *J Clin Oncol* 17: 2412–2418
- Rothenberg ML, Meropol NJ, Poplin EA, van Cutsem E, Wadler S (2001) Mortality associated with irinotecan plus bolus fluorouracil/leucovorin: summary findings of an independent panel. *J Clin Oncol* 19: 3801–3807
- Scheithauer W, Kornek GV, Raderer M, Schuell B, Schmid K, Kovats E, Schneeweiss B, Lang F, Lenauer A, Depisch D (2003) Randomized multicenter phase II trial of two different schedules of capecitabine plus oxaliplatin as first-line treatment in advanced colorectal cancer. *J Clin Oncol* 21: 1307–1312
- Scheithauer W, Kornek GV, Raderer M, Ulrich-Pur H, Fiebigler W, Gedlicka C, Schull B, Brugger S, Schneeweiss B, Lang F, Lenauer A, Depisch D (2001a) Randomized multicenter phase II trial of oxaliplatin plus irinotecan versus raltitrexed as first-line treatment in advanced colorectal cancer. *J Clin Oncol* 20: 165–172
- Scheithauer W, Kornek GV, Schuell B, Ulrich-Pur H, Penz M, Raderer M, Lang F, Schneeweiss B, Lenauer A, Depisch D (2001b) Second-line treatment with oxaliplatin+raltitrexed in patients with advanced colorectal cancer failing fluoropyrimidine/leucovorin-based chemotherapy. *Ann Oncol* 12: 709–714
- Scheithauer W, Kornek GV, Ulrich-Pur H, Penz M, Raderer M, Salek T, Haider K, Kwasny W, Depisch D (2001c) Oxaliplatin plus raltitrexed in patients with advanced colorectal carcinoma. *Cancer* 91: 1264–1271
- Scheithauer W, Rosen H, Kornek GV, Sebesta C, Depisch D (1993) Randomized comparison of combination chemotherapy plus supportive care with supportive care alone in patients with metastatic colorectal cancer. *Br Med J* 306: 752–756
- Sloan JA, Goldberg RM, Sargent DJ, Vargas-Chanes D, Nair S, Cha SS, Novotny PJ, Poon MA, O'Connell MJ, Loprinzi CL (2002) Women experience greater toxicity with fluorouracil-based chemotherapy for colorectal cancer. *J Clin Oncol* 20: 1491–1498
- Stein BN, Petrelli NJ, Douglass HO, Driscoll DL, Arcangeli G, Meropoe NJ (1995) Age and sex are independent predictors of 5-fluorouracil toxicity. Analysis of a large scale phase III trial. *Cancer* 74: 11–17
- Van Rijnsoever M, Griew F, Elsaleh H, Joseph D, Iacopetta B (2002) Characterization of colorectal cancers showing hypermethylation at multiple CpG islands. *Gut* 51: 797–802
- Williams NS, Northover JMA, Arnott SJ (1995) Colorectal tumors. In *Oxford Textbook of Oncology*, In Peckham M, Pinedo H, Veronesi U (eds) Oxford, UK: Oxford University Press, pp 1133–1168