

Full-length article

MEK inhibitor PD98059 acutely inhibits synchronized spontaneous Ca²⁺ oscillations in cultured hippocampal networksYan-fang RUI¹, Zhao-hui SUN², Jia-ping GU¹, Zhong-hua SHENG¹, Xiang-ping HE¹, Zuo-ping XIE^{1,3}

¹Department of Biological Science and Biotechnology, State Key Laboratory of Biomembrane and Membrane Biotechnology, Tsinghua University, Beijing 100084, China; ²Institute of Neurological Disorders, Life Science and Medical Department, Tsinghua University, Beijing 100084, China

Key words

PD98059; MEK; calcium oscillation

³ Correspondence to Prof Zuo-ping XIE.
Phn 86-10-6278-8677.
Fax 86-10-6277-2271.
E-mail zuopingx@mail.tsinghua.edu.cn

Received 2006-04-20

Accepted 2006-05-19

doi: 10.1111/j.1745-7254.2006.00387.x

Abstract

Aim: To investigate the changes in synchronized spontaneous Ca²⁺ oscillations induced by mitogen-activated protein kinase kinase (MEK) inhibitor PD98059 at different concentrations in cultured hippocampal network. **Methods:** Hippocampal neurons in culture for 1–2 weeks were used for this study. Spontaneous synaptic activities of these hippocampal neurons were examined by Ca²⁺ imaging using calcium-sensitive dye. MEK inhibitor PD98059 (10, 30, and 60 μmol/L) and SB202474 (10 and 60 μmol/L), a negative control for mitogen-activated protein kinase (MAPK) cascade study, were applied to the cells under the microscope while imaging was taking place. **Results:** PD98059 at a lower concentration of 10 μmol/L had little effect on the Ca²⁺ oscillation. At the higher concentration of 30 μmol/L, 5 min after application of PD98059, the spike frequency was decreased to 25.38%±7.40% (mean±SEM, n=16, P<0.01 vs medium control) of that of the control period. At an even higher concentration of 60 μmol/L, 5 min after application of PD98059, the spike frequency was decreased to 14.53%±5.34% (mean±SEM, n=16, P<0.01 vs medium control) of that of the control period. The spike amplitude underwent a corresponding decrease. However, the negative control SB202474 at concentrations of 10 and 60 μmol/L had little inhibition effect on the Ca²⁺ oscillation. **Conclusion:** These results indicate that PD98059 inhibits synchronized spontaneous Ca²⁺ oscillation through inhibition of MEK, which hints that the MAPK cascade is required to maintain synchronized spontaneous Ca²⁺ oscillation.

Introduction

The mitogen-activated protein kinase (MAPK) cascade is a ubiquitous serine/threonine kinase cascade that has been classically studied as a critical biochemical pathway involved in cell proliferation and differentiation^[1]. MAPKs constitute a superfamily of three related kinases that are activated by various extracellular stimuli including the extracellular signal regulated kinases (ERKs), the Jun N-terminal kinases (JNKs), and p38 kinases (p38)^[2,3]. ERK, JNK, and p38 can all be activated by a variety of stimuli, but these kinases are differentially affected by certain signals. For example, ERKs are most highly activated in response to mitogenic stimulation, whereas JNKs and p38 show greater activation in response

to cellular stress^[4,5]. The pathway leading to ERK activation by growth factors and other mitogens has been studied extensively. The first step involves activation of membrane-associated tyrosine kinases, followed by the sequential activation of Ras and Raf. Raf then phosphorylates the mitogen-activated protein kinase kinase (MEK), which in turn activates ERK^[6,7]. Although this cascade is mainly studied in mitotic cell regulation, its components are actually most abundantly expressed in postmitotic neurons of the developed nervous system^[8]. The hippocampus region, which is commonly used as a model to study synaptic plasticity, has highly expressed ERK^[9,10]. What are the physiologic roles of this cascade in mature neurons? PD98059, a specific inhibitor of MEK, the enzyme that activates ERK^[11], has been

shown to block induction of LTP in area CA1 of the hippocampus^[12] and attenuates multiple forms of synaptic plasticity in rat dentate gyrus *in vitro*^[13]. However, the regulation mechanism or physiological role of this cascade in the activity-dependent synaptic connections between neurons is not clear.

Calcium plays an important role in regulating a great variety of neuronal processes, especially in the transmitter release and synaptic connection. Oscillations in cytoplasmic calcium have been observed in a wide variety of neuronal cell types including cortical and hippocampal neurons^[14–16]. In primary cultured hippocampal neurons, after one week in culture, networks of interconnected neurons are formed. At approximately 9 d *in vitro*, some networks show spontaneous synchronized Ca^{2+} oscillations^[15,17]. These oscillations are believed to encode information in neural circuits^[18,19] and might play an important role during physiological or pathological events^[20,21]. Many studies have implied that the MAPK cascade might participate in $[\text{Ca}^{2+}]_i$ regulation^[22–24]. Here we used PD98059, a commercially available inhibitor of MEK, and SB202474, a negative control, to explore whether this cascade participates in the regulation of spontaneous synchronized Ca^{2+} oscillations.

Materials and methods

Drugs Dulbecco's modified Eagle's medium (DMEM) media, neurobasal medium, fetal bovine serum, B27 supplements, 0.25% trypsin-EDTA, and poly-*D*-lysine for cell culture were from Invitrogen (Carlsbad, CA, USA). Equine serum and *L*-glutamine were from Hyclone (Logan, UT, USA). PD98059 and SB202474 were purchased from Calbiochem (La Jolla, CA, USA) and were dissolved in dimethyl sulfoxide. Fluo-4-AM was from Molecular Probes (Eugene, OR, USA). Other reagents were purchased from Sigma (St Louis, MO, USA).

Hippocampal cell culture and experiment Hippocampal neurons from embryonic rats (E18) were obtained according to the method previously described^[25]. In brief, hippocampal tissues from 18-d-old fetal rats were dissected and treated with 0.25% trypsin in Ca^{2+} - Mg^{2+} -free HBSS at 37 °C for 15 min; they were then dissociated by trituration with a glass Pasteur pipette and plated in 35 mm culture dishes with glass bottoms (MatTek, Ashland, MA) for culture and subsequent microscopy. The glass surface in each dish (~15 mm diameter) was pretreated with poly-*D*-lysine for 2 h (500 $\mu\text{g}/\text{mL}$ in borate buffer), washed three times, and air-dried before cell plating. Approximately 75 000 cells were plated in the glass area of each dish in DMEM containing 5% fetal bovine se-

rum and 5% horse serum. On the second day after plating, the culture medium was replaced by serum-free Neurobasal medium containing 2% B27 supplement and 500 $\mu\text{mol}/\text{L}$ glutamine for reduced glial growth. Cells were maintained in a CO_2 incubator at 37 °C, and one-half volume of the culture medium was replaced with fresh Neurobasal medium every 3 d. The experiments were carried out on cultures after 7 d.

Ca^{2+} imaging Hippocampal cells were loaded with 4 $\mu\text{mol}/\text{L}$ Fluo-4-AM in Krebs-Ringer's saline (recording solution) (150 mmol/L NaCl, 5 mmol/L KCl, 2 mmol/L CaCl_2 , 1 mmol/L MgCl_2 , 10 mmol/L glucose, and 10 mmol/L HEPES, pH 7.4)^[19] at 37 °C for 30 min, followed by three washes and a 15-min incubation period for further de-esterification of Fluo-4-AM before imaging. Cells grown on the glass bottom in 35 mm dishes were directly imaged on a Nikon (Tokyo, Japan) TE300 inverted microscope using a 40 \times numerical aperture, 1.30 oil immersion Plan Fluor objective. A Lambda DG-4 highspeed wavelength switcher (Sutter Instruments, Novato, CA) was used for Fluo-4 excitation at 480 nm, and a cooled CCD camera (CoolSnap FX; Roper Scientific, Princeton, NJ) was used for image acquisition. MetaFluor imaging software (Universal Imaging, Downingtown, PA) was used for hardware control, image acquisition, and image analysis. The time-lapse recording of Ca^{2+} signals in hippocampal neurons was carried out for a 2-min control period before and a 6-min period after the application of different chemicals. The sampling rate was one frame every 2 s. The exposure time was 50 ms when CCD binning of 4 \times 4 was used.

Quantitative analysis of synchronized Ca^{2+} spikes Quantitative measurements of changes in intracellular Ca^{2+} concentrations ($[\text{Ca}^{2+}]_i$) were done by obtaining the average Fluo-4 fluorescence intensity of a 3 \times 3 pixel analysis box placed at the center of the cell body; the intensity values were then subtracted from the average background intensity measured in cell-free regions. Changes of $[\text{Ca}^{2+}]_i$ in each cell were then represented by the changes of relative Fluo-4 fluorescence ($\Delta F/F_0$) where F_0 was the baseline intensity obtained from the 2 min control period. Ca^{2+} spikes were defined as rapid elevation of $\Delta F/F_0$ equal to or >20%. Under our imaging settings, fields of 3–10 neurons were typically recorded and subsequently analyzed. To determine the frequency and amplitude of Ca^{2+} spikes, we counted the number of Ca^{2+} spikes and the average amplitude of these spikes over a 2-min period of the recording as a defined time point. As a result, the 2-min control period yielded only one frequency and one amplitude value, whereas the experiment period (6 min) resulted in three frequency and amplitude values at different time points after bath application of a specific molecule. To the changes in the spike frequency and

amplitude, these three frequency and amplitude values after the drug application were normalized to the control frequency or amplitude values respectively and expressed as percentages, with a value of 100% indicating no change. We quantified and examined the changes in the spike frequency and amplitude through the entire 6 min period after bath application. In order to assess the baseline changes after drug application, we calculated the average $\Delta F/F_0$ values of the rock bottom of each spike.

Bath application of different drugs To prevent adverse effects of high concentrations of drugs, a 2×working concentration of the drug was made in Krebs-Ringer’s solution and was applied to the cells to achieve the desired final concentration through 1:1 dilution (v/v). Specifically, we first recorded Ca^{2+} activities for a 2-min control period in 1 mL of Krebs-Ringer’s solution, removed 0.5 mL from the bath, added 0.5 mL of the 2×solution, and subsequently recorded for 6 min to examine the effects on spontaneous Ca^{2+} spikes. For the control, we simply carried out the same procedure to apply Krebs-Ringer’s solution to determine that there was no artifact of this application method.

Statistical analysis Data from at least three dishes from different batches of cultures were pooled together and analyzed for statistically significant differences using the paired Student’s *t*-test. Compiled data are expressed and graphed as mean±SEM, with *n* denoting the number of neurons studied for each treatment. Differences were considered significant if a *P* value was <0.05.

Results

Synchronized spontaneous Ca^{2+} spikes and the mechanisms in cultured hippocampal networks We prepared low-density hippocampal neurons culture as described by Banker *et al*^[25]. After at least one week in culture, many hippocampal neurons formed local networks which usually contained

3–10 neurons (Figure 1A, left panel). Spontaneous synaptic activities of these neurons were examined by Ca^{2+} imaging using the calcium-sensitive dye Fluo-4^[26] (Figure 1A, right panel). We observed periodic, spontaneous spike elevations of $[Ca^{2+}]_i$ and these spikes appeared to be primarily synchronized among the local group of cells (Figure 1B) without removing or reducing Mg^{2+} in medium.

The mechanisms underlying the synchronized spontaneous Ca^{2+} spikes in hippocampal networks have been studied extensively, but the results from different published reports are confusing. This may be caused by the variety of preparations used for experiments. For example, Leinekugel *et al* reported that the synchronized spontaneous Ca^{2+} spikes were mediated by the synergistic excitatory actions of gamma-aminobutyric acid ($GABA_A$) and *N*-methyl-*D*-aspartate (NMDA) receptors in the neonatal hippocampus^[27], whereas Tanaka *et al* reported that the oscillation of Ca^{2+} was mainly mediated by non-NMDA-type glutamatergic transmission^[15]. To confirm the Ca^{2+} spikes observed in our culture were driven by particular receptors, we applied different antagonists of these receptors to our cultures. We found that a non-selective antagonist of NMDA and alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA)/kainate receptors kynurenic acid 1 mmol/L completely blocked the spikes immediately after application (Figure 2A). The NMDA receptor antagonist APV at 50 μ mol/L only partially inhibited the spike amplitude (Figure 2B), whereas AMPA/kainate receptors antagonist 6,7-dinitroquinoxaline-2,3-dione (DNQX) at 20 μ mol/L completely and immediately blocked the spikes (Figure 2B, 2C). The addition of the $GABA_A$ receptor antagonist bicuculline had a mixed effect on the Ca^{2+} oscillations, which caused an increase in amplitude but a decrease in frequency (Figure 2D). Subsequently adding kynurenic acid or DNQX completely blocked the spikes (Figure 2D–F). These findings suggest that the oscillations we observed are similar to those




Figure 1. Synchronized spontaneous Ca^{2+} spikes in cultured hippocampal neurons. (A) The DIC image on the left shows five interconnected hippocampal neurons. The fluorescent image on the right shows the same region of hippocampal neurons with the DIC image loaded with Fluo-4. Scale bar, 40 μ m. (B) Traces depict synchronized spontaneous Ca^{2+} transients in these five neurons.

observed by Tanaka *et al*, which were mainly mediated by non-NMDA-type glutamatergic transmission.

PD98059 acutely inhibited synchronized spontaneous Ca²⁺ spikes at a higher concentration To test whether the MAPK cascade is required to maintain Ca²⁺ oscillations, we used the MEK inhibitor PD98059 to block MAPK activation in hippocampal neurons. As reported by Dudley *et al*, PD98059 exerts its inhibition effect on MEK at concentrations from 1 to 100 $\mu\text{mol/L}$, with the IC₅₀ value at approximately 10 $\mu\text{mol/L}$ ^[28]. We chose 10, 30, and 60 $\mu\text{mol/L}$ concentrations to test the effect on Ca²⁺ oscillations. We found that PD98059

at 10 $\mu\text{mol/L}$ had no significant effect on Ca²⁺ spike frequency and only slightly decreased the Ca²⁺ spike amplitude (Figure 3A,3D,3E). PD98059 30 $\mu\text{mol/L}$ significantly inhibited the Ca²⁺ spikes immediately after application (Figure 3B). Six minutes after application, the spike frequency was decreased to 25.38% \pm 7.40% (mean \pm SEM, $n=16$) of that of the control period (Figure 3D) and the spike amplitude was 25.16% \pm 6.99% ($n=16$) of that of the control period (Figure 3E). Application of 60 $\mu\text{mol/L}$ PD98059 caused a more rapid and severe inhibition of the Ca²⁺ spikes (Figure 3C). Six minutes after application the spike frequency was decreased to 14.53% \pm




Figure 2. Role of glutamatergic and GABAergic neurons in calcium oscillation. Traces show synchronized Ca²⁺ spikes in 4–6 neurons randomly selected from a group of synchronically firing cells. The time gap indicated by arrows is approximately 30 s, during which time different agonists were added. (A) 1 mmol/L kynurenic acid; (B) 50 $\mu\text{mol/L}$ APV for 2 min, followed by 20 $\mu\text{mol/L}$ DNQX; (C) 20 $\mu\text{mol/L}$ DNQX for 2 min followed by 50 $\mu\text{mol/L}$ APV; (D) 50 $\mu\text{mol/L}$ bicuculline for 2 min followed by 1 mmol/L kynurenic acid; (E) 50 $\mu\text{mol/L}$ bicuculline for 2 min followed by 20 $\mu\text{mol/L}$ DNQX; and (F) 50 $\mu\text{mol/L}$ bicuculline for 2 min followed by 50 $\mu\text{mol/L}$ APV, then 20 $\mu\text{mol/L}$ DNQX. Each response was repeated at least three times.




Figure 3. Acute inhibition of the synchronized Ca^{2+} oscillations by PD98059. (A–C) Traces show synchronized Ca^{2+} spikes in several neurons randomly selected from a group of synchronically firing cells. The three parts represent Ca^{2+} spikes in these cells under control conditions, after application of control medium, and after bath application of 10 $\mu\text{mol/L}$ (A), 30 $\mu\text{mol/L}$ (B) and 60 $\mu\text{mol/L}$ (C) PD98059. Changes in the frequency (D) and mean amplitude (E) of the synchronized Ca^{2+} spikes every 2 min after bath application of medium (control) or different concentrations of PD98059. Data are presented as the mean \pm SEM. $n=16$ neurons studied for each trial. ^b $P<0.05$, ^c $P<0.01$ vs control.

5.34% ($n=16$) of that of the control period (Figure 3D) and the spike amplitude was 31.81% \pm 6.27% ($n=16$) of that of the control period (Figure 3E). No significant difference was found with the control period, indicating that no artifact was produced by the bath application method. Overall, these results demonstrate that PD98059 rapidly inhibits the Ca^{2+} spikes in a dose-dependent manner.

The effect of SB202474 on synchronized spontaneous Ca^{2+} spikes SB202474 is an inactive structural analog of PD98059. We also tested its effect on the Ca^{2+} spikes to confirm whether or not the inhibition effect of PD98059 on Ca^{2+} spikes was through inhibition of the MAPK cascade. We found that 10 $\mu\text{mol/L}$ SB202474 had no inhibition effect on the Ca^{2+} oscillations frequency, but it had a small enhancement effect on the frequency immediately after application. As a consequence of the frequency increase, the amplitude

of the spikes decreased (Figure 4A,4C,4D). Application of 60 $\mu\text{mol/L}$ SB202474 had no significant effect on frequency or amplitude of the Ca^{2+} spikes (Figure 4B,4C,4D).

Discussion

We prepared the hippocampal culture based mainly on the method described by Banker *et al*[25]. We used serum-free Neurobasal medium with B27 supplement to reduce glia cell growth and increase neuron survival. As described previously, in Neurobasal/B27 medium, glia growth is reduced to less than 0.5%, resulting in a nearly pure population of neurons[29]. We used the microtubule-associated protein (MAP2) and 4',6-diamidino-2-phenylindole, dihydrochloride (DAPI) double-staining to detect the neuron proportion in our culture. We found that there were




Figure 4. SB202474 had no effect on synchronized Ca^{2+} oscillations. (A, B) Traces show synchronized Ca^{2+} spikes in several neurons randomly selected from a group of synchronically firing cells. The three parts represent Ca^{2+} spikes in these cells under control conditions, after application of control medium, and after bath application of 10 $\mu\text{mol/L}$ (A) and 60 $\mu\text{mol/L}$ (B) SB202474. Changes in the frequency (C) and mean amplitude (D) of the synchronized Ca^{2+} spikes every 2 min after bath application of medium (control) or different concentrations of SB202474. Data are presented as the mean \pm SEM. $n=16$ neurons studied for each trial. ^b $P<0.05$, ^c $P<0.01$ vs control.

more than 85% neurons in our culture, and the other cells were glia cells including astrocytes and oligocytes. In this culture, glia cells were spread beneath the neuron network. The synchronized spontaneous Ca^{2+} oscillations we observed were confined to the neuron networks. Although glia cells sometimes were observed irregularly [Ca^{2+}]_i elevation, they did not directly participate in the Ca^{2+} oscillation of neuron networks. We could deduce that glia cells might act as a supporter in maintaining and modifying the synchronized spontaneous Ca^{2+} oscillation in neuron networks. We detected the mechanism of the Ca^{2+} oscillation in our culture, and found that it was mainly mediated by non-NMDA-type glutamatergic transmission. NMDA-type glutamatergic transmission only partially inhibited the spike amplitude. GABAergic synaptic transmission might act as a regulator of the Ca^{2+} oscillation.

PD98059 has long been used as a specific MEK inhibitor to study the involvement of the ERK pathway on cellular events as diverse as growth and differentiation, cell death and survival, and synaptic plasticity^[30]. Our results show

that PD98059 acutely inhibits the synchronized Ca^{2+} oscillations in a dose-dependent manner. When the concentration (10 $\mu\text{mol/L}$) is too low to inhibit MAPK activation^[31], PD98059 has little effect on the Ca^{2+} spikes. However, at higher concentrations (30 $\mu\text{mol/L}$ and 60 $\mu\text{mol/L}$) PD98059 significantly and acutely inhibits synchronized Ca^{2+} oscillations. We also showed that PD98059 inhibited the synchronized Ca^{2+} oscillations mainly through inhibition of MEK. This conclusion is based on the observation that SB202474, a structural analog of PD98059, which is usually used as a negative control for MAPK inhibitor studies, has no effect on the Ca^{2+} spike at a higher concentration (60 $\mu\text{mol/L}$). Although SB202474 has a strange effect on the Ca^{2+} oscillations at a lower concentration (10 $\mu\text{mol/L}$), by slightly increasing the frequency of the Ca^{2+} spikes and decreasing the amplitude of the spikes, it has no effect at a higher concentration (60 $\mu\text{mol/L}$). The effect of SB202474 on Ca^{2+} spikes at a lower concentration might result from its nonspecific activation of the Ca^{2+} channel, an idea we will try to explain in future studies. The salient point of this study is that we can conclude that

SB202474 does not inhibit Ca^{2+} spikes, but activates Ca^{2+} spikes at a lower concentration. As SB202474 is an inactive analog of PD98059, these observations indicate that PD98059 inhibits Ca^{2+} oscillations mainly through the inhibition of MEK, but not its side-effect on Ca^{2+} channel^[30,32].

Many studies have implied that the MAPK cascade might participate in $[Ca^{2+}]_i$ regulation^[22,23]. The synchronized spontaneous Ca^{2+} spikes in networked neurons represent the periodic firing of action potentials, which are believed to play a major role in the development and plasticity of neuronal circuitry^[19], and the encoded information of the spontaneous Ca^{2+} oscillations was reported to lie in their frequency or amplitude^[33]. Therefore, the inhibitory effect of PD98059 on the frequency and amplitude of spontaneous Ca^{2+} oscillations reported here implied that the MAPK cascade was required to maintain the spontaneous Ca^{2+} oscillations in developing hippocampal neurons. We know that MAPKs are a family of serine/threonine protein kinases which have classically been studied as regulators of cell proliferation and differentiation. The most important and well-known member of the MAPK family is ERK, which is initiated by growth factor receptor signaling. ERKs are extensively expressed in dendrites and somas of pyramidal neurons of the adult nervous system and can be activated by several neurotransmitters in neuronal culture system^[34]. These points suggest that MAPKs might be excellent candidates for regulation of synaptic plasticity in post-mitotic neurons. MAPKs have been reported to regulate glutamate release and participate in the introduction of LTP^[12]. Previous findings have shown that the MAPK cascade regulates synaptic transmission, and our work substantiates this by providing the time-course of PD98059 actions on synaptic transmission (the synchronized Ca^{2+} transients). It is also well known that Ca^{2+} plays an important role in the epileptiform discharge^[35,36]. The synchronized spontaneous Ca^{2+} oscillations in the neuron network^[37] are usually considered a kind of spontaneous epileptiform activity. Zhao *et al* reported that ERK1/2 was required for the induction of group I metabotropic glutamate receptor-mediated epileptiform discharges. Murray *et al* reported that PD98059 protected hippocampal neurons from seizure-like events^[38]. Our results provide further evidence for the effect of PD98059 on the hippocampal network. The inhibitory effect of PD98059 on synchronized spontaneous Ca^{2+} oscillations through MAPK might be used to develop drugs for epileptiform therapy.

References

1 Seger R, Krebs EG. The MAPK signaling cascade. *FASEB J* 1995;

9: 726–35.
 2 Lazou A, Sugden PH, Clerk A. Activation of mitogen-activated protein kinases (p38-MAPKs, SAPKs/JNKs and ERKs) by the G-protein-coupled receptor agonist phenylephrine in the perfused rat heart. *Biochem J* 1998; 332: 459–65.
 3 Haq SE, Clerk A, Sugden PH. Activation of mitogen-activated protein kinases (p38-MAPKs, SAPKs/JNKs and ERKs) by adenosine in the perfused rat heart. *FEBS Lett* 1998; 434: 305–8.
 4 Bogoyevitch MA, Gillespie-Brown J, Ketterman AJ, Fuller SJ, Ben-Levy R, Ashworth A, *et al*. Stimulation of the stress-activated mitogen-activated protein kinase subfamilies in perfused heart. p38/RK mitogen-activated protein kinases and c-Jun N-terminal kinases are activated by ischemia/reperfusion. *Circ Res* 1996; 79: 162–73.
 5 Stofega MR, Yu CL, Wu J, Jove R. Activation of extracellular signal-regulated kinase (ERK) by mitogenic stimuli is repressed in v-Src-transformed cells. *Cell Growth Differ* 1997; 8: 113–9.
 6 Marais R, Marshall CJ. Control of the ERK MAP kinase cascade by Ras and Raf. *Cancer Surv* 1996; 27: 101–25.
 7 Harrisingh MC, Perez-Nadales E, Parkinson DB, Malcolm DS, Mudge AW, Lloyd AC. The Ras/Raf/ERK signalling pathway drives Schwann cell dedifferentiation. *EMBO J* 2004; 23: 3061–71.
 8 Boulton TG, Cobb MH. Identification of multiple extracellular signal-regulated kinases (ERKs) with antipeptide antibodies. *Cell Regul* 1991; 2: 357–71.
 9 Thomas KL, Hunt SP. The regional distribution of extracellularly regulated kinase-1 and -2 messenger RNA in the adult rat central nervous system. *Neuroscience* 1993; 56: 741–57.
 10 Fiore RS, Bayer VE, Pelech SL, Posada J, Cooper JA, Baraban JM. p42 mitogen-activated protein kinase in brain: prominent localization in neuronal cell bodies and dendrites. *Neuroscience* 1993; 55: 463–72.
 11 Alessi DR, Cuenda A, Cohen P, Dudley DT, Saltiel AR. PD 098059 is a specific inhibitor of the activation of mitogen-activated protein kinase kinase *in vitro* and *in vivo*. *J Biol Chem* 1995; 270: 27489–94.
 12 English JD, Sweatt JD. A requirement for the mitogen-activated protein kinase cascade in hippocampal long term potentiation. *J Biol Chem* 1997; 272: 19103–6.
 13 Coogan AN, O’Leary DM, O’Connor JJ. P42/44 MAP kinase inhibitor PD98059 attenuates multiple forms of synaptic plasticity in rat dentate gyrus *in vitro*. *J Neurophysiol* 1999; 81: 103–10.
 14 Wang X, Gruenstein EI. Mechanism of synchronized Ca^{2+} oscillations in cortical neurons. *Brain Res* 1997; 767: 239–49.
 15 Tanaka T, Saito H, Matsuki N. Intracellular calcium oscillation in cultured rat hippocampal neurons: a model for glutamatergic neurotransmission. *Jpn J Pharmacol* 1996; 70: 89–93.
 16 Liu XH, Lu GW, Cui ZJ. Calcium oscillations in freshly isolated neonatal rat cortical neurons. *Acta Pharmacol Sin* 2002; 23: 577–81.
 17 Koizumi S, Inoue K. Inhibition by ATP of calcium oscillations in rat cultured hippocampal neurones. *Br J Pharmacol* 1997; 122: 51–8.
 18 Lisman JE. Bursts as a unit of neural information: making unreliable synapses reliable. *Trends Neurosci* 1997; 20: 38–43.
 19 Bacci A, Verderio C, Pravettoni E, Matteoli M. Synaptic and intrinsic mechanisms shape synchronous oscillations in hippoc-

- ampal neurons in culture. *Eur J Neurosci* 1999; 11: 389–97.
- 20 Traub RD, Wong RK. Cellular mechanism of neuronal synchronization in epilepsy. *Science* 1982; 216: 745–7.
- 21 Miles R, Wong RK. Single neurones can initiate synchronized population discharge in the hippocampus. *Nature* 1983; 306: 371–3.
- 22 Ansari HR, Husain S, Abdel-Latif AA. Activation of p42/p44 mitogen-activated protein kinase and contraction by prostaglandin F₂α, ionomycin, and thapsigargin in cat iris sphincter smooth muscle: inhibition by PD98059, KN-93, and isoproterenol. *J Pharmacol Exp Ther* 2001; 299: 178–86.
- 23 Kupzig S, Walker SA, Cullen PJ. The frequencies of calcium oscillations are optimized for efficient calcium-mediated activation of Ras and the ERK/MAPK cascade. *Proc Natl Acad Sci USA* 2005; 102: 7577–82.
- 24 Doherty P, Williams G, Williams EJ. CAMs and axonal growth: a critical evaluation of the role of calcium and the MAPK cascade. *Mol Cell Neurosci* 2000; 16: 283–95.
- 25 Banker GA, Cowan WM. Rat hippocampal neurons in dispersed cell culture. *Brain Res* 1977; 126: 397–42.
- 26 Gee KR, Brown KA, Chen WN, Bishop-Stewart J, Gray D, Johnson I. Chemical and physiological characterization of Fluo-4 Ca²⁺-indicator dyes. *Cell Calcium* 2000; 27: 97–106.
- 27 Leinekugel X, Medina I, Khalilov I, Ben-Ari Y, Khazipov R. Ca²⁺ oscillations mediated by the synergistic excitatory actions of GABA(A) and NMDA receptors in the neonatal hippocampus. *Neuron* 1997; 18: 243–55.
- 28 Dudley DT, Pang L, Decker SJ, Bridges AJ, Saltiel AR. A synthetic inhibitor of the mitogen-activated protein kinase cascade. *Proc Natl Acad Sci USA* 1995; 92: 7686–9.
- 29 Brewer GJ, Torricelli JR, Evege EK, Price PJ. Optimized survival of hippocampal neurons in B27-supplemented Neurobasal, a new serum-free medium combination. *J Neurosci Res* 1993; 35: 567–76.
- 30 Pereira DB, Carvalho AP, Duarte CB. Non-specific effects of the MEK inhibitors PD098,059 and U0126 on glutamate release from hippocampal synaptosomes. *Neuropharmacology* 2002; 42: 9–19.
- 31 Gould MC, Stephano JL. MAP kinase, meiosis, and sperm centrosome suppression in *Urechis caupo*. *Dev Biol* 1999; 216: 348–58.
- 32 Gould MC, Stephano JL. Inactivation of Ca²⁺ action potential channels by the MEK inhibitor PD98059. *Exp Cell Res* 2000; 260: 175–9.
- 33 Gu X, Spitzer NC. Distinct aspects of neuronal differentiation encoded by frequency of spontaneous Ca²⁺ transients. *Nature* 1995; 375: 784–7.
- 34 English JD, Sweatt JD. Activation of p42 mitogen-activated protein kinase in hippocampal long term potentiation. *J Biol Chem* 1996; 271: 24329–32.
- 35 Pisani A, Bonsi P, Martella G, De Persis C, Costa C, Pisani F, *et al*. Intracellular calcium increase in epileptiform activity: modulation by levetiracetam and lamotrigine. *Epilepsia* 2004; 45: 719–28.
- 36 Shang ZC, Sun BZ, Chen ZN, Wang J, Feng Q, Wang W, *et al*. Effect of PD98059 on Ras-MAPK signal transduction pathway of chronic myelogenous leukemia. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi* 2003; 19: 54–5.
- 37 Yaari Y, Konnerth A, Heinemann U. Spontaneous epileptiform activity of CA1 hippocampal neurons in low extracellular calcium solutions. *Exp Brain Res* 1983; 51: 153–6.
- 38 Murray B, Alessandrini A, Cole AJ, Yee AG, Furshpan EJ. Inhibition of the p44/42 MAP kinase pathway protects hippocampal neurons in a cell-culture model of seizure activity. *Proc Natl Acad Sci USA* 1998; 95: 11975–80.