

## Full-length article

## Stimulation of Na<sup>+</sup>-Ca<sup>2+</sup> exchange by purified antibody against alpha-2 repeat of Na<sup>+</sup>-Ca<sup>2+</sup> exchanger in rat cardiomyocytes

Qi-long FENG<sup>1</sup>, Dong-mei WU<sup>2</sup>, Xiang-li CUI<sup>1</sup>, Hua-chen ZHAO<sup>1</sup>, Guo-quan FAN<sup>3</sup>, Lu-ying ZHAO<sup>1</sup>, Bo-wei WU<sup>1,4</sup>

Departments of <sup>1</sup>Physiology, <sup>2</sup>Pharmacology, and <sup>3</sup>Microbiology and Immunology, Shanxi Medical University, Taiyuan 030001, China

### Key words

Na<sup>+</sup>-Ca<sup>2+</sup> exchanger; patch-clamp technique; cardiac myocyte; antibody

<sup>4</sup> Correspondence to Prof Bo-wei WU.

Phn/Fax 86-351-413-5075.

E-mail wubowei299@sohu.com

Received 2008-03-18

Accepted 2008-07-08

doi: 10.1111/j.1745-7254.2008.00854.x

### Abstract

**Aim:** The aim of the present study was to investigate the effect of the antibody against alpha-2 repeat on Na<sup>+</sup>-Ca<sup>2+</sup> exchanger (NCX) current ( $I_{Na/Ca}$ ). To evaluate the functional specificity of this antibody, its effects on L-type Ca<sup>2+</sup> current ( $I_{Ca,L}$ ), voltage-gated Na<sup>+</sup> current ( $I_{Na}$ ) and delayed rectifier K<sup>+</sup> current ( $I_K$ ) were also observed. **Methods:** The whole-cell patch-clamp technique was used in this study. **Results:** The antibody against alpha-2 repeat augmented both the outward and inward Na<sup>+</sup>-Ca<sup>2+</sup> exchanger current concentration-dependently, with EC<sub>50</sub> values of 27.9 nmol/L and 24.7 nmol/L, respectively. Meanwhile, the antibody could also increase  $I_{Ca,L}$  in a concentration-dependent manner with the EC<sub>50</sub> of 33.6 nmol/L. Effects of the antibody on  $I_{Na}$  and  $I_K$  were not observed in the present study. **Conclusion:** The present results suggest that antibody against alpha-2 repeat is a stimulating antibody to NCX and could also increase  $I_{Ca,L}$  in a concentration-dependent manner, but did not have an obvious effect on  $I_{Na}$  and  $I_K$ .

### Introduction

The process of Na<sup>+</sup>-Ca<sup>2+</sup> exchange was first identified in guinea pig atria by Reuter and Seitz in 1968<sup>[1]</sup>, and in the squid giant axon by Baker *et al* shortly after in 1969<sup>[2]</sup>. The entity, the so-called Na<sup>+</sup>-Ca<sup>2+</sup> exchanger (NCX), is an ion transport protein that catalyzes electrogenic antitransport of Na<sup>+</sup> and Ca<sup>2+</sup> across the plasma membrane in a coupling ratio of 3 Na<sup>+</sup>:1 Ca<sup>2+</sup> and exists in the plasma membrane of almost all animal cells<sup>[3]</sup>. It is in cardiomyocytes, however, that the exchanger is highly expressed and plays an important role in Ca<sup>2+</sup> homeostasis. The NCX system is the primary mechanism responsible for transsarcolemmal Ca<sup>2+</sup> extrusion. There is a general agreement that the majority of Ca<sup>2+</sup> entry through voltage-dependent Ca<sup>2+</sup> channel is transported out of the cell by NCX<sup>[4,5]</sup>. Moreover, it has also been suggested that Ca<sup>2+</sup> entry mediated by NCX in Ca<sup>2+</sup> influx mode contributed directly to contraction of failing human ventricular myocytes during the early period of the cardiomyocyte action potential<sup>[6]</sup>.

The study of NCX in molecular biology started after the cloning of canine cardiac NCX by Nicoll and Philipson<sup>[7]</sup>. Later in 1997, Schwarz and Benzer<sup>[8]</sup> first identified

the highly conserved regions in all known members of the NCX family, designated the alpha-1 and alpha-2 repeats. These regions are highly conserved among different exchangers and between one another. In cardiac NCX, the alpha-1 repeat comprises most of the putative transmembrane segment 2 and 3 (TM2 and TM3) and their connecting loop, whereas alpha-2 locates in putative TM7 and its C-terminal sequence<sup>[9]</sup>.

A recent study by Nicoll and Iwamoto demonstrated that the NCX1 had oppositely oriented reentrant loop domains in alpha-1 and alpha-2 repeats, and that these reentrant domains in the alpha-repeats might be involved in the formation of the ion transport pathway<sup>[9]</sup>. Mutation analysis also showed that alpha-repeats were involved in the interaction of the exchanger with transport substrates (Na<sup>+</sup> and Ca<sup>2+</sup>), Ni<sup>2+</sup>, Li<sup>+</sup> and KB-R7943<sup>[10–12]</sup>.

Now that the alpha-repeats regions were considered important in the ion binding and translocation, it is possible that the antibodies against alpha-1 repeat and alpha-2 repeat may have a crucial action on Na<sup>+</sup>-Ca<sup>2+</sup> exchanger activity. However, the effect of the antibody against alpha-2 repeat on  $I_{NCX}$  and its specificity is unclear until now. The goal of this study is to identify the effects of the antibody on Na<sup>+</sup>-

Ca<sup>2+</sup> exchanger current using a whole-cell patch-clamp technique. Furthermore, the functional specificity of the antibody was also investigated in adult rat cardiomyocytes.

## Materials and methods

**Ventricular myocyte isolation** Single ventricular myocytes were isolated from Wistar rats (250–300 g) using an enzymatic dissociation procedure similar to that described by Mubagwa *et al*<sup>[13]</sup>. In brief, rats were anesthetized with sodium pentobarbital (30 mg/kg, ip) 30 min after having received heparin (500 U, ip). The heart was quickly removed, rinsed in ice-cold Ca<sup>2+</sup> free Tyrode's solution and perfused with oxygenated Ca<sup>2+</sup> free Tyrode's solution (at 37 °C) via aorta for approximately 7–8 min to wash out the blood. The composition of Tyrode's solution was (in mmol/L): NaCl 135, KCl 5.4, CaCl<sub>2</sub> 1.8, MgCl<sub>2</sub> 1.0, NaH<sub>2</sub>PO<sub>4</sub> 0.33, HEPES (4-(2-hydroxyethyl)piperazine-1-ethanesulfonic acid) 10, glucose 10 (pH adjusted to 7.4 with NaOH). The composition of Ca<sup>2+</sup>-free Tyrode's solution was the same to Tyrode's solution except for the absence of CaCl<sub>2</sub>. Then the perfusate was switched to enzyme solution for 8–10 min. Enzyme solution contained (in mmol/L): NaCl 135, KCl 5.4, CaCl<sub>2</sub> 75 μmol/L, MgCl<sub>2</sub> 1.0, NaH<sub>2</sub>PO<sub>4</sub> 0.33, HEPES 10, glucose 10, taurine 20, collagenase P (Boehringer Mannheim, Mannheim, Germany) 100 mg/L (pH adjusted to 7.4 with NaOH). The ventricle was then separated and minced with a pair of surgical scissors in the Kraftbrühe (KB) solution. The isolated myocytes were stored in KB solution at room temperature (22 °C) at least 4 h before use. KB solution was composed of (in mmol/L): KOH 85, L-glutamic acid 50, KCl 30, MgCl<sub>2</sub> 1.0, KH<sub>2</sub>PO<sub>4</sub> 30, glucose 10, taurine 20, HEPES 10, EGTA[ethyleneglycol-bis(β-amino-ethylether)-N,N,N',N'-tetraacetic acid] 0.5 (pH adjusted to 7.4 with KOH 1 mol/L).

**Electrophysiological measurement** Voltage-clamp recording was carried out in the whole-cell configuration of the patch-clamp method<sup>[14]</sup> using a Patchclamp Amplifier (Axopatch-200A, Axon Instruments, Foster City, CA, USA). Patch electrodes were made from thin-walled glass capillaries and the electrodes with resistance of 2–4 MΩ were filled with the pipette solution. Cell capacitance was measured by the method described by Coetzee *et al*<sup>[15]</sup>. Analysis was carried out using pClampfit 8.0 software (Axon Instruments).

For the measurement of Na<sup>+</sup>-Ca<sup>2+</sup> exchange current ( $I_{Na/Ca}$ ), the extracellular (test) solution contained (in mmol/L): NaCl 140, CaCl<sub>2</sub> 2.0, MgCl<sub>2</sub> 2.0, glucose 10, HEPES 5.0 (pH adjusted to 7.4 with CsOH). In addition,

the Na<sup>+</sup>-K<sup>+</sup> pump, K<sup>+</sup> channel and Ca<sup>2+</sup> channel were blocked with ouabain (Sigma Chemical, St Louis, MO, USA) 20 μmol/L, BaCl<sub>2</sub> 1.0 mmol/L, CsCl 2.0 mmol/L and nifedipine (Sigma) 1.0 μmol/L. The pipette solution contained (in mmol/L): EGTA 42, CaCl<sub>2</sub> 29, MgCl<sub>2</sub> 13, aspartate, K<sub>2</sub>ATP 10, Na<sub>2</sub>-cretinophosphate 5.0, TEA (tetraethylammonium) (Sigma) 20, HEPES 5.0 (pH adjusted to 7.4 with CsOH). To measure L-type Ca<sup>2+</sup> current ( $I_{Ca,L}$ ), the extracellular (test) solution contained (in mmol/L): NaCl 135, CaCl<sub>2</sub> 1.8, MgCl<sub>2</sub> 1.0, KCl 5.4, glucose 10, NaH<sub>2</sub>PO<sub>4</sub> 0.33, HEPES 10 (pH adjusted to 7.4 with NaOH). The pipette solution contained (in mmol/L): EGTA 10, KCl 140, Na<sub>2</sub>ATP 2.0, HEPES 5.0, 4-AP 5.0, MgCl<sub>2</sub> 1.0 (pH adjusted to 7.3 with KOH). To record voltage-gated Na<sup>+</sup> current ( $I_{Na}$ ), the extracellular (test) solution contained (in mmol/L): NaCl 60, CsCl 5.0, CdCl<sub>2</sub> 0.1, MgCl<sub>2</sub> 2.5, glucose 10, 4-AP 5.0, HEPES 5.0, saccharose 80 (pH adjusted to 7.4 with NaOH). The pipette solution contained (in mmol/L): EGTA 11, KCl 130, Na<sub>2</sub>ATP 5.0, HEPES 10, MgCl<sub>2</sub> 2.0, CaCl<sub>2</sub> 1.0, 4-AP 5.0 (pH adjusted to 7.2 with CsOH). For the measurement of delayed rectifier K<sup>+</sup> current ( $I_K$ ), the extracellular (test) solution contained (in mmol/L): NaCl 145, KCl 4.0, MgCl<sub>2</sub> 1.0, HEPES 10, glucose 5.0, CaCl<sub>2</sub> 0.1, CdCl<sub>2</sub> 0.1 (pH adjusted to 7.4 with NaOH). The pipette solution contained (in mmol/L): KCl 130, MgCl<sub>2</sub> 2.0, CaCl<sub>2</sub> 1.0, EGTA 11, MgATP 5, K<sub>2</sub>ATP 5.0, HEPES 10 (pH adjusted to 7.4 with KOH).

**Antibody preparation** The antibody against alpha-2 repeat of Na<sup>+</sup>-Ca<sup>2+</sup> exchanger was prepared in our laboratory<sup>[16]</sup>. Briefly, peptide corresponding to alpha-2 repeat (815 TFASKVAATQDQYADASIGNVTGSN 839) in cardiac NCX was synthesized by CL (Xi'an) Bioscientific Incorporation. Then the rats were randomly divided into two groups: control and immunized groups. Rats in immunized groups were immunized with the synthesized alpha-2 repeat emulsified in equal volume of Freund's adjuvant (CFA, Sigma). The positive antiserum with high titer (≥1:640 by enzyme-linked immunosorbent assay [ELISA]) were affinity-purified using a Mab Trap Kit (Amersham Pharmacia Biotech, Uppsala, Sweden). The concentration of the purified antibody was determined using the method described by Bradford<sup>[17]</sup>. The control group received the same disposal as described above except that the peptide was substituted for saline solution.

**Data analysis** Results were expressed as mean±SD, and analyzed with least-significant difference (LSD) test of ANOVA in SPSS 11.  $P < 0.05$  was considered significant. EC<sub>50</sub> values were determined using GraphPad Prism 4


software.

## Results

### Measurement of the Na<sup>+</sup>-Ca<sup>2+</sup> exchange current

( $I_{Na/Ca}$ )  $I_{Na/Ca}$  was measured as the current sensitive to 5.0 mmol/L Ni<sup>2+</sup>[18] by the voltage protocol shown at the top of Figure 1. Ramp voltage-clamp pulse from 60 to -120 mV (90 mV/s) was applied from a holding potential of -40 mV. The current-voltage relationship was constructed from the declining slope of the ramp pulse. After application of Ni<sup>2+</sup> at the concentration of 5.0 mmol/L, the current immediately decreased, at both positive and negative potentials (Figure 1). The difference between current-voltage relationships in the absence and presence of Ni<sup>2+</sup> reflected  $I_{Na/Ca}$  (Ni<sup>2+</sup>-sensitive current). Significant run-down of the Ni<sup>2+</sup>-sensitive current was not observed during the experiment.


**Stimulating effect of antibody against alpha-2 repeat on Na<sup>+</sup>-Ca<sup>2+</sup> exchange current ( $I_{Na/Ca}$ )** Antibody against



**Figure 1.** Measurement of Ni<sup>2+</sup>-sensitive Na<sup>+</sup>-Ca<sup>2+</sup> exchange current in rat ventricular myocytes. The voltage protocol is shown in the top panel (see text for details). (A) Current-voltage relationship before (trace a) and after (trace b) application of 5.0 mmol/L NiCl<sub>2</sub>. (B) Ni<sup>2+</sup>-sensitive Na<sup>+</sup>-Ca<sup>2+</sup> exchange current (numerical subtraction of a-b).

alpha-2 repeat had a stimulating effect on performance of Na<sup>+</sup>-Ca<sup>2+</sup> exchange, as demonstrated by the present study. It was shown that this antibody increased both the outward current and inward current concentration-dependently with EC<sub>50</sub> values of 27.9 nmol/L and 24.7 nmol/L, respectively (Figure 2, Table 1). The stimulating effects on both outward and inward current of  $I_{Na/Ca}$  were abolished when the antibody was incubated with synthesized alpha-2 repeat before it was applied to cells.

**Effects of the antibody against alpha-2 repeat on  $I_{Ca,L}$ ,  $I_{Na}$  and  $I_K$**  To evaluate the functional selectivity of the antibody against alpha-2 repeat, its effects on  $I_{Ca,L}$ ,  $I_{Na}$  in adult rat hearts and  $I_K$  in adult guinea-pig hearts were




**Figure 2.** Representative traces showing effect of antibody against alpha-2 on Na<sup>+</sup>-Ca<sup>2+</sup> exchange current in rat ventricular myocytes. (A) Current-voltage relationship before (trace a, control) and after application of antibody against alpha-2 at 10 nmol/L (trace b), 20 nmol/L (trace c), 40 nmol/L (trace d), 80 nmol/L (trace e) and 160 nmol/L (trace f), respectively. Trace g was recorded after application of 5 mmol/L Ni<sup>2+</sup>. (B) Ni<sup>2+</sup>-sensitive Na<sup>+</sup>-Ca<sup>2+</sup> exchange current before (a-g) and after application of antibody against alpha-2 at concentration of 10 nmol/L (b-g), 20 nmol/L (c-g), 40 nmol/L (d-g), 80 nmol/L (e-g) and 160 nmol/L (f-g), respectively. The voltage protocol is shown in the top panel (see text for details).

**Table 1.** Stimulating effect of antibody against alpha-2 repeat of Na<sup>+</sup>-Ca<sup>2+</sup> exchanger on  $I_{Na/Ca}$ . Values were expressed as mean±SD. <sup>b</sup>*P*<0.05, <sup>c</sup>*P*<0.01 vs control. Membrane current density is expressed as membrane current (pA) per cell capacitance (pF).

Antibody against alpha-2 repeat/nmol·L <sup>-1</sup>		$I_{Na/Ca}/pA \cdot pF^{-1}$			
repeat/nmol·L <sup>-1</sup>	<i>n</i>	Increment (%)		Increment (%)	
		+ 50 mV		- 100 mV	
0 (Control)	10	0.43±0.09		0.37±0.08	
10	8	0.62±0.09	44	0.51±0.11	38
20	8	0.73±0.10 <sup>b</sup>	70	0.64±0.08 <sup>b</sup>	73
40	8	1.04±0.11 <sup>c</sup>	142	0.86±0.10 <sup>c</sup>	132
80	8	1.12±0.13 <sup>c</sup>	161	0.92±0.10 <sup>c</sup>	148
160	8	1.17±0.14 <sup>c</sup>	172	0.95±0.11 <sup>c</sup>	157

also observed. It was shown that this antibody could also increase  $I_{Ca,L}$  in a concentration-dependent manner and EC<sub>50</sub> was 33.6 nmol/L. After washing with Tyrode solution, the effect could be partly abolished. Nifedipine could inhibit the above inward current completely, which proved that the current was  $I_{Ca,L}$  (Figure 3, Table 2). Furthermore, the current-voltage (*I-V*) relationship curve did not shift after application of the 40 nmol/L antibody, although peak Ca<sup>2+</sup> current increased at +10 mV (Figure 4). Effects of the antibody on  $I_{Na}$  (Figure 5) and  $I_K$  (Figure 6) were not observed in the present study.



**Figure 3.** Representative traces showing effect of antibody against alpha-2 on L-type Ca<sup>2+</sup> current in rat ventricular myocytes. Trace a, control. Traces b, c, d, e and f, after application of antibody against alpha-2 at 10 nmol/L, 20 nmol/L, 40 nmol/L, 80 nmol/L and 160 nmol/L, respectively. Trace g, after application of 1 μmol/L nifedipine. The voltage protocol is shown in the top panel.


**Table 2.** Effect of antibody against alpha-2 repeat of Na<sup>+</sup>-Ca<sup>2+</sup> exchanger on  $I_{Ca,L}$ . Values are presented as mean±SD. <sup>b</sup>*P*<0.05, <sup>c</sup>*P*<0.01 vs control. Membrane current density is expressed as membrane current (pA) per cell capacitance (pF).

Antibody against alpha-2 repeat/nmol·L <sup>-1</sup>	<i>n</i>	$I_{Ca,L}/pA \cdot pF^{-1}$
0 (Control)	11	2.51±0.43
10	6	2.98±0.26
20	5	3.30±0.41 <sup>b</sup>
40	5	3.74±0.40 <sup>c</sup>
80	5	4.14±0.36 <sup>c</sup>
160	5	4.37±0.30 <sup>c</sup>

## Discussion

Under physiological conditions, NCX operates mainly in Ca<sup>2+</sup> efflux mode (Na<sup>+</sup> influx), and only a very small quantity of Ca<sup>2+</sup> enters the cell during the very early rising phase (1–4 ms) of the action potential *via* NCX in Ca<sup>2+</sup> influx mode<sup>[19]</sup>. This is not the case, however, in pathological conditions such as heart failure (HF). It was shown that NCX could bring a larger amount of Ca<sup>2+</sup> into the cell in HF than in normal conditions during the action potential, which partly compensated for the downregulated SR Ca-ATPase function and supported contraction in the patients with HF<sup>[20–22]</sup>. Because the antibody against alpha-2 repeat was proven to increase both  $I_{Na/Ca}$  and  $I_{Ca,L}$  in the present study, it may have the therapeutic potential to improve systolic function in HF patients by increasing Ca<sup>2+</sup> entry *via* NCX and also the L-type Ca<sup>2+</sup> channel.


Our investigation showed that the antibody against alpha-2 repeat could also increase  $I_{Ca,L}$  besides enhancement of Na<sup>+</sup>-Ca<sup>2+</sup> exchange current. Moreover, the current-voltage (*I-V*) relationship curve of  $I_{Ca,L}$  was not shifted by the antibody (40 nmol/L) and EC<sub>50</sub> of  $I_{Ca,L}$  was similar to that of  $I_{Na/Ca}$ . An early study has shown that in cardiac muscle where Ca influx across the sarcolemma is essential for contraction, the L-type Ca<sup>2+</sup> channel has four homologous domains (I–IV), each comprising six transmembrane segments (S1–S6)<sup>[23]</sup>. Mutational analysis indicated that S5–S6 linkers were highly conserved in domains I–IV<sup>[23]</sup> and contributed to formation of the ion pore<sup>[24]</sup>. Residues 1079–1110 were supposed to be located on the S5–S6 linker in domain III<sup>[25]</sup>. Meanwhile, mutation analysis showed that residues 815–839 of alpha-2 repeat in NCX were involved in the interaction of the exchanger with Na<sup>+</sup> and Ca<sup>2+</sup><sup>[9–12]</sup>. We compared the amino acid alignment of the alpha-2 repeat in NCX with the residues 1079–1110




**Figure 4.** Current traces and current-voltage (*I-V*) relationship of L-type Ca<sup>2+</sup> current. (A, B) Original Ca<sup>2+</sup> current traces in the absence (A) and presence (B) of antibody against alpha-2 repeat (40 nmol/L). The currents were measured in response to depolarizing voltage clamp steps of 500 ms in the voltage range between -40 and 50 mV from a holding potential of -40 mV. (C) Current-voltage (*I-V*) relationship of Ca<sup>2+</sup> currents in the absence and presence of 40 nmol/L antibody against alpha-2 repeat. *n*=13 cells from 10 hearts.

of the L-type Ca<sup>2+</sup> channel using EMBOSS Pairwise Alignment Algorithms (European Bioinformatics Institute), which showed that the degree of amino acid similarity was 28.1% between these two functional segments (Figure 7), providing a clue for the non-specific action of the antibody on I<sub>Ca,L</sub>.

From the genetic and evolutionary points of view, intramolecular repeats are thought to arise from intra-genetic duplications, and can not survive throughout evolution unless they are essential to protein function<sup>[8]</sup>. As we know, alpha-repeats exist in all known members of the NCX family, which implies that alpha-repeats arose early in the evolutionary history and furthermore, their existing as a tandem pair is essential for the protein to operate properly. In the present study, we first observed the effect of the antibody against alpha-2 repeat on I<sub>Na/Ca</sub> in adult rat



**Figure 5.** Representative traces showing effect of antibody against alpha-2 on voltage-gated Na<sup>+</sup> current at 10 nmol/L, 20 nmol/L, 40 nmol/L, 80 nmol/L and 160 nmol/L, respectively in rat ventricular myocytes.



**Figure 6.** Traces showing effect of antibody against alpha-2 on delayed rectifier K<sup>+</sup> current at 10 nmol/L, 20 nmol/L, 40 nmol/L, 80 nmol/L and 160 nmol/L, respectively in guinea pig ventricular myocytes.

```

815   TFASKVAATQD---QYADASIGNV----TGSN   839
1079  LFKGKLYTCS DSSKQTEAESKGN YITYTKTEV  1110
    
```

**Figure 7.** Amino acid alignment of alpha-2 repeat of Na<sup>+</sup>-Ca<sup>2+</sup> exchanger (upper) and pore region of L-type Ca<sup>2+</sup> channel (lower). Residues identical between two segments are in bold. Alignment insertions are indicated with a dash.

cardiomyocytes with the whole-cell patch clamp technique. Our results showed that antibody against alpha-2 repeat was a stimulating antibody because it increased I<sub>Na/Ca</sub> in a concentration-dependent manner, which provided supplemental evidence that alpha-2 repeat was essential to translocation of sodium and calcium by NCX.

Recent topological research showed that the alpha-2 repeat comprised putative transmembrane 7 and its

C-terminal segment and formed a domain mostly accessible from the cytoplasm<sup>[26]</sup>. However, our results in this study indicated that the antibody against alpha-2 repeat could stimulate Na<sup>+</sup>-Ca<sup>2+</sup> exchange from the external side of the cardiomyocytes. Then why did the antibody play its role from the extracellular side? One possibility is that the interaction between antibody against alpha-2 repeat and NCX might lead to conformation alteration of the exchanger molecule, just as what happens to KB-R7943<sup>[27]</sup>.

The present study showed that the antibody against alpha-2 repeat of Na<sup>+</sup>-Ca<sup>2+</sup> exchanger was a stimulating antibody to NCX and could also increase  $I_{Ca,L}$  in a concentration-dependent manner, whereas it did not have an obvious effect on  $I_{Na}$  and  $I_K$ .

### Author contribution

Prof Bo-wei WU designed research; Dr Qi-long FENG, Dong-mei WU, Hua-chen ZHAO and Guo-quan FAN performed research; Lu-ying ZHAO contributed new analytical tools and reagents; Dr Qi-long FENG and Xiang-li CUI analyzed data; Dr Qi-long FENG, Dong-mei WU, and Prof Bo-wei WU wrote the paper.

### References

- 1 Reuter H, Seitz N. The dependence of calcium efflux from cardiac muscle on temperature and external ion composition. *J Physiol* 1968; 195: 451–70.
- 2 Baker PF, Blaustein MP, Hodgkin AL, Steinhardt RA. The influence of calcium on sodium efflux in squid axons. *J Physiol* 1969; 200: 431–58.
- 3 Kang TM, Hilgemann DW. Multiple transport modes of the cardiac Na<sup>+</sup>/Ca<sup>2+</sup> exchanger. *Nature* 2004; 427: 544–8.
- 4 Bers DM. Altered cardiac myocyte Ca regulation in heart failure. *Physiology* 2006; 21: 380–7.
- 5 Reppel M, Fleischmann BK, Reuter H, Pillekamp F, Schunkert H, Hescheler J. Regulation of Na<sup>+</sup>/Ca<sup>2+</sup> exchange current in the normal and failing heart. *Ann NY Acad Sci* 2007; 1099: 361–72.
- 6 Weisser TJ, Piacentino V III, Gaughan JP, Margulies K, Houser SR. Calcium entry via Na/Ca exchange during the action potential directly contributes to contraction of failing human ventricular myocytes. *Cardiovasc Res* 2003; 57: 974–85.
- 7 Nicoll DA, Longoni S, Philipson KD. Molecular cloning and functional expression of the cardiac sarcolemmal Na<sup>+</sup>-Ca<sup>2+</sup> exchanger. *Science* 1990; 250: 562–5.
- 8 Schwarz EM, Benzer S. Calx, a Na-Ca exchanger gene of *Drosophila melanogaster*. *Proc Natl Acad Sci USA* 1997; 94: 10 249–54.
- 9 Iwamoto T, Nakamura TY, Pan Y, Uehara A, Imanaga I, Shigekawa M. Unique topology of the internal repeats in the cardiac Na<sup>+</sup>/Ca<sup>2+</sup> exchanger. *FEBS Lett* 1999; 446: 264–8.
- 10 Nicoll DA, Hryshko LV, Matsuoka S, Frank JS, Philipson KD. Mutation of amino acid residues in the putative transmembrane segments of the cardiac sarcolemmal Na<sup>+</sup>-Ca<sup>2+</sup> exchanger. *J Biol Chem* 1996; 271: 13 385–91.
- 11 Doering AE, Nicoll DA, Lu Y, Lu L, Weiss JN, Philipson KD. Topology of a functionally important region of the cardiac Na<sup>+</sup>/Ca<sup>2+</sup> exchanger. *J Biol Chem* 1998; 273: 778–83.
- 12 Iwamoto T, Uehara A, Nakamura TY, Imanaga I, Shigekawa M. Chimeric analysis of Na<sup>+</sup>/Ca<sup>2+</sup> exchangers NCX1 and NCX3 reveals structural domains important for differential sensitivity to external Ni<sup>2+</sup> or Li<sup>+</sup>. *J Biol Chem* 1999; 274: 23 094–102.
- 13 Mubagwa K, Stengl M, Flameng W. Extracellular divalent cations block a cation non-selective conductance unrelated to calcium channels in rat cardiac muscle. *J Physiol* 1997; 502: 235–47.
- 14 Hamill OP, Marty A, Neher E, Sakmann B, Sigworth FJ. Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflugers Arch* 1981; 391: 85–100.
- 15 Coetzee WA, Ichikawa H, Hearse DJ. Oxidant stress inhibits Na-Ca-exchange current in cardiac myocytes: mediation by sulfhydryl groups? *Am J Physiol Heart Circ Physiol* 1994; 266: H909–19.
- 16 Feng QL, Fan GQ, Zhao JH, Zhao HC, Wu BW. Acquisition and purification of the antibody against  $\alpha$ -1 and  $\alpha$ -2 repeats of Na<sup>+</sup>-Ca<sup>2+</sup> exchanger. *J Shanxi Med Univ* 2005; 36: 395–8.
- 17 Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976; 72: 248–54.
- 18 Kimura J, Miyamae S, Noma A. Identification of sodium-calcium exchange current in single ventricular cells of guinea-pig. *J Physiol* 1987; 384: 199–222.
- 19 Bers DM. Cardiac excitation-contraction coupling. *Nature* 2002; 415: 198–205.
- 20 Despa S, Islam M, Weber CR, Pogwizd SM, Bers DM. Intracellular Na<sup>+</sup> concentration is elevated in heart failure but Na/K pump function is unchanged. *Circulation* 2002; 105: 2543–8.
- 21 Weber CR, Piacentino V III, Ginsburg KS, Houser SR, Bers DM. Na<sup>+</sup>-Ca<sup>2+</sup> exchange current and submembrane [Ca<sup>2+</sup>] during the cardiac action potential. *Circ Res* 2002; 90: 182–9.
- 22 Weber CR, Piacentino V III, Houser SR, Bers DM. Dynamic regulation of sodium/calcium exchange function in human heart failure. *Circulation* 2003; 108: 2224–9.
- 23 Mikami A, Imoto K, Tanabe T, Niidome T, Mori Y, Takeshima H, *et al*. Primary structure and functional expression of the cardiac dihydropyridine-sensitive calcium channel. *Nature* 1989; 340: 230–3.
- 24 Nakai J, Adams BA, Imoto K, Beam KG. Critical roles of the S3 segment and S3–S4 linker of repeat I in activation of L-type calcium channels. *Proc Natl Acad Sci USA* 1994; 91: 1014–8.
- 25 Murphy BJ, Tuana BS. Calcium ions inhibit the allosteric interaction between the dihydropyridine and phenylalkylamine binding site on the voltage-gated calcium channel in heart sarcolemma but not in skeletal muscle transverse tubules. *Can J Physiol Pharmacol* 1990; 68: 1389–95.
- 26 Ren X, Nicoll DA, Philipson KD. Helix packing of the cardiac Na<sup>+</sup>-Ca<sup>2+</sup> exchanger: proximity of transmembrane segments 1, 2, and 6. *J Biol Chem* 2006; 281: 22 808–14.
- 27 Iwamoto T, Kita S, Uehara A. Structural domains influencing sensitivity to isothiourea derivative inhibitor KB-R7943 in cardiac Na<sup>+</sup>/Ca<sup>2+</sup> exchanger. *Mol Pharmacol* 2001; 59: 524–31.