

# Notch-mediated induction of N-cadherin and $\alpha 9$ -integrin confers higher invasive phenotype on rhabdomyosarcoma cells

A Masià<sup>1</sup>, A Almazán-Moga<sup>1</sup>, P Velasco<sup>2</sup>, J Reventós<sup>1</sup>, N Torán<sup>3</sup>, J Sánchez de Toledo<sup>2</sup>, J Roma<sup>\*,1</sup> and S Gallego<sup>\*,1,2</sup>

<sup>1</sup>Research Unit in Biomedicine and Translational and Pediatric Oncology, Vall d'Hebron Institut de Recerca (VHIR), Universitat Autònoma de Barcelona, Passeig Vall d'Hebron 119, Barcelona 08035, Spain; <sup>2</sup>Pediatric Oncology and Hematology Unit, Universitat Autònoma de Barcelona, Barcelona, Spain;

<sup>3</sup>Department of Pathology, Hospital Universitari Vall d'Hebron, Universitat Autònoma de Barcelona, Barcelona, Spain

**BACKGROUND:** Rhabdomyosarcoma (RMS) is the commonest type of soft-tissue sarcoma in children. Patients with metastatic RMS continue to have very poor prognosis. Recently, several works have demonstrated a connection between Notch pathway activation and the regulation of cell motility and invasiveness. However, the molecular mechanisms of this possible relationship remain unclear.

**METHODS:** The Notch pathway was manipulated pharmacologically and genetically. The mRNA changes were analysed by quantitative PCR and protein variations by western blot and immunofluorescence. Finally, the capabilities of RMS cells to adhere, heal a wound and invade were assessed in the presence of neuronal cadherin (N-cadherin)- and  $\alpha 9$ -integrin-blocking antibodies.

**RESULTS:** Cells treated with  $\gamma$ -secretase inhibitor showed lower adhesion capability and downregulation of N-cadherin and  $\alpha 9$ -integrin. Genetic manipulation of the Notch pathway led to concomitant variations in N-cadherin and  $\alpha 9$ -integrin. Treatment with anti-N-cadherin-blocking antibody rendered marked inhibition of cell adhesion and motility, while anti- $\alpha 9$ -integrin-blocking antibody exerted a remarkable effect on cell adhesion and invasiveness.

**CONCLUSION:** Neuronal cadherin and  $\alpha 9$ -integrin are postulated as leading actors in the association between the Notch pathway and promotion of cell adhesion, motility and invasion, pointing to these proteins and the Notch pathway itself as interesting putative targets for new molecular therapies against metastases in RMS.

*British Journal of Cancer* (2012) **107**, 1374–1383. doi:10.1038/bjc.2012.411 www.bjcancer.com

Published online 13 September 2012

© 2012 Cancer Research UK

**Keywords:** rhabdomyosarcoma; Notch; NCAD; ITGA9; invasion; soft-tissue sarcomas

Rhabdomyosarcoma (RMS), a malignant tumour of early onset, is the most common type of soft-tissue sarcoma in children. Regarding histopathological criteria, RMS can be divided into two main subtypes: embryonal and alveolar (eRMS and aRMS, respectively). The majority of aRMS (80–85%) contain one of the reciprocal chromosomal translocations: t(2;13)(q35;q14) or t(1;13)(p36;q14). These translocations generate the novel fusion genes *PAX3-FOXO1* and *PAX7-FOXO1*, respectively (Barr *et al*, 1993; Davis *et al*, 1994). However, no characteristic translocations have been described in eRMS. Patients with metastatic RMS have very poor prognosis and more intense therapies are thus indicated. Moreover, the major cause of death in these patients is the formation of distant metastases. The cellular components that control mobility, invasiveness and metastasis in RMS remain largely unknown. The molecules responsible for these processes should be identified before the development of targeted therapies focused on reducing metastasis in this neoplasia. Recently, we described the role of Notch pathway activation in controlling the migration and invasiveness processes in RMS (Roma *et al*, 2011). Similarly, the role of the Notch pathway in invasion and metastasis

has also been reported in osteosarcoma (Zhang *et al*, 2008; Zhang *et al*, 2010; Engin *et al*, 2009) and some adult cancers, such as colon, prostate, cervical carcinoma and glioblastoma among others (Zayzafoon *et al*, 2004; Pang *et al*, 2010; Chigurupati *et al*, 2010; Christofori, 2011).

The acquisition of invasive properties by cancer cells during tumour progression is closely linked to changes in the expression of adhesion molecules that mediate the interaction of cancer cells with their surrounding cells and extracellular matrix. During tumour progression, malignant cells typically downregulate proteins that mediate adhesion to basal membrane and neighbouring cells and, at the same time, promote the expression of adhesion molecules that facilitate dynamic interaction with extracellular matrix components (Mariotti *et al*, 2007). A paradigmatic example of this phenomenon is the cadherin switch, frequently observed in the epithelial–mesenchymal transition of epithelial cancer cells (Cavallaro *et al*, 2002; Hazan *et al*, 2004). During this transition, E-cadherin expression is downregulated with concomitant enhancement of neuronal cadherin (N-cadherin) expression, which correlates with the acquisition of greater invasiveness and metastatic potential by malignant cells.

Neuronal cadherin (N-cadherin or cadherin-2) is a member of the classical type I cadherin family, which is critically involved in tissue development and maintenance. Like other classical type I cadherins, N-cadherin contains a large N-terminal extracellular domain composed of five tandem-repeated domains, a transmembrane segment

\*Correspondence: Dr J Roma or Dr S Gallego;

E-mail: josep.roma@vhir.org or sgallego@vhebron.net

Received 23 March 2012; revised 26 June 2012; accepted 14 August 2012; published online 13 September 2012

and a cytoplasmatic C-terminal domain able to interact with catenins (Mariotti *et al*, 2007). Although the mechanism by which increased N-cadherin expression promotes malignancy is not fully understood, in a subset of primary tumour cells, it is thought to promote adhesion of malignant cells to N-cadherin-expressing stromal or endothelial cells, thereby facilitating invasion and migration of tumour cells to distant sites (Sandig *et al*, 1997; Qi *et al*, 2005). Several experiments have demonstrated that the expression of N-cadherin suffices to promote invasion and metastasis in some cancers (Nieman *et al*, 1999; Hazan *et al*, 2000; Li *et al*, 2009).

The superfamily of integrins includes several transmembrane cell adhesion proteins that bind to extracellular matrix, cell surface and soluble ligands. All integrins act as heterodimers composed of  $\alpha$ - and  $\beta$ - subunits. The  $\alpha$ - and  $\beta$ -polypeptidic chains are distinct with no detectable homology between them (Takada *et al*, 2007). The heterodimer  $\alpha 9 \beta 1$  has been implicated in several processes that entail dynamic interaction between cells and extracellular matrix, such as re-epithelialisation during cutaneous wound healing (Singh *et al*, 2009), cell adhesion in medulloblastoma and in hematopoietic progenitor cells (Fiorilli *et al*, 2008; Schreiber *et al*, 2009) and finally, cell migration in neoplastic and non-neoplastic cells (Gupta and Vlahakis, 2009; Oommen *et al*, 2011).

The main objective of this work was to demonstrate the existence of a connection between Notch pathway activation and the regulation of mechanisms underlying cell motility and invasiveness in RMS. It has been previously suggested that Notch1 and Notch3 signalling may promote an upregulation of N-cadherin expression in primary melanoma (Liu *et al*, 2006; Wang *et al*, 2007); however, the molecular mechanisms of this possible relationship remain unclear. Moreover, as we demonstrated recently, the Notch pathway mediates RMS cell motility and invasiveness (Roma *et al*, 2011). The oncogenic potential of the Notch pathway was first described in acute T-cell lymphoblastic leukaemia in the late 1980s (Radtke *et al*, 1999). An abnormal upregulation of the Notch pathway has also been reported in ovarian (Park *et al*, 2006), breast (Glahan and Callahan, 1997) and other cancers (Nickoloff *et al*, 2003). With respect to paediatric malignancies, Notch signalling appears to contribute essentially to osteosarcoma metastasis (Zhang *et al*, 2008; Zhang *et al*, 2010) and proliferation (Tanaka *et al*, 2009); Notch signalling also promotes medulloblastoma cancer stem cell survival (Fan *et al*, 2006) and contributes to angiogenesis in neuroblastoma (Funahashi *et al*, 2008). The four Notch receptors are synthesised as precursor proteins that are processed by furin-like enzymes in the trans-Golgi compartment. This cleavage leads to the production of a heterodimeric Notch receptor, which is further processed at the cell surface in a ligand-dependent manner by  $\alpha$ - and  $\gamma$ -secretase. The  $\alpha$ -secretase cleavage occurs at an extracellular site and removes the Notch ectodomain, whereas  $\gamma$ -secretase cleavage occurs in the Notch transmembrane domain, leading to the release of the intracellular Notch fragment (NICD) that translocates to the nucleus where it binds to CSL (CBF1-Su(H)-Lag1) transcription repressors, converting them into transcriptional activators. The targets of these transcription factors in vertebrates are typified by *Hes* and *Hey* genes (Artavanis-Tsakonas *et al*, 1999; Luo *et al*, 2005). The possible demonstration of a connection between Notch signalling and the expression of adhesion molecules may shed light on the unknown molecular mechanism of the Notch-mediated invasive phenotype of RMS cells and allow us to adopt new therapeutic approaches to the fight against metastases in this neoplasia.

## MATERIALS AND METHODS

### Cell culture, transfections and drug treatments

RH30, CW9019 and HTB82 RMS cell lines were grown in MEM medium with Earle's Salts, supplemented with 10% FCS, 2 mM

L-glutamine, 1 mM sodium pyruvate,  $1 \times$  non-essential aminoacids, 100 U ml<sup>-1</sup> penicillin and 0.1 mg ml<sup>-1</sup> streptomycin (all culture reagents supplied by PAA Laboratories, Haidmannweg, Austria) and maintained at 37 °C in a 5% CO<sub>2</sub> water-jacketed incubator. Full-length dnMAML1, Delta and Hes1 complementary DNAs (cDNAs) were generously provided by Dr Anna Bigas (Institut Municipal d'Investigació Mèdica, Barcelona, Spain). The full-length NICD cDNA was a kind gift from Dr Rosanna Paciucci (Institut de Recerca Hospital Vall d'Hebron, Barcelona, Spain). All cDNAs used were cloned into a pCDNA3.1 vector, except dnMAML1 that was cloned into a pEGFP vector. All cDNA transfections were carried out in parallel with the respective empty vector as a control. Transient transfections were performed using 8 µg of plasmid DNA and FuGENE 6 transfection reagent according to the manufacturer's protocol (Roche, Basel, Switzerland). Transfection efficiency was approximately 70% for the three cell lines and was assessed by calculating the percentage of GFP-positive cells referred to the total of cells. Transfected cells were harvested between 48 and 72 h and processed for subsequent assays. Gamma-secretase inhibitor (GSI)-XXI (Calbiochem, San Diego, CA, USA) was diluted in DMSO and added to the culture medium for 3 days at a final concentration of 4 nM as previously described (Roma *et al*, 2011). Control plates were supplemented with an equivalent volume of DMSO.

### Complementary DNA microarrays

Microarrays were carried out using the Affymetrix microarray platform and the Genechip Human Gene 1.0 ST Array; 100 ng of total RNA were used per sample. Two plates of RH30 cells previously treated with GSI were compared with their control (DMSO). Quality of total RNA was assessed by Bioanalyzer Assay (Agilent). Chips were processed on an Affymetrix GeneChip Fluidics Station 450 and Scanner 3000. Results were analysed using TMEV software (Saeed *et al*, 2006).

### Western blotting

Cells were homogenised in lysis buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1 mM PMSF, 1 mM EDTA, 5 mg ml<sup>-1</sup> aprotinin, 5 mg ml<sup>-1</sup> leupeptin, 1% Triton X-100, 1% sodium deoxycholate and 0.1% SDS) and incubated for 4 min at 90 °C. The total protein content was measured using the DC assay kit (Bio-Rad Laboratories, Hercules, CA, USA), a 10% SDS-PAGE was performed and the resolved proteins were in turn transferred onto PVDF membranes (Bio-Rad Laboratories). Membranes were then incubated with anti-Hes1 polyclonal antibody AB5702 (Millipore, Billerica, MA, USA) diluted 1:200, anti-N-cadherin monoclonal antibody C2542 (Sigma, St Louis, MO, USA) diluted 1:1000, anti- $\alpha 9$ -integrin monoclonal antibody clone 3E4 (Abnova, Aachen, Germany) diluted 1:250 and anti-Notch1 (C-20) sc-6014 (Santa Cruz Biotechnology, Santa Cruz, CA, USA) diluted 1:500. An anti- $\alpha$ -tubulin monoclonal antibody (Cell Signaling, Danvers, MA, USA) diluted 1:2000 was used as a loading control.

### Immunofluorescence

Each well was plated with  $3 \times 10^5$  cells on 12-mm round glass coverslips pre-coated with a 4% collagen-I solution (BD Biosciences, Erembodegem, Belgium). After 24 h, cells were transfected with the appropriate plasmids as described above. After 48 h, cells were washed with PBS, fixed in 4% paraformaldehyde, incubated with 50 mM NH<sub>4</sub>Cl and permeabilised with 0.1% Triton X-100. Samples were then blocked with 5% FCS prior to incubation with the following primary antibodies: anti-Hes1 AB5702 (Millipore) diluted 1:50, anti-N-cadherin C2542 (Sigma) diluted 1:250 and anti- $\alpha 9$ -integrin clone 3E4 (Abnova) diluted 1:125. Cells were then incubated with suitable secondary antibodies labelled with

AlexaFluor-647 or AlexaFluor-488 (Molecular probes, Eugene, OR, USA). Nuclei were stained with  $5 \text{ ng ml}^{-1}$  of Hoechst33342 (Sigma). Fluorescence was visualised in a FV1000 confocal microscope (Olympus, Hamburg, Germany) and quantified with Olympus Fluorview software. A minimum of 100 cells obtained from random selected fields was evaluated.

### Immunohistochemistry

Patient samples ( $n=9$ ) were obtained from the Vall d'Hebron Hospital Pathology Department (kindly provided by Dr Nuria Toran). Paraffin-embedded tissues were sliced, deparaffinised and rehydrated, followed by antigen retrieval in 10 mM citrate buffer (DAKO). Endogenous peroxidase activity was quenched using 1% hydrogen peroxide. Samples were blocked and incubated with antibodies against Hes1 (AB5702, Millipore), N-cadherin (C2542, Sigma) and  $\alpha 9$ -integrin (3E4, Abnova), all diluted 1:200. After incubation with peroxidase-conjugated secondary antibody and peroxidase substrate, samples were counterstained with hematoxylin for 10 s, dehydrated and mounted. Labelling of each antibody was evaluated using a semiquantitative method regarding the percentage of positive cells ( $\leq 25\%$ , 26–50%, 51–75% and  $> 75\%$  corresponding to a score of 1, 2, 3 and 4 respectively) and intensity of the labelling (0–3). The assigned final score (1–7) corresponds to the sum of the labelling and percentage of positive cell scores.

### RNA isolation, retrotranscription and real-time PCR

RNA was isolated using the Quick-prep micro RNA isolation kit following the manufacturer's instructions (Qiagen, Valencia, CA, USA). Samples of  $2 \mu\text{g}$  of total RNA were reverse transcribed using random primers (Invitrogen, Carlsbad, CA, USA). The reaction mixture was incubated for 1 h at  $37^\circ\text{C}$  with 200 U of Moloney murine leukaemia virus reverse transcriptase (Promega, Madison, WI, USA). A 40-cycle PCR based on the TaqMan assay (Applied Biosystems, Foster City, CA, USA) was performed to detect N-cadherin and  $\alpha 9$ -integrin (Hs00983056\_m1 and Hs00979865\_m1 TaqMan assays, respectively). The downstream Notch effector Hes1 (assay Hs00172878\_m1) was used to assess Notch activation. The housekeeping gene *TBP* (assay Hs00172424\_m1) was used as internal control. Quantification of relative levels of each mRNA analysed was performed by the method of (Livak and Schmittgen, 2001). All samples were tested in triplicate.

### Fibronectin and collagen adhesion assay

A total of  $3 \times 10^5$  GSI-pre-treated cells per well was seeded in a 24-well plate previously coated with collagen or fibronectin. Cells were then incubated at  $37^\circ\text{C}$  for 15 min (RH30), 30 min (CW9019) or 1 h (HTB82), respectively, fixed with 4% paraformaldehyde and stained with 0.2% crystal violet. Cells were then lysed in 10% acetic acid and the absorbance at 590 nm was determined as a value proportional to the number of cells adhered to the substrate. To establish the specificity of the assay for N-cadherin and  $\alpha 9$ -integrin, some wells were treated with  $20 \mu\text{g ml}^{-1}$  of either N-cadherin- (GC-4, Sigma) or  $\alpha 9$ -integrin-blocking antibodies (Y9A2, Chemicon International, Temecula, CA, USA). The control wells were treated with an anti- $\alpha$ -tubulin monoclonal antibody (Cell Signaling) at the same concentration, as isotypic control. All measurements were taken in triplicate.

### Transwell assay

Cells were trypsinised, seeded at  $10^5$  cells per well and incubated in the presence or absence of  $20 \mu\text{g ml}^{-1}$  of either N-cadherin or  $\alpha 9$ -integrin-blocking antibodies for 30 min on ice. Following the incubation, cells were plated in the upper chamber previously coated with BD Matrigel (BD Biosciences) in an 8- $\mu\text{m}$  pore-size transwell (Corning). Basal medium containing 1% FBS in the

presence or absence of the aforementioned N-cadherin and  $\alpha 9$ -integrin-blocking antibodies was added to each well and incubated at  $37^\circ\text{C}$  for 6, 12 and 48 h (for RH30, CW9019 and HTB82 cell lines, respectively). The control wells were treated with  $20 \mu\text{g ml}^{-1}$  anti- $\alpha$ -tubulin monoclonal antibody (Cell Signaling). Remaining cells were removed from the upper chamber with a cotton swab, and cells migrated to the lower surface were stained with 0.2% crystal violet, lysed in 10% acetic acid and the absorbance at 590 nm was determined as a value proportional to the number of cells on the lower surface of each membrane. All analyses were made in triplicate.

### Wound-healing assay

Cells were seeded on a 24-well plate ( $3 \times 10^5$  cells per well) and incubated in the presence or absence of  $20 \mu\text{g ml}^{-1}$  of either N-cadherin- or  $\alpha 9$ -integrin-blocking antibodies as described above. The control wells were treated with  $20 \mu\text{g ml}^{-1}$  anti- $\alpha$ -tubulin monoclonal antibody as isotypic control (Cell Signaling). A day later, cell monolayers were scratched with a pipette tip and placed in complete growth medium with or without the suitable blocking antibody. The initial scratched areas were measured using the ImageJ software (NIH, freely available at <http://rsb.info.nih.gov/ij/>) and serial images of random selected fields were acquired until the healing of the wounds was complete. Healing velocity was calculated as a quotient between the scratched area and the time required for the wound to heal. All analyses were made in triplicate.

### Chromatin immunoprecipitation (ChIP) assay

Chromatin immunoprecipitation was performed following the manufacturer's (Millipore) instructions using  $5 \mu\text{g}$  of polyclonal rabbit antibodies against Hes1 AB5702 (Millipore) and Notch1 sc-6014 (Santa Cruz Biotechnology). The same amount of rabbit immunoglobulin (IgG) was used as a control (Abcam). The co-immunoprecipitated DNAs were used for PCR to amplify N-cadherin and  $\alpha 9$ -integrin promoter regions. The primers used were as follows: N-cadherin promoter (forward 5'-ACCCAGAGA TCAAGGAGCTG-3'; reverse 5'-CTCCACTTCCACCTCCACAT-3') and  $\alpha 9$ -integrin promoter (forward 5'-GAGCTCAAAAGTGCCT CTC-3'; reverse 5'-TGAGGGAGGAAAAAGAAGCA-3').

## RESULTS


### Pharmacological inhibition of Notch signalling impairs cell adhesion in RMS cell lines

Cells treated with GSI showed a clear reduction in cell adhesiveness detected in either collagen- or fibronectin-coated plates (Figure 1). The effect produced by GSI-XXI on cell adhesiveness varied among the cell lines analysed and according to the coating of the plate surface; however, in general, treated cells showed a significant reduction in adhesion that ranged from 20 to 50% in fibronectin-coated plates (Figure 1A) and from 30 to 60% in collagen-coated plates (Figure 1B). Cell viability was assessed by flow cytometry with no substantial differences observed between GSI-treated and untreated (DMSO) cells (Supplementary Figure S1). These observations fitted our hypothesis that Notch pathway may be able to regulate some of the molecular processes that lead to cell adhesion and migration. However, they did not rule out the possibility that the effects observed may have been caused by effects of the drug on other pathways or directly on cell adhesion molecules.

### Pharmacological inhibition of Notch signalling significantly reduces N-cadherin and $\alpha 9$ -integrin expression

A cDNA microarray permitted preliminary identification of adhesion proteins downregulated under GSI treatment. Neuronal





**Figure 1** Cell adhesion to fibronectin- or collagen-coated plates. Images showing a representative field of attached cells for each cell line in the presence (GSI) or absence (DMSO) of  $\gamma$ -secretase inhibitor and plots representing the quantification of cells attached to fibronectin- (**A**) or collagen-coated (**B**) plates. All samples were evaluated in triplicate. Student's *t*-test significance: \* $P < 0.05$ , \*\* $P < 0.005$ .


cadherin and  $\alpha 9$ -integrin were selected for their downregulation under GSI treatment, their significant expression (other proteins downregulated but barely expressed in control samples were ruled out) and their previously well-established involvement in cell adhesion, motility and metastasis (Figure 2A). With the aim of confirming the results obtained in the cDNA microarray, we specifically analysed the expression of the two selected genes by western blotting and qRT-PCR. Western blotting (Figure 2B) revealed a reduction in both full-length N-cadherin and the CTF2 fragment levels. The CTF1 fragment was not detected in the RMS cell lines analysed (not shown), thereby suggesting a negligible role of this isoform in RMS. Levels of  $\alpha 9$ -integrin were also reduced in cells treated with GSI. Levels of  $\beta 1$ -integrin, the usual partner of  $\alpha 9$ -integrin, showed no changes (not shown). The reduction in Hes1, as a marker of effective Notch inhibition in treated cells, is also shown in Figure 2B. Similarly, mRNA levels of both N-cadherin and  $\alpha 9$ -integrin showed statistically significant reductions in the three cell lines, down to between 60 and 85% for N-cadherin and to 20–50% for  $\alpha 9$ -integrin, compared with untreated cells (100%). The mRNA expression of Hes1 was also analysed as a marker of correct Notch inhibition and showed a

statistically significant reduction, as expected, in cells treated with GSI (Figure 2C). Although pharmacological Notch inhibition was not complete, reductions in N-cadherin and  $\alpha 9$ -integrin expressions were significant, particularly for  $\alpha 9$ -integrin.

Although these results confirmed the N-cadherin and  $\alpha 9$ -integrin reduction in cells treated with GSI, they failed to demonstrate unequivocally that the effects observed could be exclusively Notch-dependent.

#### Genetic modification of Notch pathway activation gives rise to a parallel regulation of N-cadherin and $\alpha 9$ -integrin

Seeking to rule out the possibility of the observed effects on cells treated with Notch inhibitors being attributable to the involvement of other pathways or to a direct effect of the drugs on N-cadherin or  $\alpha 9$ -integrin – and therefore demonstrate that these effects resulted from specific Notch activation – we genetically manipulated Notch signalling activity by transfecting constructs containing the Notch ligand Delta1 (Delta) and a dominant negative form of MAML1 (dnMAML1). Activation of the Notch pathway by transfection of Delta promotes N-cadherin and  $\alpha 9$ -integrin



**Figure 2** Pharmacological inhibition of Notch signalling significantly reduced N-cadherin and  $\alpha 9$ -integrin expression. **(A)** Complementary DNA (cDNA) expression microarray showing the relative levels of  $\alpha 9$ -integrin (ITGA9), N-cadherin (N-Cad) and Hes1 expression in GSI-treated samples (GSI) compared with the control (DMSO). **(B)** Panel of western blots showing the effects of GSI on full-length N-cadherin (N-Cad/FL) and CTF2 fragment,  $\alpha 9$ -integrin (ITGA9), NICD and Hes1 levels compared with untreated samples (DMSO). Alpha-tubulin was used as a loading control. Below the western blot composition, a plot depicting densitometric analysis of each band of the western blot. **(C)** Relative mRNA levels evaluated by quantitative PCR in the presence (GSI) or absence (DMSO) of  $\gamma$ -secretase inhibitor. Neuronal cadherin,  $\alpha 9$ -integrin and Hes1 mRNA levels were evaluated in triplicate. Student's *t*-test significance: \**P* < 0.05, \*\**P* < 0.005.

increases in both protein and mRNA levels (Figures 3A and C). Conversely, inhibition of the pathway by overexpression of the dominant negative dnMAML1 produces a parallel reduction in protein and mRNA levels of N-cadherin and  $\alpha 9$ -integrin (Figures 3B and C). Western blot revealed the presence of full-length N-cadherin (N-Cad FL) and the CTF2 fragment, whereas the CTF1 fragment was not detected (not shown). Both forms of N-cadherin detected (N-Cad FL and CTF2) and  $\alpha 9$ -integrin showed induction when the Notch pathway was activated (overexpression of Delta) and a clear reduction when repressed (overexpression of dnMAML1) (Figures 3A and B). No changes were observed in the amount of  $\beta 1$ -integrin, the usual partner of  $\alpha 9$ -integrin (not shown).

### Immunofluorescence


Fluorescent immunocytochemistry against N-cadherin revealed moderate expression of this protein in non-transfected cells (Figures 4A and C). When cells underwent overactivation of the Notch pathway (by Delta1 transfection), N-cadherin expression was significantly increased in the three cell lines analysed and conserved the localisation pattern of the endogenous protein in each cell line. The staining was localised in cytoplasm and cell membrane in RH30 and CW9019 cell lines, but more prominently

in the nucleus in HTB82, which suggests a major presence of the CTF2 form in this cell line (also observed by western blot, Figure 3).

The distribution of  $\alpha 9$ -integrin staining was general and showed prominent nuclear and cytoplasmic staining. As in the results found with N-cadherin transfections, RMS cells transfected with Delta also showed a statistically significant increase in  $\alpha 9$ -integrin staining compared with controls (Figures 4B and C).

### Specific anti-N-cadherin- and anti- $\alpha 9$ -integrin-blocking antibodies reduce invasive properties of RMS cells

Seeking to demonstrate that the role of N-cadherin and  $\alpha 9$ -integrin is critical for the Notch-mediated increase in RMS cell capability to migrate and invade, we performed adhesion, wound-healing and invasion assays in the presence or absence of specific blocking antibodies against these two proteins. The main aim of these experiments was to demonstrate whether inhibition of N-cadherin and/or  $\alpha 9$ -integrin suffices to compensate for the increases in adhesion, wound-healing and invasiveness detected in cells with constitutive Notch activation. Cells transfected with Delta or NICD showed an increase in their adhesion to fibronectin-coated plates, which could be reverted using either anti-N-cadherin- or anti- $\alpha 9$ -integrin-blocking antibodies (Figure 5A). The effects of the



**Figure 3** Genetic modifications of the Notch pathway influenced N-cadherin and  $\alpha 9$ -integrin expression. Western blots showing the concomitant variations of Notch pathway activation (NICD and Hes1) and N-cadherin (N-Cad/FL and CTF2) and  $\alpha 9$ -integrin (ITGA9) expression in RMS cells transfected with Delta (**A**) or dnMAML1 (**B**). Below each western blot composition, a plot depicting densitometric analysis of each band of the western blot. The variations associated with genetic modification of the Notch pathway (Delta and MAML1 transfections) observed in protein are also seen in the mRNAs of N-cadherin,  $\alpha 9$ -integrin and Hes1 determined by quantitative PCR (**C**). All RNA measurements were taken in triplicate. Student's *t*-test significance: \* $P < 0.05$ , \*\* $P < 0.005$ .

blocking antibodies on healing velocity differed for N-cadherin and  $\alpha 9$ -integrin. Thus, while N-cadherin-blocking antibody gave rise to a significant reduction in healing velocity in the three cell lines analysed, the anti- $\alpha 9$ -integrin-blocking antibody did not promote any significant decrease in the capability of cells to heal a wound (Figure 5B). Finally, the effects observed with both antibodies on cell invasiveness in a matrigel/transwell assay also differed (Figure 5C). Interestingly, the  $\alpha 9$ -integrin-blocking antibody produced marked invasiveness impairment in this assay, whereas only a very weak effect of N-cadherin-blocking antibody on invasiveness was observed.

#### Correlation among Hes1, N-cadherin and $\alpha 9$ -integrin in RMS archival tumour samples

Patient sections were stained using antibodies against Hes1, N-cadherin and  $\alpha 9$ -integrin with the aim of demonstrating the correlation among these proteins observed in cultured RMS cell lines. The intensity of labelling was evaluated using a semiquantitative method as described above. The results obtained showed a correlation between the expression of Hes1 with both N-cadherin and  $\alpha 9$ -integrin (Figure 6). Although the number of samples was limited ( $n = 9$ ), the correlation between Hes1 and N-cadherin yielded a Pearson correlation of 0.584 ( $P < 0.1$ ) and the correlation

for Hes1 and  $\alpha 9$ -integrin a value of 0.771 ( $P < 0.05$ ), thereby suggesting a dependency of Hes1 expression on N-cadherin and  $\alpha 9$ -integrin. This correlation among these proteins confirms, in patients, the aforementioned findings in cultured cells and suggests that Notch activation may also influence the expression of these two adhesion proteins in tumours.


#### NICD and Hes1 bind to N-cadherin and $\alpha 9$ -integrin promoters

Chromatin immunoprecipitation assay revealed co-immunoprecipitation of both NICD and Hes1 with the N-cadherin and  $\alpha 9$ -integrin promoter sequences. The control IgGs rendered no amplification (Figure 7). These results pointed to a direct binding of these Notch downstream effectors to the promoter regions of N-cadherin and  $\alpha 9$ -integrin genes.

#### DISCUSSION

Rhabdomyosarcoma patients with metastatic disease continue to have poor prognosis and the major cause of death in these patients is the formation of distant metastases. The identification of molecular components and mechanisms that control metastasis in






**Figure 4** N-cadherin and  $\alpha 9$ -integrin immunocytochemistry. Immunocytochemistry with anti-Hes1 and N-cadherin (N-Cad) antibodies (**A**) or Hes1 and  $\alpha 9$ -integrin (ITGA9) antibodies (**B**) in cells transfected with Delta or control vector (CTRL). Staining was as follows: Hes1 (green), N-cadherin and  $\alpha 9$ -integrin (red) and nuclei (blue). (**C**) Relative fluorescence quantification for Hes1, N-cadherin and  $\alpha 9$ -integrin. Student's *t*-test significance: \**P* < 0.05.

RMS may be useful for developing targeted therapies focused on reducing metastasis in this neoplasia. Although the mechanism by which increased N-cadherin expression promotes malignancy is not fully understood, it is thought to promote adhesion of malignant cells to N-cadherin-expressing stromal or endothelial cells, thereby facilitating invasion and migration of tumour cells to distant sites (Sandig *et al*, 1997; Qi *et al*, 2005). Although several works have demonstrated that N-cadherin expression suffices to promote metastasis in some cancers (Nieman *et al*, 1999; Hazan *et al*, 2000; Li *et al*, 2009) and an aberrant N-cadherin expression has been described in RMS (Soler *et al*, 1993), no previous publications have shown the importance of N-cadherin for RMS invasiveness. Herein, a central role of N-cadherin in RMS cell adhesion and motility is demonstrated; furthermore, a new mechanism of N-cadherin induction, mediated by Notch pathway activation, is also proposed. Likewise,  $\alpha 9$ -integrin – forming a heterodimer with  $\beta 1$ -integrin – has previously been implicated in processes, such as cell migration, involving dynamic interaction among cells and their microenvironment (Gupta and Vlahakis, 2009; Oommen *et al*, 2011). However, its possible role in RMS remains unknown. Therefore, this work represents the first description of a pro-invasive role of  $\alpha 9$ -integrin in RMS.

We and others recently implicated the Notch pathway in paediatric sarcoma motility and invasiveness (Zhang *et al*, 2008;

Roma *et al*, 2011); however, the molecular connection between the Notch pathway and the process that leads to cell adhesion remains largely obscure. A previous work described the requirement of Notch activation in the epithelial-to-mesenchymal transition of epithelial cells, which finally rendered a disassembly of E-cadherin junctions, thereby permitting an increase in cell motility (Zavadil *et al*, 2004). The results herein presented point to the existence of a Notch-mediated mechanism able to positively modulate the expression of adhesion proteins, such as N-cadherin and  $\alpha 9$ -integrin, which may drive motility and invasiveness in neoplastic cells, thereby permitting cancer progression. The three cell lines studied belong to different RMS subtypes and are known to have great biological differences. Two contain PAX/FOXO1 translocations and one bears no translocation. The fact that we consistently found a relationship between Notch and N-cadherin and  $\alpha 9$ -integrin in such different cell lines suggests that the mechanism here described is a general mechanism in RMS – and perhaps in other neoplasias – and not a mere characteristic of a particular RMS subtype. Therefore, Notch-controlled changes in the levels of proteins implicated in tumour cell spread, such as a down-regulation of E-cadherin in epithelial tumours (Zavadil *et al*, 2004) and an upregulation of N-cadherin and  $\alpha 9$ -integrin in a mesenchymal tumour such as RMS, support the role of Notch as an upstream key signalling pathway in the control of cancer




**Figure 5** Anti-N-cadherin- and anti- $\alpha 9$ -integrin-blocking antibodies reduced invasive properties induced by Notch activation. Three RMS cell lines were transfected with the constructions indicated below the bottom of each bar (pCDNA, Delta) in the presence (+) or absence (–) of either N-cadherin-blocking antibody (N-Cad Ab) or  $\alpha 9$ -integrin-blocking antibody ( $\alpha 9$ -ITG Ab) as indicated in the figure. **(A)** Adhesion assay on fibronectin-coated plates. The percentage of cells attached was quantified per cell line and condition. **(B)** Wound-healing assay. All bars indicate the healing velocity of each cell line, transfection and treatment **(C)** Matrigel/Transwell assay. Invasiveness is represented as a percentage of cells in the lower compartment, compared with the control. The three assays were performed in three independent wells per cell line and condition. Student's *t*-test significance: \**P* < 0.05.

dissemination. Furthermore, the promoter-binding study, herein provided, pointed to a direct interaction of Hes1 and NICD to N-cadherin and  $\alpha 9$ -integrin gene regulatory regions. NICD and Hes1 are known to be proteins able to bind directly (Hes1) or indirectly (NICD) to DNA, thereby influencing gene transcription. The binding of NICD to the N-cadherin and  $\alpha 9$ -integrin promoter regions strongly suggests a role in their transcriptional activation, as NICD is an archetypal transcriptional activator. On the other hand, Hes1 is a known transcriptional repressor; however, it may also act as a transcriptional activator of certain genes (Yan *et al*, 2002). It is not clear whether Hes1 can exert a repressive or activating role in the promoters of the two genes included in this study. It is also conceivable that the relative amounts of both NICD and Hes1 may cooperatively regulate – inducing or repressing, depending on their relative proportion – the expression of the N-cadherin and  $\alpha 9$ -integrin genes.


The use of pharmacological Notch pathway inhibitors, such as GSIs, has been proposed as a therapeutic alternative in several cancers, focused essentially on reducing cell proliferation. Several GSIs have been actively studied as potential inhibitors of the generation of the  $\beta$ -amyloid peptide associated with Alzheimer's disease (Selkoe and Kopan, 2003). More recently, some GSIs have begun to be studied in phase I–II trials for patients with advanced breast cancer and acute T-cell leukaemias (Deangelo *et al*, 2006; Krop *et al*, 2006). Beyond the use of GSIs as anti-proliferation agents, the results of our experiments point to GSIs as compounds able to reduce the invasive features of tumour cells and, therefore, suitable for use in the fight against metastasis.

Taken together, the results of this work reveal that the Notch pathway has a significant role in the regulation of N-cadherin and  $\alpha 9$ -integrin expression in RMS and, therefore, the Notch pathway should be considered a key regulator of the mechanisms, such as





**Figure 6** The expression of Hes1, N-cadherin and  $\alpha 9$ -integrin correlate in archival RMS tumour samples. **(A)** Representative sections of three patients (34, 37 and 41) stained with anti-Hes1, N-Cad and  $\alpha 9$ -integrin antibodies. **(B and C)** Plots depicting the correlations between Hes1 and both N-cadherin and  $\alpha 9$ -integrin semiquantitative scores. Pearson correlation value and significance are shown in the upper left corner in each plot.



**Figure 7** Binding of NICD and Hes1 to N-cadherin and  $\alpha 9$ -integrin promoters. Lysates from RH30 cells were analysed by ChIP assay using specific antibodies against Hes1, NICD and rabbit IgG as a control. Immunoprecipitated chromatin was amplified by PCR using N-cadherin and  $\alpha 9$ -integrin promoter region-specific primers. IgG: negative control rabbit IgG; Input: 5% of the sonicated cell lysate used for IP; H<sub>2</sub>O: PCR negative control; M: 100 bp DNA ladder.

cell motility and invasiveness, which lead to tumour progression. Hence, Notch-inhibiting molecules can be considered as possible therapeutic agents against metastasis. Furthermore, the importance of N-cadherin and  $\alpha 9$ -integrin as players directly involved in cell attachment, migration and invasiveness in RMS is also

demonstrated, consequently pointing to these proteins as new candidates for target-specific therapies focused on reducing metastasis in this neoplasia.

## ACKNOWLEDGEMENTS

The authors wish to thank Ms Marta Rebull and Mr Isaac Vidal for laboratory technical support, Ms Marta Valeri for technical support with the *in vivo* wound-healing assay and confocal microscopy, Ms Paqui Gallego for technical support with the real-time PCR assays, Mr Ricardo Gonzalo for cDNA microarrays hybridisation and Ms Christine O'Hara for help with the English version of this manuscript. This work was supported by grants from Institut Català d'Oncologia (ICO), Instituto de Salud Carlos III (RD06/0020/1021 and PI11/00740), Fundació la Marató de TV3, Asociación Española Contra el Cáncer, Fundació SMALL and Fundació A. BOSCH.

## Conflict of interest

The authors declare no conflict of interest.

Supplementary Information accompanies the paper on British Journal of Cancer website (<http://www.nature.com/bjc>)

## REFERENCES

- Artavanis-Tsakonas S, Rand MD, Lake RJ (1999) Notch signaling: cell fate control and signal integration in development. *Science* **268**: 225–232
- Barr FG, Gallili N, Holich J, Biegel JA, Rovera G, Emanuel BS (1993) Rearrangement of the PAX3 paired box gene in the paediatric solid tumour alveolar rhabdomyosarcomas. *Nat Genet* **3**: 113–117
- Cavallaro U, Schaffhausen B, Christofori G (2002) Cadherins and the tumour progression: is it all in a switch? *Cancer Lett* **176**(2): 123–128
- Chigurupati S, Venkataraman R, Barrera D, Naganathan A, Madan M, Paul L, Pattisapu JV, Kyriazis GA, Sugaya K, Bushnev S, Lathia JD, Rich JN, Chan SL (2010) Receptor channel TRPC6 is a key mediator of Notch-driven glioblastoma growth and invasiveness. *Cancer Res* **70**(1): 418–427
- Christofori G. Metastatic colon cancer cells negotiate the intravasation notch (2011) *Cancer Cell* **19**(1): 6–8
- Davis RJ, D'Cruz CM, Lowell MA, Biegel JA, Barr FG (1994) Fusion of PAX7 to the FOXO1 by the variant t(1;13) (p36;q14) translocation in alveolar rhabdomyosarcomas. *Cancer Res* **54**: 2869–2872
- Deangelo DJ, Stone RM, Silverman LB, Stock W, Attar EC, Fearon I, Dallob A, Matthews C, Stone J, Freedman SJ, Aster J (2006) A phase I clinical trial of the Notch inhibitor MK-0752 in patients with T-cell acute lymphoblastic leukemia/lymphoma (T-ALL) and other leukemias. *J Clin Oncol*. ASCO Annual meeting Proceedings **24**(18S): abstract 6585
- Engin F, Bertin T, Ma O, Jiang MM, Wang L, Sutton RE, Donehower LA, Lee B (2009) Notch signaling contributes to the pathogenesis of human osteosarcomas. *Hum Mol Genet* **18**(8): 1464–1470
- Fan X, Matsui W, Khaki L, Stearns D, Chun J, Li YM, Eberhart CG (2006) Notch pathway inhibition depletes stem-like cells and blocks engraftment in embryonal brain tumors. *Cancer Res* **66**: 7445–7452
- Fiorilli P, Partridge D, Staniszewska I, Wang JY, Grabacka M, So K, Marcinkiewicz C, Reiss K, Khalili K, Croul SE (2008) Integrins mediate adhesion of medulloblastoma cells to tenascin and activate pathways associated with survival and proliferation. *Lab Invest* **88**(11): 1143–1156
- Funahashi Y, Hernandez SL, Das I, Ahn A, Huang J, Vorontchikhina M, Sharma A, Kanamaru E, Borisenko V, Desilva DM, Suzuki A, Wang X, Shawber CJ, Kandel JJ, Yamashiro DJ, Kitajewski J (2008) A Notch1 ectodomain construct inhibits endothelial notch signaling, tumor growth, and angiogenesis. *Cancer Res* **68**: 4727–4735
- Glahan D, Callahan R (1997) The mouse mammary tumor associated gene INT3 is a unique member of the NOTCH gene family (NOTCH4). *Oncogene* **14**: 1883–1890
- Gupta SK, Vlahakis NE (2009) Integrin alpha9beta1 mediates enhanced cell migration through nitric oxide synthase activity regulated by Src tyrosine kinase. *J Cell Sci* **122**(12): 2043–2054
- Hazan RB, Phillips GR, Qiao RF, Norton L, Aaronson SA (2000) Exogenous expression of N-Cadherin in breast cancer cells induces cell migration, invasion, and metastasis. *J Cell Biol* **148**(4): 779–790
- Hazan RB, Qiao R, Keren R, Badano I, Suyama K (2004) Cadherin switch in tumor progression. *Ann N Y Acad Sci* **1014**: 155–163
- Krop IE, Kosh M, Fearon I, Savoie J, Dallob A, Matthews C, Stone J, Winer E, Freedman SJ, Lorusso P (2006) Phase I pharmacokinetic (PK), and pharmacodynamic (PD) trial of the novel oral Notch inhibitor MK-0752 in patients (pts) with advanced breast cancer (BC) and other solid tumors. *J Clin Oncol*. ASCO Annual meeting Proceedings **24**(18S): abstract 10574
- Li K, He W, Lin N, Wang X, Fan QX (2009) N-Cadherin knock-down decreases invasiveness of esophageal squamous cell carcinoma in vitro. *World J Gastroenterol* **15**(6): 697–704
- Liu ZJ, Xiao M, Balint K, Smalley KS, Brafford P, Qiu R, Pinnix CC, Li X, Herlyn M (2006) Notch1 signaling promotes primary melanoma progression by activating mitogen-activated protein kinase/phosphatidylinositol 3-kinase-Akt pathways and up-regulating N-cadherin expression. *Cancer Res* **66**(8): 4182–4190
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2-DDCT method. *Methods* **25**(4): 402–408
- Luo D, Renault VM, Rando TA (2005) The regulation of notch signaling in muscle stem cell activation and postnatal myogenesis. *Semin Cell Dev Biol* **16**: 612–622
- Mariotti A, Perotti A, Sessa C, Ruegg C (2007) N-Cadherin as a therapeutic target in cancer. *Expert Opin Investig Drugs* **16**(4): 451–465
- Nickoloff BJ, Osborne BA, Miele L (2003) Notch signaling as a therapeutic target in cancer: a new approach to the development of cell fate modifying agents. *Oncogene* **22**: 6598–6608
- Nieman MT, Prudoff RS, Johnson KR, Wheelock MJ (1999) N-Cadherin promotes motility in human breast cancer cells regardless of their E-cadherin expression. *J Cell Biol* **147**(3): 631–644
- Oommen S, Gupta SK, Vlahakis NE (2011) Vascular endothelial growth factor A (VEGF-A) induces endothelial and cancer cell migration through direct binding to integrin alpha9beta1: identification of a specific alpha9beta1 binding site. *J Biol Chem* **286**(2): 1083–1092
- Pang RT, Leung CO, Ye TM, Liu W, Chiu PC, Lam KK, Lee KF, Yeung WS (2010) MicroRNA-34a suppresses invasion through down-regulation of Notch1 and Jagged1 in cervical carcinoma and choriocarcinoma cells. *Carcinogenesis* **31**(6): 1037–1044
- Park JT, Li M, Nakayama K, Mao TL, Davidson B, Zhang Z, Kurman RJ, Eberhart CG, IeM Shih, Wang TL (2006) Notch3 gene amplification in ovarian cancer. *Cancer Res* **66**: 6312–6318
- Qi J, Chen N, Wang J, Siu CH (2005) Transendothelial migration of melanoma cells involves N-cadherin-mediated adhesion and activation of the beta-catenin signaling pathway. *Mol Biol Cell* **16**(9): 4386–4397
- Radtko F, Wilson A, Stark G, Bauer M, van Meerwijk J, MacDonald HR, Aguet M (1999) Deficient T cell fate specification in mice with an induced inactivation of Notch1. *Immunity* **10**: 547–558
- Roma J, Masià A, Reventós J, Sánchez de Toledo J, Gallego S (2011) Notch pathway inhibition significantly reduces rhabdomyosarcoma invasiveness and mobility in vitro. *Clin Cancer Res* **17**(3): 505–513
- Saeed AI, Bhagabati NK, Braisted JC, Liang W, Sharov V, Howe EA, Li J, Thiagarajan M, White JA, Quackenbush J (2006) TM4 microarray software suite. *Methods Enzymol* **411**: 134–193
- Sandig M, Voura EB, Kalnins VI, Siu CH (1997) Role of cadherins in the transendothelial migration of melanoma cells in culture. *Cell Motil Cytoskeleton* **38**(4): 351–364
- Schreiber TD, Steinhilber C, Essl M, Abele H, Geiger K, Müller CA, Aicher WK, Klein G (2009) The integrin alpha9beta1 on hematopoietic stem and progenitor cells: involvement in cell adhesion, proliferation and differentiation. *Haematologica* **94**(11): 1493–1501
- Selkoe D, Kopan R (2003) Notch and presenilin: regulated intramembrane proteolysis links development and degeneration. *Annu Rev Neurosci* **26**: 565–597
- Singh P, Chen C, Pal-Ghosh S, Stepp MA, Sheppard D, Van De Water L (2009) Loss of integrin alpha9beta1 results in defects in proliferation, causing poor re-epithelialization during cutaneous wound healing. *J Invest Dermatol* **129**(1): 217–228
- Soler AP, Johnson KR, Wheelock MJ, Knudsen KA (1993) Rhabdomyosarcoma-derived cell lines exhibit aberrant expression of the cell-cell adhesion molecules N-CAM, N-Cadherin, and cadherin-associated proteins. *Exp Cell Res* **208**(1): 84–93
- Takada Y, Ye X, Simon S (2007) The integrins. *Genome Biol* **8**(5): 215
- Tanaka M, Setoguchi T, Hirotsu M, Gao H, Sasaki H, Matsunoshita Y, Komiya S (2009) Inhibition of Notch pathway prevents osteosarcoma growth by cell cycle regulation. *Br J Cancer* **100**: 1957–1965
- Wang T, Holt CM, Xu C, Ridley C, P O Jones R, Baron M, Trump D (2007) Notch3 activation modulates cell growth behaviour and cross-talk to Wnt/TCF signaling pathway. *Cell Signal* **19**(12): 2458–2467
- Yan B, Raben N, Plotz PH (2002) Hes-1, a known transcriptional repressor, acts as a transcriptional activator for the human acid alpha-glucosidase gene in human fibroblast cells. *Biochem Biophys Res Commun* **291**(3): 582–587
- Zavadij I, Cermak L, Soto-Nieves N, Böttinger EP (2004) Integration of TGF-beta/Smad and Jagged1/Notch signalling in epithelial-to-mesenchymal transition. *EMBO J* **23**(5): 1155–1165
- Zayzafoon M, Abdulkadir SA, McDonald JM (2004) Notch signaling and ERK activation are important for the osteomimetic properties of prostate cancer bone metastatic cell lines. *J Biol Chem* **279**(5): 3662–3670
- Zhang P, Yang Y, Nolo R, Zweidler-McKay PA, Hughes DP (2010) Regulation of NOTCH signaling by reciprocal inhibition of HES1 and Deltex1 and its role in osteosarcoma invasiveness. *Oncogene* **29**(20): 2916–2926
- Zhang P, Yang Y, Zweidler-McKay PA, Hughes DP (2008) Critical role of notch signalling in osteosarcoma invasion and metastasis. *Clin Cancer Res* **14**(10): 2962–2969

This work is published under the standard license to publish agreement. After 12 months the work will become freely available and the license terms will switch to a Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported License.