

Keywords: physical activity; renal cancer; meta-analysis

# The association between physical activity and renal cancer: systematic review and meta-analysis

G Behrens<sup>1</sup> and M F Leitzmann<sup>1</sup>

<sup>1</sup>Department of Epidemiology and Preventive Medicine, Regensburg University Medical Center, Franz-Josef-Strauss-Allee 11, 93053 Regensburg, Germany

**Background:** Physical activity may decrease renal cancer risk by reducing obesity, blood pressure, insulin resistance, and lipid peroxidation. Despite plausible biologic mechanisms linking increased physical activity to decreased risk for renal cancer, few epidemiologic studies have been able to report a clear inverse association between physical activity and renal cancer, and no meta-analysis is available on the topic.

**Methods:** We searched the literature using PubMed and Web of Knowledge to identify published non-ecologic epidemiologic studies quantifying the relationship between physical activity and renal cancer risk in individuals without a cancer history. Following the PRISMA guidelines, we conducted a systematic review and meta-analysis, including information from 19 studies based on a total of 2327322 subjects and 10756 cases. The methodologic quality of the studies was examined using a comprehensive scoring system.

**Results:** Comparing high vs low levels of physical activity, we observed an inverse association between physical activity and renal cancer risk (summary relative risk (RR) from random-effects meta-analysis=0.88; 95% confidence interval (CI)=0.79–0.97). Summarising risk estimates from high-quality studies strengthened the inverse association between physical activity and renal cancer risk (RR=0.78; 95% CI=0.66–0.92). Effect modification by adiposity, hypertension, type 2 diabetes, smoking, gender, or geographic region was not observed.

**Conclusion:** Our comprehensive meta-analysis provides strong support for an inverse relation of physical activity to renal cancer risk. Future high-quality studies are required to discern which specific types, intensities, frequencies, and durations of physical activity are needed for renal cancer risk reduction.

Renal cancer is one of the top 10 cancer sites in the United States and Europe. Each year, about 65 000 new cases are diagnosed in the United States (Howlader *et al*, 2012) and about 60 000 new cases are diagnosed in the European Union (Boyle and Ferlay, 2005). Well-established unfavourable risk factors for renal cancer include smoking, obesity, hypertension, and type 2 diabetes mellitus (Scelo and Brennan, 2007). In contrast, physical activity may prevent the development of renal cancer, partly because it helps reduce obesity (Wing, 1999), blood pressure (Blair *et al*, 1984), and insulin resistance (Rosenthal *et al*, 1983). Physical activity may also

independently decrease renal cancer risk by lowering lipid peroxidation levels (Vincent *et al*, 2002). However, few available epidemiologic studies have been able to report a clear inverse association between physical activity and renal cancer (Leitzmann, 2011). Moreover, no meta-analysis is available on the relation between physical activity and renal cancer. To address this research gap, we conducted a systematic literature search and meta-analysis to quantify the association between physical activity and renal cancer, taking into account the methodologic quality of the studies.

\*Correspondence: Dr G Behrens; E-mail: gundula.behrens@klinik.uni-regensburg.de

Received 18 October 2012; revised 3 January 2013; accepted 7 January 2013; published online 14 February 2013

© 2013 Cancer Research UK. All rights reserved 0007–0920/13

## MATERIALS AND METHODS

**Literature search.** Our systematic review and meta-analysis adhered to the PRISMA guidelines (Moher *et al*, 2009). Both authors searched the literature using PubMed (see Appendix B for PubMed search options) and Web of Knowledge (see Appendix C for Web of Knowledge search options) to identify published non-ecologic epidemiologic studies quantifying the relationship between physical activity and renal cancer risk in individuals without a cancer history. That search was complemented by a scan of the reference lists of the identified studies and a scan of the reference list of a previous systematic review (Leitzmann, 2011). We considered all human research articles published in English through the end of September 2012 not classified as review, meta-analysis, editorial, comment, letter, practice guideline, or news. Our search strategy included the terms physical activity, exercise, cardiorespiratory fitness, cardiovascular fitness, resistance training, endurance training, aerobic, sport, athletes, players, lifestyle, kidney cancer, renal cancer, renal cell cancer, renal carcinoma, renal cell carcinoma, cancer, risk, incidence, and mortality. The search strategy excluded research on cancer survivors and research on specific types of cancer other than renal cancer. That search yielded 586 potential articles. Irrelevant articles were eliminated after screening titles and abstracts ( $n=477$ ) or manuscripts ( $n=82$ ). The 27 remaining studies (Goodman *et al*, 1986; Paffenbarger *et al*, 1987; Brownson *et al*, 1991; Lindblad *et al*, 1994; Mellemegaard *et al*, 1994, 1995; Bergstrom *et al*, 1999, 2001; Parker *et al*, 2002; Menezes *et al*, 2003; Mahabir *et al*, 2004; Nicodemus *et al*, 2004; van Dijk *et al*, 2004; Washio *et al*, 2005; Chiu *et al*, 2006; Pan *et al*, 2006; Setiawan *et al*, 2007; Tavani *et al*, 2007; Hu *et al*, 2008, 2009; Moore *et al*, 2008; Thompson *et al*, 2008; Yun *et al*, 2008; Spyridopoulos *et al*, 2009; Wilson *et al*, 2009; George *et al*, 2011; Parent *et al*, 2011) proved to be relevant.

To avoid duplicate information from overlapping studies, we removed eight of the 27 identified studies because their results were pooled (Lindblad *et al*, 1994; Mellemegaard *et al*, 1994) or updated (Parker *et al*, 2002; Menezes *et al*, 2003; Pan *et al*, 2006) in studies (Mellemegaard *et al*, 1995; Chiu *et al*, 2006; Hu *et al*, 2008) using the same database, because they reported results (Hu *et al*, 2009; Wilson *et al*, 2009) presented earlier (Mahabir *et al*, 2004; Hu *et al*, 2008), or because their investigations of total sitting time (George *et al*, 2011) were closely related to a previous study (Moore *et al*, 2008) on physical activity from the same cohort. The remaining 19 studies (Goodman *et al*, 1986; Paffenbarger *et al*, 1987; Brownson *et al*, 1991; Mellemegaard *et al*, 1995; Bergstrom *et al*, 1999, 2001; Mahabir *et al*, 2004; Nicodemus *et al*, 2004; van Dijk *et al*, 2004; Washio *et al*, 2005; Chiu *et al*, 2006; Setiawan *et al*, 2007; Tavani *et al*, 2007; Hu *et al*, 2008; Moore *et al*, 2008; Thompson *et al*, 2008; Yun *et al*, 2008; Spyridopoulos *et al*, 2009; Parent *et al*, 2011) were included in the meta-analysis.

**Quality score.** The magnitude and heterogeneity of risk estimates may depend on the methodologic quality associated with the underlying study and with the risk estimate derivation. Similar to three previous systematic reviews (Monninkhof *et al*, 2007; Voskuil *et al*, 2007; Liu *et al*, 2011) on the association between physical activity and specific types of cancer, both authors employed a quality score proposed by Voskuil *et al* (2007) to assess the methodologic quality of the studies and the consistency of the available evidence. Please refer to Appendix A for a description of the items covered by the quality score.

**Main statistical analysis.** Because some studies presented risk estimates for men and women and some studies investigated more than one physical activity domain, the 19 identified studies reported a total of 37 risk estimates. If separate risk estimates were

available for men and women, both risk estimates were included in the meta-analysis because they were based on independent samples. To prevent potential bias arising from the fact that the risk estimates for the various physical activity domains were based on the same study population, both authors allowed only one estimate per study and gender in the main analysis. Specifically, if more than one physical activity domain was studied, we selected the risk estimate with the highest quality score in the main analysis. Of the 37 risk estimates, 25 were included in the main analysis.

In the meta-analysis, we interpreted odds ratios and hazard ratios as relative risk estimates (RR<sub>*i*</sub>), computed the natural logarithms of those risk estimates  $\log(\text{RR}_i)$  with corresponding standard errors  $s_i = (\log(\text{upper } 95\% \text{ confidence interval (CI) bound of RR}) - \log(\text{RR}_i))/1.96$ , and employed a random-effects model to determine the weighted average of those  $\log(\text{RR}_i)$ s while allowing for heterogeneity of effects. In the random-effects model, the  $\log(\text{RR}_i)$ s were weighted by  $w_i = 1/(s_i^2 + t^2)$  where  $s_i$  represented the standard error of  $\log(\text{RR}_i)$  and  $t^2$  represented the restricted maximum-likelihood estimate of the overall variance (Higgins and Thompson, 2002). In one case (Paffenbarger *et al*, 1987), we derived the standard error of the  $\log(\text{RR}_i)$  using the *P*-value accompanying the RR estimate. In five additional cases (Brownson *et al*, 1991; Mellemegaard *et al*, 1995; Bergstrom *et al*, 1999, 2001; Chiu *et al*, 2006), the reported RRs used the highest rather than the lowest activity level as the reference category, so we reversed those RRs for comparability. Heterogeneity of the risk estimates was assessed using the *Q*- and *I*<sup>2</sup>-statistics (Higgins and Thompson, 2002). Publication bias was tested using funnel plots (Egger *et al*, 1997), Egger's regression test (Egger *et al*, 1997), and Begg's rank correlation test (Begg and Mazumdar, 1994).

**Statistical subanalyses.** If a study presented separate risk estimates for recreational, occupational, and total physical activity, in a subanalysis all those risk estimates were included in the meta-analysis. Also, in a subanalysis we used all 37 risk estimates to investigate the impact of prespecified potentially influential methodologic factors on the summary risk estimate.

On the basis of pre-existing evidence, we hypothesised that the relations of physical activity to renal cell cancer would differ according to study design (cohort or case-control), physical activity domain (recreational, occupational, or total physical activity), and gender (men, women, or men and women combined). Thus, we conducted subanalyses within categories of those variables. We also performed exploratory analyses that were stratified by geographic region (North America, Europe, Asia), type of physical activity assessment (energy expenditure, physical fitness, moderate-to-vigorous physical activity duration, moderate-to-vigorous physical activity frequency, and qualitative assessments using categories, such as 'sedentary', 'light', 'moderate', or 'high' physical activity), timing in life of physical activity (recent physical activity, past physical activity, or consistent physical activity over time), number of adjustment factors (in quartiles), adjustments for smoking and obesity (adjusted for smoking and obesity, adjusted for smoking but not obesity, adjusted neither for smoking nor obesity; the option of adjusting for obesity but not smoking was not included because it did not occur), adjustment for hypertension (yes, no), adjustment for type 2 diabetes mellitus (yes, no), or methodologic quality score (in tertiles). To assess the influence of those factors, we applied random-effects meta-analysis regression comparing the model including the current factor of interest as a single explanatory variable with the null model not including any explanatory variables.

All statistical analyses were performed in R (R Development Core Team, 2011) using the R-package 'metafor' (Viechtbauer, 2010). Risk estimates are reported with 95% CIs. Statistical significance is based on the 5% significance level.

Table 1. Characteristics of the 19 studies on physical activity and renal cancer risk included in the meta-analysis

Authors, year, gender	Subjects	Cases	Region	Adjustment factors (excluding age, sex)	PA domain	Timing in life of PA	Relative risk (95% CI), high vs low PA	Low PA defined by	High PA defined by	Quality score (%)
<b>Case-control studies</b>										
Brownson et al (1991) Men	17 147	449	North America	Smoking	Occupational	Recent PA	<b>0.77 (0.50, 1.11)</b>	Low PA	High PA	45
Chiu et al (2006) Men	1660	225	North America	History of hypertension (yes/no), family history of renal cancer, marital status, red meat intake, smoking, total energy intake, vegetable intake	Recreational	Consistent PA over time	<b>0.83 (0.48, 1.43)</b>	MVPA (10 min) less than once per month	MVPA (10 min) every day	76
Women	829	123			Recreational	Consistent PA over time	<b>0.40 (0.19, 0.83)</b>	MVPA (10 min) less than once per month	MVPA (10 min) every day	76
Goodman et al (1986) Men	378	189	North America	—	Recreational	Consistent PA over time	<b>1.14 (0.65, 2.05)</b>	None/occasional PA	Strenuous PA	66
Women	156	78			Occupational	Consistent PA over time	0.88 (0.48, 1.55)	None/occasional PA	Strenuous PA	62
					Recreational	Consistent PA over time	<b>1.11 (0.44, 2.97)</b>	None/occasional PA	Strenuous PA	66
					Occupational	Consistent PA over time	1.20 (0.41, 4.06)	None/occasional PA	Strenuous PA	62
Hu et al (2008) Men and women	6177	1138	North America	Residential area	Recreational	Recent PA	<b>0.90 (0.71, 1.14)</b>	No PA	55 min or more of MVPA per week	60
Meillemgaard et al (1995) Men	1994	864	Europe	Obesity (BMI), smoking, study centre	Recreational	Past PA	<b>1.11 (0.56, 2.50)</b>	Not physically active	Very active	62
Women	1308	572			Occupational	Past PA	1.11 (0.71, 1.67)	Not physically active	Very active	58
					Recreational	Past PA	<b>1.67 (0.71, 3.33)</b>	Not physically active	Very active	62
					Occupational	Past PA	1.67 (0.91, 3.33)	Not physically active	Very active	58
Parent et al (2011) Men	533	177	North America	Alcohol intake, coffee intake, obesity (BMI), proxy, race/ethnicity, recreational/occupational activity (mutual adjustment), SES (socio-economic status, education), smoking	Total	Consistent PA over time	1.02 (0.70, 1.49)	Less than 1.5 MET at work independent of leisure time PA or 1.5–3.9 MET at work and less than once per week engaged in leisure time MVPA	Energy expenditure of 4 MET per day or more at work independent of leisure time PA or 1.6–3.9 MET per day at work and at least once per week engaged in leisure time MVPA	68
					Recreational	Past PA	<b>1.11 (0.76, 1.64)</b>	MVPA less than once per week	MVPA at least once per week	69
					Occupational	Consistent PA over time	0.84 (0.38, 1.89)	1.5 MET or less of energy expenditure	Energy expenditure of 4 MET or more per day	64

Table 1. (Continued)

Authors, year, gender	Subjects	Cases	Region	Adjustment factors (excluding age, sex)	PA domain	Timing in life of PA	Relative risk (95% CI), high vs low PA	Low PA defined by	High PA defined by	Quality score (%)
Spyridopoulos et al (2009) Men and women	350	70	Europe	Alcohol intake, coffee intake, history of diabetes (yes/no), obesity (serum adiponectin, serum leptin, waist-hip ratio), protein intake, SES (education), smoking, vegetarian diet	Recreational	Recent PA	<b>0.62 (0.48, 0.82)</b>	—	Increment of 3.5 h of MVPA per week	78
Tavani et al, 2007 Men and women	2301	767	Europe	Calendar year of interview, history of hypertension (yes/no), obesity (BMI), smoking, study centre	Recreational Occupational	Past PA Past PA	<b>1.03 (0.78, 1.36)</b> 0.71 (0.55, 0.92)	Less than 2 h of MVPA per week Low PA	More than 7 h of MVPA per week High PA	66 55
<b>Cohort studies</b>										
Bergstrom et al, 1999 Men	674 025	2704	Europe	Calendar year of follow-up, residential area, SES (job title)	Occupational	Past PA	<b>0.80 (0.65, 0.98)</b>	Sedentary activities	High PA	58
Women	253 336	587			Occupational	Past PA	<b>1.25 (0.79, 1.96)</b>	Sedentary activities	High PA	58
Bergstrom et al, 2001 Men and women	17 241	102	Europe	History of hypertension (yes/no), obesity (BMI), smoking	Recreational Occupational	Consistent PA over time Consistent PA over time	<b>1.67 (0.83, 3.33)</b> 1.25 (0.63, 2.50)	Sedentary activities Sedentary activities	Strenuous PA Strenuous PA	83 71
Mahabir et al (2004) Men	29 133	210	Europe	Alcohol intake, dietary fat intake, fruit and vegetable intake, history of hypertension (blood pressure), intervention group, recreational/occupational activity (mutual adjustment), obesity (BMI), residential area, serum cholesterol, SES (education), smoking, total energy intake	Recreational Occupational	Recent PA	<b>0.46 (0.18, 1.13)</b>	Light PA	Heavy PA	75
					Occupational	Recent PA	1.08 (0.54, 2.15)	Sedentary activities	Heavy PA	63

Table 1. (Continued)

Authors, year, gender	Subjects	Cases	Region	Adjustment factors (excluding age, sex)	PA domain	Timing in life of PA	Relative risk (95% CI), high vs low PA	Low PA defined by	High PA defined by	Quality score (%)
Moore et al (2008) Men and women	482 386	1238	North America	Body height, history of diabetes (yes/no), history of hypertension (yes/no), obesity (BMI), protein intake, race/ethnicity, smoking.	Recreational  Occupational	Recent PA  Recent PA	<b>0.77 (0.64, 0.92)</b>  0.84 (0.57, 1.22)	Never/rarely engaging in VPA  Mostly sitting	Five times per week or more engaged in VPA (more than 20 min)  Heavy PA	76  65
Nicodemus et al (2004) Women	34 637	124	North America	—	Recreational	Recent PA	<b>0.37 (0.14, 0.99)</b>	Low VPA frequency	High VPA frequency	71
Paffenbarger et al (1987) Men and women	56 683	53	North America	Birth year	Recreational	Past PA	<b>0.95 (0.47, 1.94)</b>	Less than 5 h of VPA per week	5 h of VPA per week or more	61
Setiawan et al (2007) Men  Women	75 162  85 964	220  127	North America	Alcohol intake, history of hypertension (yes/no), obesity (BMI), smoking	Total  Total	Recent PA  Recent PA	<b>1.09 (0.75, 1.58)</b>  <b>0.66 (0.40, 1.10)</b>	1.4 MET per day or less  1.4 MET per day or less	Energy expenditure of 1.8 MET per day or more  Energy expenditure of 1.8 MET per day or more	77  77
Thompson et al (2008) Men	21 663	31	North America	Alcohol intake, examination year, history of cancer, history of diabetes (fasting glucose level), obesity (BMI), smoking	Total	Recent PA	<b>0.91 (0.45, 2.68)</b>	Lowest physical fitness quintile	Upper two physical fitness quintiles	69
van Dijk et al (2004) Men  Women	2335  2444	179  96	Europe	Obesity (BMI), smoking, total energy intake	Recreational  Occupational  Recreational	Recent PA  Consistent PA over time  Recent PA	<b>0.74 (0.44, 1.23)</b>  0.82 (0.46, 1.47)  <b>1.13 (0.56, 2.29)</b>	Less than 30 min of MVPA per day  Energy expenditure of <8 kJ min <sup>-1</sup>  Less than 30 min of MVPA per day	More than 10.5 h of MVPA per week  Energy expenditure of >12 kJ min <sup>-1</sup>  More than 10.5 h of MVPA per week	77  76  77
Washio et al (2005) Men and women	114 517	38	Asia	—	Recreational  Occupational	Recent PA  Recent PA	<b>0.54 (0.25, 1.18)</b>  1.44 (0.72, 2.88)	MVPA less than once per week  Sedentary activities	MVPA once per week or more  Physically active	58  51
Yun et al (2008) Men	444 963	395	Asia	Alcohol intake, dietary pattern, history of diabetes (fasting glucose level), obesity (BMI), SES (employment), smoking	Recreational	Recent PA	<b>1.01 (0.83, 1.23)</b>	Combination of MVPA frequency and duration: MVPA less than 4 times per week and less than 30 min per session	Combination of MVPA frequency and duration: MVPA at least five times per week and at least 30 min per session	71

Abbreviations: BMI = body mass index; MET = metabolic equivalent of task; MVPA = moderate-to-vigorous physical activity; PA = physical activity; RR = relative risk; SES = socioeconomic status; VPA = vigorous physical activity. The 19 studies are grouped by study design. The main meta-analysis considered just one risk estimate (in bold) per study and gender.

Table 2. Summary risk estimates and  $I^2$  measures of heterogeneity from random-effects models stratified by selected study characteristics


Stratification criterion	Number of included RRs	RR (95% CI) (high vs low PA) from random-effects model	$I^2$ (%)	P-value <sup>a</sup>
<b>Methodologic quality<sup>b</sup></b>				
RRs within upper tertile of quality score	11	0.78 (0.66, 0.92)	33	0.02
RRs within intermediate tertile of quality score	12	1.00 (0.89, 1.13)	0	
RRs within lower tertile of quality score	14	0.93 (0.80, 1.07)	30	
<b>PA assessment</b>				
RRs based on qualitative PA assessments	18	0.98 (0.85, 1.14)	35	0.24
RRs based on energy expenditure	6	0.97 (0.84, 1.12)	0	
RRs based on MVPA duration	6	0.85 (0.69, 1.04)	43	
RRs based on MVPA frequency	6	0.72 (0.53, 0.97)	53	
<b>PA domain</b>				
RRs based on total activity	4	0.95 (0.76, 1.20)	0	0.84
RRs based on occupational activity	14	0.91 (0.79, 1.04)	21	
RRs based on recreational activity	19	0.88 (0.77, 1.00)	40	
<b>Timing in life of PA</b>				
RRs based on recent PA	16	0.83 (0.74, 0.93)	28	0.18
RRs based on consistent PA over time	11	0.96 (0.79, 1.15)	0	
RRs based on past PA	10	1.01 (0.84, 1.20)	46	
<b>Gender</b>				
RRs among men	17	0.93 (0.84, 1.02)	2	0.41
RRs among women	9	0.95 (0.66, 1.36)	57	
RRs among men and women	11	0.85 (0.73, 0.98)	42	
<b>Study design</b>				
RRs from case-control studies	18	0.91 (0.79, 1.04)	36	0.93
RRs from cohort studies	19	0.89 (0.80, 0.99)	19	
<b>Study region</b>				
RRs from studies in North America	18	0.85 (0.77, 0.94)	0	0.63
RRs from studies in Europe	16	0.95 (0.80, 1.12)	51	
RRs from studies in Asia	3	1.00 (0.83, 1.20)	0	
<b>Number of adjustment factors<sup>c</sup></b>				
RRs within upper tertile of number of adjustment factors	12	0.83 (0.71, 0.97)	40	0.28
RRs within intermediate tertile of number of adjustment factors	4	0.87 (0.68, 1.10)	52	
RRs within lower tertile of number of adjustment factors	21	0.96 (0.85, 1.08)	14	
<b>Adjustment for smoking and obesity</b>				
RRs adjusted for smoking and obesity	23	0.92 (0.82, 1.03)	37	0.31
RRs adjusted for smoking but not obesity	3	0.71 (0.54, 0.94)	0	
RRs adjusted neither for smoking nor obesity	11	0.89 (0.78, 1.01)	2	
<b>Adjustment for hypertension</b>				
RRs adjusted for hypertension	12	0.85 (0.73, 0.97)	30	0.30
RRs not adjusted for hypertension	25	0.93 (0.83, 1.03)	24	
<b>Adjustment for diabetes</b>				
RRs adjusted for diabetes	5	0.81 (0.66, 0.99)	57	0.18
RRs not adjusted for diabetes	32	0.92 (0.84, 1.01)	14	

Abbreviations: CI = confidence interval; MVPA = moderate-to-vigorous physical activity; PA = physical activity; RR = relative risk.

<sup>a</sup>P-values for effect heterogeneity across strata were obtained from random-effects meta-regression comparing the model including the stratification variable as a single explanatory variable with the null model not including any explanatory variables.

<sup>b</sup>The quality scores ranged from 45 to 83 percentage points (out of 100 percentage points), with lower and upper tertile cutoffs of 62 percentage points and 71 percentage points, respectively.

<sup>c</sup>The number of adjustment factors (not counting adjustments for age and sex) ranged between 0 and 12, with lower and upper tertile cutoffs of 3 and 5, respectively.



**Figure 1.** Forest plot corresponding to the main random-effects meta-analysis including 25 risk estimates quantifying the relationship between high physical activity and renal cancer risk. Relative risks (RRs) compare high vs low levels of physical activity and are grouped by study design. The size of the box representing each risk estimate is proportional to the weight that the risk estimate contributed to the summary risk estimate.

## RESULTS


**Description of underlying study characteristics.** Table 1 presents the 19 studies on physical activity and renal cancer risk included in the meta-analysis. Because six studies stratified results by gender and nine studies investigated more than one physical activity domain, the 19 studies reported a total of 37 risk estimates.

When grouping studies by potentially effect modifying factors (Table 2), we noted that there was an equal number of risk estimates from case-control and prospective cohort studies, with the vast majority of studies originating in the United States or Europe. Half of the risk estimates were based on recreational physical activity, one-third of the risk estimates were based on occupational activity, and four risk estimates were based on total physical activity. Half of the physical activity assessments were of a qualitative type and the remaining half were of a quantitative type. Nearly two-thirds of the risk estimates were adjusted for smoking and obesity, one-third of the risk estimates were adjusted for hypertension, and one-sixth of the risk estimates were adjusted for history of type 2 diabetes mellitus.

**Meta-analysis.** The random-effects model summarising the 25 risk estimates with the highest quality scores from each of the 19 studies (Figure 1) revealed a statistically significant 12% reduction in renal cancer risk when comparing a high with a low level of physical activity (RR = 0.88; 95% CI = 0.79–0.97;  $I^2 = 33\%$ ). The magnitude of that summary risk estimate did not materially change when grouping those 25 risk estimates by study design (Figure 1), physical activity domain (Figure 2), or gender (Figure 3). That meta-analysis combined a total of 2 327 322 subjects and 10 756 cases. No publication bias was indicated by the funnel plot (Figure 4), Egger's regression test ( $P = 0.89$ ), or Begg's rank correlation test ( $P = 0.53$ ).

**Renal cancer end point.** Because physical activity may differentially impact the incidence vs mortality of kidney cancer, we repeated the main analysis after excluding the two risk estimates from the papers on kidney cancer mortality (Washio *et al*, 2005; Thompson *et al*, 2008). The summary risk estimate remained unchanged (RR = 0.88; 95% CI = 0.80–0.98).

**Potentially influential methodologic factors.** In subanalyses investigating potentially influential methodologic factors, all 37



**Figure 2.** Forest plot corresponding to the main random-effects meta-analysis including 25 risk estimates quantifying the relationship between high physical activity and renal cancer risk. Relative risks (RRs) compare high vs low levels of physical activity and are grouped by physical activity domain. The size of the box representing each risk estimate is proportional to the weight that the risk estimate contributed to the summary risk estimate.

risk estimates were used. The random-effects summary risk estimate (RR = 0.90; 95% CI = 0.82–0.98;  $I^2 = 26\%$ ) of those 37 risk estimates did not substantially differ from that of the main analysis. We found that the methodologic quality score significantly influenced the magnitude of the summary risk estimate ( $P = 0.02$ ; Table 2) but not the underlying overall variation  $I^2$ . The best evidence synthesis of studies that fell into the high tertile of the quality score yielded a meta-analysis estimate for the relation of physical activity to renal cancer of 0.78 (95% CI = 0.66–0.92;  $I^2 = 0.02$ ). In contrast, the meta-analysis RRs for studies falling into the intermediate (RR = 1.00; 95% CI = 0.89–1.13;  $I^2 = 0$ ) and lower (RR = 0.93; 95% CI = 0.80–1.07;  $I^2 = 0.02$ ) tertiles of the quality score were statistically nonsignificant.

When stratifying by the type of physical activity assessment, the summary risk estimates based on frequency of moderate-to-vigorous physical activity (RR = 0.72; 95% CI = 0.53–0.97) or duration of moderate-to-vigorous physical activity (RR = 0.85; 95% CI = 0.69–1.04) appeared to be stronger than those based on energy expenditure (RR = 0.97; 95% CI = 0.84–1.12) or qualitative physical activity assessments (RR = 0.98; 95% CI = 0.85–1.14). However, that variation was not statistically significant ( $P = 0.24$ ). Similarly, the magnitude of the inverse association between physical activity and renal cancer appeared to be stronger with a larger number of adjustment factors, although that difference was not statistically significant ( $P = 0.28$ ). The meta-analysis RR for studies in the top tertile of the number of adjustment factors was






Figure 3. Forest plot corresponding to the main random-effects meta-analysis including 25 risk estimates quantifying the relationship between high physical activity and renal cancer risk. Relative risks (RRs) compare high vs low levels of physical activity and are grouped by gender. The size of the box representing each risk estimate is proportional to the weight that the risk estimate contributed to the summary risk estimate.




Figure 4. Funnel plot corresponding to the main random-effects meta-analysis including 25 risk estimates quantifying the relationship between high physical activity and renal cancer risk.

0.83 (95% CI = 0.71–0.97), whereas the RR for studies in the bottom tertile of the number of adjustment factors was 0.96 (95% CI = 0.85–1.08). There was no difference in risk estimates between study designs (RR for case-control studies = 0.91; 95% CI = 0.79–1.04; RR for cohort studies = 0.89; 95% CI = 0.80–0.99; *P*-value for interaction = 0.93). Similarly, none of the following remaining study characteristics affected the summary risk estimates: physical activity domain (*P* = 0.84), timing in life of physical activity (*P* = 0.18), gender (*P* = 0.41), geographic region (*P* = 0.63), joint adjustment for smoking and obesity (*P* = 0.31), hypertension adjustment (*P* = 0.30), and diabetes adjustment (*P* = 0.18).

We further examined study characteristics according to the quality score (Table 3). Studies that fell into the top tertile of the quality score tended to employ quantitative physical activity assessments, to investigate recreational activity, to examine recent physical activity, to use a cohort design, and to adjust for smoking, obesity, hypertension, and diabetes. In contrast, studies in the bottom tertile of the quality score tended to employ qualitative physical activity assessments, to investigate

Table 3. Distribution of methodologic characteristics (absolute frequencies) of all 37 risk estimates by tertile of quality score<sup>a</sup>

Methodologic characteristics	RRs within upper tertile of quality score	RRs within intermediate tertile of quality score	RRs within lower tertile of quality score
<b>PA assessment</b>			
RRs based on qualitative PA assessments	2	5	11
RRs based on energy expenditure	3	3	0
RR based on physical fitness	0	1	0
RRs based on MVPA duration	3	1	2
RRs based on MVPA frequency	3	2	1
<b>PA domain</b>			
RRs based on total activity	2	2	0
RRs based on occupational activity	1	4	9
RRs based on recreational activity	8	6	5
<b>Timing in life of PA</b>			
RRs based on recent PA	7	5	4
RRs based on consistent PA over time	4	5	2
RRs based on past PA	0	2	8
<b>Gender</b>			
RRs among men	5	7	5
RRs among women	3	2	4
RRs among men and women	3	3	5
<b>Study design</b>			
RRs from case-control studies	3	6	9
RRs from cohort studies	8	6	5
<b>Study region</b>			
RRs from studies in North America	5	8	5
RRs from studies in Europe	6	3	7
RRs from studies in Asia	0	1	2
<b>Number of adjustment factors<sup>b</sup></b>			
RRs within upper tertile of number of adjustment factors	5	7	0
RRs within intermediate tertile of number of adjustment factors	2	1	1
RRs within lower tertile of number of adjustment factors	4	4	13
<b>Adjustment for smoking and obesity</b>			
RRs adjusted for smoking and obesity	9	9	5
RRs adjusted for smoking but not obesity	2	0	1
RRs adjusted neither for smoking nor obesity	0	3	8
<b>Adjustment for hypertension</b>			
RRs adjusted for hypertension	7	4	1
RRs not adjusted for hypertension	4	8	13
<b>Adjustment for diabetes</b>			
RRs adjusted for diabetes	2	3	0
RRs not adjusted for diabetes	9	9	14

Abbreviations: MVPA = moderate-to-vigorous physical activity; PA = physical activity; RR = relative risk.

<sup>a</sup>The quality scores ranged from 45 to 83 percentage points (out of 100 percentage points), with lower and upper tertile cutoffs of 62 percentage points and 71 percentage points, respectively.

<sup>b</sup>The number of adjustment factors (not counting adjustments for age and sex) ranged between 0 and 12, with lower and upper tertile cutoffs of 3 and 5, respectively.

occupational activity, to examine past physical activity, to use a case-control design, and to not adjust for smoking, obesity, hypertension, or diabetes.

After adjusting the main random-effects model for study quality (in tertiles), the previously observed heterogeneity ( $P$ -heterogeneity = 0.03) of risk estimates was no longer evident ( $P$ -heterogeneity = 0.12).

## DISCUSSION

**Main results.** This comprehensive meta-analysis revealed a statistically significant 12% reduction in renal cancer risk associated with a high vs low level of physical activity.

**Potentially influential factors.** The summary RR estimate was not affected by individual potentially influential factors, such as type of physical activity assessment, physical activity domain, timing in life of physical activity, gender, study design, study region, number of adjustment factors, and adjustments for smoking, obesity, hypertension, or diabetes. However, the quality score representing a combination of specific factors affected the summary risk estimate. Summary risk estimates based on studies that fell into the top quality score tertile were statistically significantly inverse, whereas summary risk estimates based on studies that fell into the intermediate or bottom quality score tertiles were not. After adjusting for study quality, the previously observed heterogeneity in the random-effects model was no longer statistically significant.

The influence of individual factors was examined in previous meta-analyses of physical activity and cancers of the colorectum (Harriss *et al*, 2009; Boyle *et al*, 2012), pancreas (Bao and Michaud, 2008), and prostate (Liu *et al*, 2011). In agreement with our observations, no statistically significant heterogeneity across gender (Bao and Michaud, 2008; Boyle *et al*, 2012), study design (Liu *et al*, 2011), geographic region (Bao and Michaud, 2008; Harriss *et al*, 2009), physical activity domain (Boyle *et al*, 2012), or obesity adjustment (Bao and Michaud, 2008; Harriss *et al*, 2009) was reported.

The influence of a quality score combining several factors was previously studied with respect to the associations between physical activity and cancers of the breast (Monninkhof *et al*, 2007), endometrium (Voskuil *et al*, 2007), prostate (Liu *et al*, 2011), colon (Boyle *et al*, 2012), and pancreas (O'Rorke *et al*, 2010). In agreement with our findings, the meta-analysis on physical activity and breast cancer (Monninkhof *et al*, 2007) detected a more pronounced risk reduction with increased quality score, while the remaining analyses (Voskuil *et al*, 2007; O'Rorke *et al*, 2010; Liu *et al*, 2011; Boyle *et al*, 2012) did not detect any statistically significant association between quality score and summary risk estimates. Two reviews (O'Rorke *et al*, 2010; Boyle *et al*, 2012), however, described decreased variation in risk estimates with increasing quality score. No such observation was made in this study.

**Potential biological mechanisms.** A high level of physical activity has been shown to reduce adiposity (Wing, 1999), hypertension (Blair *et al*, 1984), insulin resistance (Rosenthal *et al*, 1983), circulating levels of insulin-like growth factor 1 (Eliakim *et al*, 1996, 1998), and lipid peroxidation (Vincent *et al*, 2002) – factors positively associated with the development of renal carcinoma (Kellerer *et al*, 1995; Chow *et al*, 2000; Gago-Dominguez *et al*, 2002; van Dijk *et al*, 2004; Vatten *et al*, 2007; Yuen *et al*, 2009). Further potential cancer preventing mechanisms include the beneficial effects of physical activity on chronic inflammation and immune function (McTiernan, 2008). It is hypothesised, however, that the effects of physical activity on chronic inflammation are mediated, in part, through avoidance of adiposity. The exact mechanisms linking physical activity to immune function related to tumour suppression have not yet been established, but it is thought that physical activity improves the number or the function of natural killer cells.

**Strengths and limitations.** This is the first meta-analysis of physical activity and renal cancer. It bears the strengths and limitations inherent in any meta-analysis combining results from studies with heterogeneous study designs (Greenland and O'Rourke, 2008). Particular strengths of the current meta-analysis are that it is based on an extensive systematic literature review, that

it rigorously excluded duplicate information induced by overlapping studies, and that it combined information from 19 studies, including a total of 2 327 322 subjects and 10 756 cases. A further strength is that it is among the few meta-analyses of physical activity and a specific type of cancer to assess the heterogeneity of summary estimates by potentially influential factors underlying the RR estimates. The employed quality score addressed potential selection, misclassification, and confounding biases, and accounted for heterogeneity of the results. An inverse association between physical activity and renal cancer risk was observed in analyses including all risk estimates and in analyses including only risk estimates from high-quality studies. In addition, no publication bias was detected.

One limitation of this meta-analysis is the large variation in the underlying studies regarding their definitions of exposure to physical activity – ranging from 'physically very active' to '5 h of vigorous physical activity per week or more'. Similarly, the definitions of physical activity referent groups ranged from 'not physically active' to '<5 h of vigorous physical activity per week'. Such variation did not allow us to conduct stratified analyses according to comparable groups of exposed and non-exposed individuals. Thus, we were not able to identify the specific type, intensity, frequency, and duration of physical activity required to lower renal cancer risk.

**Conclusion.** In conclusion, our comprehensive meta-analysis provides strong support for an inverse relation of physical activity to the risk of renal cancer. On the basis of high-quality studies, physical activity may decrease the risk of renal cancer by 22%. Future research is required to discern which specific types, intensities, frequencies, and durations of physical activity are needed for renal cancer risk reduction. High-quality studies that employ standardised physical activity assessments and uniform definitions of physical activity are warranted.

## REFERENCES

- Bao Y, Michaud DS (2008) Physical activity and pancreatic cancer risk: a systematic review. *Cancer Epidemiol Biomarkers Prev* **17**: 2671–2682.
- Begg CB, Mazumdar M (1994) Operating characteristics of a rank correlation test for publication bias. *Biometrics* **50**: 1088–1101.
- Bergstrom A, Moradi T, Lindblad P, Nyren O, Adami HO, Wolk A (1999) Occupational physical activity and renal cell cancer: a nationwide cohort study in Sweden. *Int J Cancer* **83**: 186–191.
- Bergstrom A, Terry P, Lindblad P, Lichtenstein P, Ahlbom A, Feychting M, Wolk A (2001) Physical activity and risk of renal cell cancer. *Int J Cancer* **92**: 155–157.
- Blair SN, Goodyear NN, Gibbons LW, Cooper KH (1984) Physical fitness and incidence of hypertension in healthy normotensive men and women. *JAMA* **252**: 487–490.
- Boyle P, Ferlay J (2005) Cancer incidence and mortality in Europe, 2004. *Ann Oncol* **16**: 481–488.
- Boyle T, Keegel T, Bull F, Heyworth J, Fritschi L (2012) Physical activity and risks of proximal and distal colon cancers: a systematic review and meta-analysis. *J Natl Cancer Inst* **104**(20): 1548–1561.
- Brownson RC, Chang JC, Davis JR, Smith CA (1991) Physical activity on the job and cancer in Missouri. *Am J Public Health* **81**: 639–642.
- Chiu BC, Gapstur SM, Chow WH, Kirby KA, Lynch CF, Cantor KP (2006) Body mass index, physical activity, and risk of renal cell carcinoma. *Int J Obes (Lond)* **30**: 940–947.
- Chow WH, Gridley G, Fraumeni Jr JF, Jarvholm B (2000) Obesity, hypertension, and the risk of kidney cancer in men. *N Engl J Med* **343**: 1305–1311.
- Egger M, Davey Smith G, Schneider M, Minder C (1997) Bias in meta-analysis detected by a simple, graphical test. *BMJ* **315**: 629–634.
- Eliakim A, Brasel JA, Mohan S, Barstow TJ, Berman N, Cooper DM (1996) Physical fitness, endurance training, and the growth hormone-insulin-like growth factor I system in adolescent females. *J Clin Endocrinol Metab* **81**: 3986–3992.

- Eliakim A, Brasel JA, Mohan S, Wong WL, Cooper DM (1998) Increased physical activity and the growth hormone-IGF-I axis in adolescent males. *Am J Physiol* **275**: R308–R314.
- Gago-Dominguez M, Castela JE, Yuan JM, Ross RK, Yu MC (2002) Lipid peroxidation: a novel and unifying concept of the etiology of renal cell carcinoma (United States). *Cancer Causes Control* **13**: 287–293.
- George SM, Moore SC, Chow WH, Schatzkin A, Hollenbeck AR, Matthews CE (2011) A prospective analysis of prolonged sitting time and risk of renal cell carcinoma among 300,000 older adults. *Ann Epidemiol* **21**: 787–790.
- Goodman MT, Morgenstern H, Wynder EL (1986) A case-control study of factors affecting the development of renal cell cancer. *Am J Epidemiol* **124**: 926–941.
- Greenland S, O'Rourke K (2008) Meta-analysis. In *Modern Epidemiology*, Rothman KJ, Greenland S, Lash TL (eds) 3rd edn pp 652–682. Lippincott Williams & Wilkins: Philadelphia, PA.
- Harriss DJ, Atkinson G, Batterham A, George K, Cable NT, Reilly T, Haboubi N, Renehan AG (2009) Lifestyle factors and colorectal cancer risk (2): a systematic review and meta-analysis of associations with leisure-time physical activity. *Colorectal Dis* **11**: 689–701.
- Higgins JP, Thompson SG (2002) Quantifying heterogeneity in a meta-analysis. *Stat Med* **21**: 1539–1558.
- Howlader N, Noone A, Krapcho M, Neyman N, Aminou R, Waldron W, Altekruse S, Kosary C, Ruhl J, Tatalovich Z, Cho H, Mariotto A, Eisner M, Lewis D, Chen H, Feuer E, Cronin K (eds) (2012) *SEER Cancer Statistics Review, 1975-2009 (Vintage 2009 Populations)*. National Cancer Institute: Bethesda, MD, available at [http://seer.cancer.gov/csr/1975\\_2009\\_pops09/](http://seer.cancer.gov/csr/1975_2009_pops09/) based on November 2011 SEER data submission, posted to the SEER web site, April 2012).
- Hu J, La Vecchia C, DesMeules M, Negri E, Mery L. Canadian Cancer Registries Epidemiology Research G (2008) Nutrient and fiber intake and risk of renal cell carcinoma. *Nutr Cancer* **60**: 720–728.
- Hu J, Mao Y, DesMeules M, Csizmadia I, Friedenreich C, Mery L. Canadian Cancer Registries Epidemiology Research G (2009) Total fluid and specific beverage intake and risk of renal cell carcinoma in Canada. *Cancer Epidemiol* **33**: 355–362.
- Kellerer M, von Eye Corleta H, Muhlhofer A, Capp E, Mosthaf L, Bock S, Petrides PE, Haring HU (1995) Insulin- and insulin-like growth-factor-I receptor tyrosine-kinase activities in human renal carcinoma. *Int J Cancer* **62**: 501–507.
- Leitzmann MF (2011) Physical activity and genitourinary cancer prevention. *Recent Results Canc Res* **186**: 43–71.
- Lindblad P, Wolk A, Bergstrom R, Persson I, Adami HO (1994) The role of obesity and weight fluctuations in the etiology of renal cell cancer: a population-based case-control study. *Cancer Epidemiol Biomarkers Prev* **3**: 631–639.
- Liu Y, Hu F, Li D, Wang F, Zhu L, Chen W, Ge J, An R, Zhao Y (2011) Does physical activity reduce the risk of prostate cancer? A systematic review and meta-analysis. *Eur Urol* **60**: 1029–1044.
- Mahabir S, Leitzmann MF, Pietinen P, Albanes D, Virtamo J, Taylor PR (2004) Physical activity and renal cell cancer risk in a cohort of male smokers. *Int J Cancer* **108**: 600–605.
- McTiernan A (2008) Mechanisms linking physical activity with cancer. *Nat Rev Cancer* **8**: 205–211.
- Mellemegaard A, Engholm G, McLaughlin JK, Olsen JH (1994) Risk factors for renal-cell carcinoma in Denmark. III. Role of weight, physical activity and reproductive factors. *Int J Cancer* **56**: 66–71.
- Mellemegaard A, Lindblad P, Schlehofer B, Bergstrom R, Mandel JS, McCredie M, McLaughlin JK, Niwa S, Odaka N, Pommer W, Olsen JH (1995) International renal-cell cancer study. III. Role of weight, height, physical activity, and use of amphetamines. *Int J Cancer* **60**: 350–354.
- Menezes RJ, Tomlinson G, Kreiger N (2003) Physical activity and risk of renal cell carcinoma. *Int J Cancer* **107**: 642–646.
- Moher D, Liberati A, Tetzlaff J, Altman DG (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* **339**: b2535.
- Monninkhof EM, Elias SG, Vlems FA, van der Tweel I, Schuit AJ, Voskuil DW, van Leeuwen FE (2007) Physical activity and breast cancer: a systematic review. *Epidemiology* **18**: 137–157.
- Moore SC, Chow WH, Schatzkin A, Adams KF, Park Y, Ballard-Barbash R, Hollenbeck A, Leitzmann MF (2008) Physical activity during adulthood and adolescence in relation to renal cell cancer. *Am J Epidemiol* **168**: 149–157.
- Nicodemus KK, Sweeney C, Folsom AR (2004) Evaluation of dietary, medical and lifestyle risk factors for incident kidney cancer in postmenopausal women. *Int J Cancer* **108**: 115–121.
- O'Rourke MA, Cantwell MM, Cardwell CR, Mulholland HG, Murray LJ (2010) Can physical activity modulate pancreatic cancer risk? a systematic review and meta-analysis. *Int J Cancer* **126**: 2957–2968.
- Paffenbarger Jr RS, Hyde RT, Wing AL (1987) Physical activity and incidence of cancer in diverse populations: a preliminary report. *Am J Clin Nutr* **45**: 312–317.
- Pan SY, DesMeules M, Morrison H, Wen SW (2006) Obesity, high energy intake, lack of physical activity, and the risk of kidney cancer. *Cancer Epidemiol Biomarkers Prev* **15**: 2453–2460.
- Parent ME, Rousseau MC, El-Zein M, Latreille B, Desy M, Siemiatycki J (2011) Occupational and recreational physical activity during adult life and the risk of cancer among men. *Cancer Epidemiol* **35**: 151–159.
- Parker AS, Cerhan JR, Lynch CF, Ershov AG, Cantor KP (2002) Gender, alcohol consumption, and renal cell carcinoma. *Am J Epidemiol* **155**: 455–462.
- R Development Core Team (2011) *R: A Language and Environment for Statistical Computing*. R Development Core Team: Vienna, Austria.
- Rosenthal M, Haskell WL, Solomon R, Widstrom A, Reaven GM (1983) Demonstration of a relationship between level of physical training and insulin-stimulated glucose utilization in normal humans. *Diabetes* **32**: 408–411.
- Scelo G, Brennan P (2007) The epidemiology of bladder and kidney cancer. *Nat Clin Pract Urol* **4**: 205–217.
- Setiawan VW, Stram DO, Nomura AM, Kolonel LN, Henderson BE (2007) Risk factors for renal cell cancer: the multiethnic cohort. *Am J Epidemiol* **166**: 932–940.
- Spyridopoulos TN, Petridou ET, Dessypris N, Terzidis A, Skalkidou A, Deliveliotis C, Chrousos GP. Obesity, Cancer Oncology G (2009) Inverse association of leptin levels with renal cell carcinoma: results from a case-control study. *Hormones (Athens)* **8**: 39–46.
- Tavani A, Zucchetto A, Dal Maso L, Montella M, Ramazzotti V, Talamini R, Franceschi S, La Vecchia C (2007) Lifetime physical activity and the risk of renal cell cancer. *Int J Cancer* **120**: 1977–1980.
- Thompson AM, Church TS, Janssen I, Katzmarzyk PT, Earnest CP, Blair SN (2008) Cardiorespiratory fitness as a predictor of cancer mortality among men with pre-diabetes and diabetes. *Diabetes Care* **31**: 764–769.
- van Dijk BA, Schouten LJ, Kiemeny LA, Goldbohm RA, van den Brandt PA (2004) Relation of height, body mass, energy intake, and physical activity to risk of renal cell carcinoma: results from the Netherlands Cohort Study. *Am J Epidemiol* **160**: 1159–1167.
- Vatten LJ, Trichopoulos D, Holmen J, Nilsen TI (2007) Blood pressure and renal cancer risk: the HUNT Study in Norway. *Br J Cancer* **97**: 112–114.
- Viechtbauer W (2010) Conducting meta-analyses in R with the metafor package. *J Stat Softw* **36**: 1–48.
- Vincent KR, Vincent HK, Braith RW, Lennon SL, Lowenthal DT (2002) Resistance exercise training attenuates exercise-induced lipid peroxidation in the elderly. *Eur J Appl Physiol* **87**: 416–423.
- Voskuil DW, Monninkhof EM, Elias SG, Vlems FA, van Leeuwen FE (2007) Physical activity and endometrial cancer risk, a systematic review of current evidence. *Cancer Epidemiol Biomarkers Prev* **16**: 639–648.
- Washio M, Mori M, Sakauchi F, Watanabe Y, Ozasa K, Hayashi K, Miki T, Nakao M, Mikami K, Ito Y, Wakai K, Tamakoshi A, Group JS (2005) Risk factors for kidney cancer in a Japanese population: findings from the JACC Study. *J Epidemiol* **15**(Suppl 2): S203–S211.
- Wilson RT, Wang J, Chinchilli V, Richie JP, Virtamo J, Moore LE, Albanes D (2009) Fish, vitamin D, and flavonoids in relation to renal cell cancer among smokers. *Am J Epidemiol* **170**: 717–729.
- Wing RR (1999) Physical activity in the treatment of the adulthood overweight and obesity: current evidence and research issues. *Med Sci Sports Exerc* **31**: S547–S552.
- Yuen JS, Akkaya E, Wang Y, Takiguchi M, Peak S, Sullivan M, Protheroe AS, Macaulay VM (2009) Validation of the type 1 insulin-like growth factor receptor as a therapeutic target in renal cancer. *Mol Cancer Ther* **8**: 1448–1459.
- Yun YH, Lim MK, Won YJ, Park SM, Chang YJ, Oh SW, Shin SA (2008) Dietary preference, physical activity, and cancer risk in men: national health insurance corporation study. *BMC Cancer* **8**: 366.

This work is published under the standard license to publish agreement. After 12 months the work will become freely available and the license terms will switch to a Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported License.

## APPENDIX A

**Description of quality score**

To assess the methodologic quality of studies, we employed a quality score proposed by Voskuil *et al* (2007). The maximum score was 105. The score addressed selection bias (up to 42 points), misclassification bias (up to 42 points), and confounding bias (up to 21 points). Specifically, the score covered the following 19 items: (1) percentage lost to follow-up (cohort studies, up to 8 points)/percentage response (case-control studies, up to 8 points); (2) difference between percentage response among cases and controls (case-control studies: up to 6 points; cohort studies: 10 points by default); (3) percentage incident cases with known incidence date (*vs* inclusion of prevalent/fatal cases with unknown incidence date: up to 7 points, independent of study design); (4) cases and controls from the same population (case-control studies: up to 10 points; cohort studies: 10 points by default); (5) same exclusion criteria for cases and controls (case-control studies: up to 7 points; cohort studies: 7 points by default); (6) complete list of recreational physical activities (up to 4 points independent of study design); (7) assessment of total physical activity (up to 4 points); (8) assessment of physical activity intensity and frequency or assessment of physical activity intensity and duration (up to 5 points); (9) source of physical activity information (proxy/self-administered questionnaire/interview) (up to 4 points); (10) definition of physical activity score provided (up to 2 points); (11) examination of past physical activity (up to 4 points); (12) examination of change in physical activity (up to 4 points); (13) use of a valid or reliable physical activity assessment (up to 2 points); (14) minimisation of recall bias (up to 7 points); (15) valid cancer diagnosis (up to 4 points); (16) exclusion of benign cases (up to 2 points); (17) statistical adjustment for potential confounding variables (up to 4 points); (18) comprehensive adjustment for potential confounding variables (up to 9 points); (19) mutual adjustment for recreational and occupational physical activity (up to 8 points). With respect to item 18, we considered the well-established (Scelo and Brennan, 2007) renal cancer risk factors, such as smoking, obesity, hypertension, and type 2 diabetes mellitus, and awarded points based on how many of those factors had been included in the model (0 points for 0 factors, 4 points for 1 or 2 factors, and 9 points for 3 or 4 factors).

## APPENDIX B

**PubMed search strategy**

For the PubMed search, last performed on 30 September 2012, we pasted the following search terms all at once into the PubMed search command line:

(physical activity[title/abstract] OR exercise[title/abstract] OR cardiorespiratory fitness[title/abstract] OR cardiovascular fitness[title/abstract] OR resistance training[title/abstract] OR endurance training[title/abstract] OR aerobic[title/abstract] OR sport[title/abstract] OR athletes[title/abstract] OR players[title/abstract] OR lifestyle[title/abstract]) AND (kidney cancer[title] OR renal cancer[title] OR renal cell cancer[title] OR renal carcinoma[title] OR renal cell carcinoma[title] OR cancer[title])

NOT (lung\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (bronchial[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (breast\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (mamma\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (ovar\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT

(endometr\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (uter\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (cervi\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (gynecolog\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (prostat\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (testic\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (urinary\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (bladder\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (urothelial\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (colon\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (rectal\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (colorectal\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (bowel\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (\*digestive\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (gastric\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (stomach\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (oesophag\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (esophag\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (pancrea\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (tract\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (duct\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (tube\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (liver\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (hepato\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (gallbladder\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (oral\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (pharyn\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (nasopharyn\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (laryn\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (lymphoid\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (bone\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (head and neck\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (skin\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (melanoma\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (brain\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (thora\*[-title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (thyroid\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (squamous cell\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (basal cell\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (adenoma\*[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (multiple[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title]) NOT (mouth[title] NOT renal cancer[title] NOT renal cell cancer[title] NOT kidney cancer[title])

AND (risk[title/abstract] OR incidence[title/abstract] OR mortality[title/abstract])

NOT (surviv\*[title] OR prognosis\*[title] OR quality of life\*[title] OR fatigue\*[title] OR pallia\*[title] OR cancer patient\*[title] OR cancer care\*[title] OR recurrence\*[title] OR progression\*[title] OR clinical outcome\*[title] OR chemotherapy\*[title] OR radiation\*[title] OR radiotherapy\*[title] OR rehabilitation\*[title] OR recovery\*[title] OR cancer diagnosis[title] OR cancer treatment[title] OR cancer surgery[title] OR with cancer[title])

AND English[lang]

NOT (review[ptyp] OR meta-analysis[ptyp] OR editorial[ptyp] OR comment[ptyp] OR letter[ptyp] OR guideline[ptyp] OR news[ptyp]) AND humans[MeSH Terms]

## APPENDIX C

### Web of Knowledge search strategy

For the Web of Knowledge search, last performed on 30 September 2012, we specified the following search steps:

#STEP 1 SET LEMMATIZATION = OFF (ADJUST YOUR SEARCH SETTINGS)

#STEP 2 PUT THE FOLLOWING TERMS INTO TOPIC SEARCH

(physical activity OR exercise OR cardiorespiratory fitness OR cardiovascular fitness OR resistance training OR endurance training OR aerobic OR sport OR athletes OR players OR lifestyle) AND (kidney cancer OR renal cancer OR renal cell cancer OR renal carcinoma OR renal cell carcinoma OR cancer)

AND (risk OR incidence OR mortality)

#STEP 3 PUT THE FOLLOWING TERMS INTO TITLE SEARCH

(kidney cancer OR renal cancer OR renal cell cancer OR renal carcinoma OR renal cell carcinoma OR cancer) NOT (lung\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (bronchial NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (breast\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (mamma\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (ovar\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (endometr\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (uter\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (cervi\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (gynecolog\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (prostat\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (testic\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (urinary\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (bladder\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (urothelial\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (colon\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT

(rectal\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (colorectal\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (bowel\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (\*digestive\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (gastric\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (stomach\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (oesophag\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (esophag\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (pancrea\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (tract\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (duct\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (tube\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (liver\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (hepato\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (gallbladder\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (oral\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (pharyn\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (nasopharyn\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (laryn\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (lymphoid\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (bone\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (head and neck\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (skin\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (melanoma\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (brain\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (thora\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (thyroid\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (squamous cell\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (basal cell\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (adenoma\* NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (multiple NOT renal cancer NOT renal cell cancer NOT kidney cancer) NOT (mouth NOT renal cancer NOT renal cell cancer NOT kidney cancer)

NOT (surviv\* OR prognosis\* OR quality of life\* OR fatigue\* OR pallia\* OR cancer patient\* OR cancer care\* OR recurrence\* OR progression\* OR clinical outcome\* OR chemotherapy\* OR radiation\* OR radiotherapy\* OR rehabilitation\* OR recovery\* OR cancer diagnosis OR cancer treatment OR cancer surgery OR with cancer)

#STEP 4 SET RESEARCH AREA TO PUBLIC ENVIRONMENTAL OCCUPATIONAL HEALTH

#STEP 5 EXCLUDE ALL DOCUMENT TYPES APART FROM ARTICLE

#STEP 6 SET LANGUAGE TO ENGLISH