

Reply to 'Comment on 'Distinct clinical outcomes of two CIMP-positive colorectal cancer subtypes based on a revised CIMP classification system''

Jeong Mo Bae^{1,2}, Jung Ho Kim^{1,3}, Yoonjin Kwak^{3,4}, Dae-Won Lee⁵, Yongjun Cha⁵, Xianyu Wen¹, Tae Hun Lee¹, Nam-Yun Cho¹, Seung-Yong Jeong⁶, Kyu Joo Park⁶, Sae Won Han⁵, Hye Seung Lee^{3,4}, Tae-You Kim⁵ and Gyeong Hoon Kang^{*1,3}

¹Laboratory of Epigenetics, Cancer Research Institute, Seoul National University College of Medicine, Seoul 03080, South Korea; ²Department of Pathology, SMG-SNU Boramae Medical Center, Seoul 07061, South Korea; ³Department of Pathology, Seoul National University College of Medicine, Seoul 03080, South Korea; ⁴Department of Pathology, Seoul National University Bundang Hospital, Seongnam 13620, South Korea; ⁵Department of Internal Medicine, Seoul National University Hospital, Seoul 03080, South Korea and ⁶Department of Surgery, Seoul National University Hospital, Seoul 03080, South Korea

Sir,

We thank Tapial *et al*, 2017 for their interest in our recent study. The authors applied our revised CIMP classification to their three different clinical data sets, which are composed of patients with early-onset colorectal cancer (EOCRC) (younger than 45 years), patients with late-onset CRC (LOCRC) (older than 70 years), and individuals diagnosed with synchronous CRC (SCRC). They addressed that only LOCRC cases showed similar tendency of increasing *BRAF* mutation, MSI-high and *MLH1* methylation along with the increase in the number of methylated genes. Moreover, they insisted that prognostic results were only partially confirmed in SCRC.

Non-linearity of molecular alterations including *BRAF* mutation, MSI-high and *MLH1* methylation in EOCRC and SCRC might originate from two reasons. First, EOCRC and SCRC have strong germline predispositions to CRC, even though they are not either familial adenomatous polyposis or Lynch syndrome (Cybulski *et al*, 2014; de Voer *et al*, 2016). Recent studies revealed that germline predisposition in EOCRC is greater than expected (Pearlman *et al*, 2017); these germline predispositions are mainly associated with chromosomal instability rather than CIMP (Chan *et al*, 2001; McGivern *et al*, 2004). Second, CIMP-P2 CRCs usually occur in elderly patients. Two recent studies showed similar tendency in CRCs with *BRAF* mutation and concurrent MSI or *MLH1* methylation (Seppala *et al*, 2015; Vedeld *et al*, 2017).

The authors tried to validate prognostic value of our revised CIMP classification in their EOCRC, LOCRC and SCRC subgroups. However, considering their previous publication, the sample size of each subgroup is too small to get enough statistical power for the subgroup of low prevalence, such as CIMP-P1 and CIMP-P2 (Ogino *et al*, 2011; Perea *et al*, 2015; Arriba *et al*, 2017). Considering the low prevalence of EOCRC and SCRC, a multi-centre study might be necessary to validate the prognostic value of our revised CIMP classification.

Overall, Tapial *et al*'s results emphasise the fact that EOCRC and SCRC have different molecular landscapes compared with sporadic CRCs. Further comprehensive study might shed light on the complex interaction between germline predisposition, accumulation of somatic mutation and epigenetic alteration.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

Arriba M, Sanchez R, Rueda D, Gomez L, Garcia JL, Rodriguez Y, Pajares JA, Perez J, Urioste M, Sarmiento RG, Perea J (2017) Toward a molecular

- classification of synchronous colorectal cancer: clinical and molecular characterization. *Clin Colorectal Cancer* **16**(1): 31–37.
- Chan TL, Curtis LC, Leung SY, Farrington SM, Ho JW, Chan AS, Lam PW, Tse CW, Dunlop MG, Wyllie AH, Yuen ST (2001) Early-onset colorectal cancer with stable microsatellite DNA and near-diploid chromosomes. *Oncogene* **20**(35): 4871–4876.
- Cybulski C, Nazari S, Narod SA (2014) Multiple primary cancers as a guide to heritability. *Int J Cancer* **135**(8): 1756–1763.
- de Voer RM, Hahn MM, Weren RD, Mensenkamp AR, Gilissen C, van Zelst-Stams WA, Spruijt L, Kets CM, Zhang J, Venselaar H, Vreede L, Schubert N, Tychon M, Derks R, Schackert HK, Geurts van Kessel A, Hoogerbrugge N, Ligtenberg MJ, Kuiper RP (2016) Identification of novel candidate genes for early-onset colorectal cancer susceptibility. *PLoS Genet* **12**(2): e1005880.
- McGivern A, Wynter CV, Whitehall VL, Kambara T, Spring KJ, Walsh MD, Barker MA, Arnold S, Simms LA, Leggett BA, Young J, Jass JR (2004) Promoter hypermethylation frequency and *BRAF* mutations distinguish hereditary non-polyposis colon cancer from sporadic MSI-H colon cancer. *Fam Cancer* **3**(2): 101–107.
- Ogino S, Chan AT, Fuchs CS, Giovannucci E (2011) Molecular pathological epidemiology of colorectal neoplasia: an emerging transdisciplinary and interdisciplinary field. *Gut* **60**(3): 397–411.
- Pearlman R, Frankel WL, Swanson B, Zhao W, Yilmaz A, Miller K, Bacher J, Bigley C, Nelsen L, Goodfellow PJ, Goldberg RM, Paskett E, Shields PG, Freudenheim JL, Stanich PP, Lattimer I, Arnold M, Liyanarachchi S, Kalady M, Heald B, Greenwood C, Paquette I, Prues M, Draper DJ, Lindeman C, Kuebler JP, Reynolds K, Brell JM, Shaper AA, Mahesh S, Buie N, Weeman K, Shine K, Haut M, Edwards J, Bastola S, Wickham K, Khanduja KS, Zacks R, Pritchard CC, Shirts BH, Jacobson A, Allen B, de la Chapelle A, Hampel H (2017) Prevalence and Spectrum of germline cancer susceptibility gene mutations among patients with early-onset colorectal cancer. *JAMA Oncol* **3**(4): 464–471.
- Perea J, Cano JM, Rueda D, Garcia JL, Inglada L, Osorio I, Arriba M, Perez J, Gaspar M, Fernandez-Miguel T, Rodriguez Y, Benitez J, Gonzalez-Sarmiento R, Urioste M (2015) Classifying early-onset colorectal cancer according to tumor location: new potential subcategories to explore. *Am J Cancer Res* **5**(7): 2308–2313.
- Seppala TT, Bohm JP, Friman M, Lahtinen L, Vayrynen VM, Liipo TK, Ristimaki AP, Kairaluoma MV, Kellokumpu IH, Kuopio TH, Mecklin JP (2015) Combination of microsatellite instability and *BRAF* mutation status for subtyping colorectal cancer. *Br J Cancer* **112**(12): 1966–1975.
- Tapial S, Rueda D, Arriba M, Garcia JL, Brandariz L, Perez J, Rodriguez Y, Garcia-Olmo D, Gonzalez-Sarmiento R, Urioste M, Perea J (2017) Comment on 'Distinct clinical outcomes of two CIMP-positive colorectal cancer subtypes based on a revised CIMP classification system'. *Br J Cancer*; e-pub ahead of print 2 January 2018; doi:10.1038/bjc.2017.343.
- Vedeld HM, Merok M, Jeanmougin M, Danielsen SA, Honne H, Presthus GK, Svindland A, Sjo OH, Hektoen M, Eknaes M, Nesbakken A, Lothe RA, Lind GE (2017) CpG island methylator phenotype identifies high risk patients among microsatellite stable *BRAF* mutated colorectal cancers. *Int J Cancer* **141**(5): 967–976.

This work is published under the BJC's standard license to publish agreement. After 12 months the license terms will change to a Creative Commons AttributionNonCommercial-Share Alike 4.0 Unported License.

*Correspondence: GH Kang; E-mail: ghkang@snu.ac.kr

Published online 2 January 2018

© 2018 Cancer Research UK. All rights reserved 0007–0920/18