

Phospho- Δ Np63 α is a key regulator of the cisplatin-induced microRNAome in cancer cells

Y Huang^{1,3}, A Chuang^{1,3}, H Hao², C Talbot², T Sen³, B Trink³, D Sidransky³ and E Ratovitski^{*1}

Head and neck squamous cell carcinoma (HNSCC) cells exposed to cisplatin (CIS) displayed a dramatic ATM-dependent phosphorylation of Δ Np63 α that leads to the transcriptional regulation of downstream mRNAs. Here, we report that phospho (p)- Δ Np63 α transcriptionally deregulates miRNA expression after CIS treatment. Several p- Δ Np63 α -dependent microRNA species (miRNAs) were deregulated in HNSCC cells upon CIS exposure, including miR-181a, miR-519a, and miR-374a (downregulated) and miR-630 (upregulated). Deregulation of miRNA expression led to subsequent modulation of mRNA expression of several targets (TP53-S46, HIPK2, ATM, CDKN1A and 1B, CASP3, PARP1 and 2, DDIT1 and 4, BCL2 and BCL2L2, TP73, YES1, and YAP1) that are involved in the apoptotic process. Our data support the notion that miRNAs are critical downstream targets of p- Δ Np63 α and mediate key pathways implicated in the response of cancer cells to chemotherapeutic drugs.

Cell Death and Differentiation (2011) 18, 1220–1230; doi:10.1038/cdd.2010.188; published online 28 January 2011

Cisplatin (CIS) is a chemotherapeutic agent often used in the treatment of human cancers because of its ability to induce cell death in neoplastic cells.¹ However, tumors eventually become resistant to platinum chemotherapy.^{1,2} CIS induces DNA damage, leading to the accumulation of activated members of the tumor protein (TP) 53 family (TP53, TP63, and TP73).^{3–7} When induced, TP53, TP63, and TP73 alter the transcription of a large set of downstream target genes, controlling cell-cycle arrest, cell death, increased DNA repair, inhibition of angiogenesis, and so on.^{8–12} MicroRNA species (miRNAs) are small 18–24-nucleotide non-coding RNAs that act through the RNA interference pathway and repress target gene expression largely by modulating translation and mRNA stability.^{13,14} miRNA precursors (pri-miRNAs) are processed in the nucleus, and the hairpin products are then cleaved by the double-stranded ribonuclease, DICER1, in the cytoplasm to generate mature miRNAs.^{14,15} Altered expression of miRNA genes has been found in a variety of tumor types and has shown the oncogenic, tumor-suppressive, or apoptotic potential of specific miRNAs.^{16–19} Certain miRNAs were shown to mediate the induction of cell death, cell cycle arrest, and senescence and contribute to epithelial stem cell maturation.^{18–20}

We previously observed that the head and neck squamous cell carcinoma (HNSCC) cells exposed to CIS displayed a dramatic downregulation of Δ Np63 α through an ATM-dependent phosphorylation mechanism.^{21,22} We also showed that phospho (p)- Δ Np63 α is critical for the transcriptional

regulation of downstream mRNAs in HNSCC cells.^{21,22} In the current study, we present evidence that p- Δ Np63 α regulates miRNA expression in CIS-treated HNSCC cells through both transcriptional and post-transcriptional mechanisms.

Results

CIS induces the p- Δ Np63 α -dependent expression of DICER1. We employed the JHU-029 stable clones, which have been shown to exclusively produce wild-type Δ Np63 α or Δ Np63 α -S385G with an altered ability to be phosphorylated by ATM kinase.^{21,22} Using mRNA expression and ChIP-on-chip arrays, we previously showed that exposure of wild-type Δ Np63 α cells to 10 μ g/ml CIS for 16 h led to an approximately fivefold upregulation of the double-stranded RNA-specific endoribonuclease, DICER1.²² We found here that upon CIS exposure, the DICER1 expression level is higher (~4.0- to 4.2-fold) in wild-type Δ Np63 α cells than in Δ Np63 α -S385G cells (Figure 1).

CIS was previously shown to induce binding of the NF- κ B/p- Δ Np63 α protein complexes to the CCAAT promoter elements.²² The specific CCAAT elements (1, 2, and 3) along with the responsive elements (REs) for p63 (see ref. 23) were found in the 2700-bp human DICER1 promoter identified in the UCSC server using the TFSEARCH software (<http://www.cbrc.jp/research/db/TFSEARCH.html>; Computational Biology Research Center, Parallel Application Laboratory, RWCP, Tokyo, Japan). A few cognate REs for various

¹Department of Dermatology, Johns Hopkins University School of Medicine, Baltimore, MD, USA; ²Functional Genomics/Proteomics Core Facility, Johns Hopkins University School of Medicine, Baltimore, MD, USA; ³Department of Otolaryngology/Head and Neck Surgery, Johns Hopkins University School of Medicine, Baltimore, MD, USA

*Corresponding author: E Ratovitski, Department of Dermatology, Johns Hopkins University School of Medicine, Cancer Research Building II, Room 2M05, 1550 Orleans Street, Baltimore, MD 21231, USA. Tel: +1 410 502 5130; Fax: +1 410 614 1411; E-mail: eratovi1@jhmi.edu

Keywords: p63; cisplatin; squamous cell carcinomas; DICER1; microRNA

Abbreviations: HNSCC, head and neck squamous cell carcinoma; ATM, ataxia-telangiectasia mutated; DDIT, DNA-damage-inducible transcript; CIS, cisplatin; ChIP, chromatin immunoprecipitation; IP, immunoprecipitation; PARP, poly(ADP-ribose) polymerase; BCL, B-cell CLL/lymphoma; NF, nuclear factor; CDKN, cyclin-dependent kinase inhibitor; HIPK, homeodomain interacting protein kinase; YES, Yamaguchi sarcoma viral oncogene homolog; YAP, YES-associated protein; HMG, high-mobility group; PCR, polymerase chain reaction; RISC, RNA-induced silencing microprocessor complex; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; p, phospho

Received 2.7.10; revised 17.12.10; accepted 17.12.10; Edited by M Oren; published online 28.1.11




Figure 1 p- Δ Np63 α upregulates DICER1 expression upon CIS exposure. Wild-type Δ Np63 α cells (p63wt) or Δ Np63 α -S385G cells (p63mut) were exposed to Con (–) or 10 μ g/ml CIS (+) for 24 h. Total RNA and protein were isolated and analyzed for DICER1 expression. (a) PCR assay. (b) qPCR assay of DICER1 expression. Data were normalized against GAPDH levels and plotted as relative units (RU), with measurements obtained from wild-type Δ Np63 α cells treated with Con set as 1. Experiments were performed in triplicate with \pm S.D. as indicated ($P < 0.01$). (c) Immunoblotting with anti-DICER1 antibodies (levels of DICER1 were quantified and normalized against β -actin protein level, and the values obtained from untreated wild-type Δ Np63 α cells are designated as (1)). As loading controls, we used the GAPDH mRNA levels for PCR and the β -actin protein levels for immunoblotting

transcription factors (e.g. E2F, C/EBP β , STAT, Oct-1, and P53) were found present in the DICER1 promoter (Supplementary Figure S1). Using antibodies against both Δ Np63 and p- Δ Np63 α for chromatin immunoprecipitation (ChIP) assay, we found that Δ Np63 α (in its phosphorylated form) binds to the NF-Y-REs, CCAAT elements 1 and 2 of the DICER1 promoter, whereas no detectable binding was found to the nonspecific region (Figure 2a). We further examined the effect of endogenous p- Δ Np63 α on the DICER1 (DCR) promoter (+49 to –871, containing p63RE and CCAAT element 3, Supplementary Figure S1) in wild-type Δ Np63 α and Δ Np63 α -S385G cells by monitoring luciferase activity from the pDCR-Luc reporter plasmid. We found that the CIS exposure led to a dramatic \sim 2.7-fold increase in the DCR-driven luciferase activity in wild-type Δ Np63 α cells, whereas no significant changes were observed in Δ Np63 α -S385G cells (Figure 2b, graph panel, sample 3 *versus* 2). Next, wild-type Δ Np63 α and Δ Np63 α -S385G cells were transfected with the exogenous Δ Np63 α -S385G-FL and Δ Np63 α -FL constructs, respectively. In contrast to Δ Np63 α -S385G-FL, Δ Np63 α -FL has undergone the CIS-induced phosphorylation, which was detected with

the anti-p- Δ Np63 α antibody (Figure 2b, immunoblot panel). We further showed that the competition of exogenous Δ Np63 α -S385G with endogenous p- Δ Np63 α decreased the CIS-mediated DCR-Luc activity by \sim 2.3-fold in wild-type Δ Np63 α cells (Figure 2b, graph panel, sample 5 *versus* 3). However, exogenous p- Δ Np63 α -FL increased the CIS-mediated DCR-Luc activity in Δ Np63 α -S385G cells, supporting the competition of exogenous p- Δ Np63 α with endogenous Δ Np63 α -S385G (Figure 2b, graph panel, sample 10 *versus* 8). Altogether, these data strongly support that the DICER1 promoter is a potential transcriptional target for p- Δ Np63 α in HNSCC cells upon CIS exposure.

CIS deregulates the p- Δ Np63 α -dependent expression of miRNAs. Given that DICER1 is a well-known regulator of miRNA biosynthesis and has a central role in the maturation of the miRNAs that contribute to apoptosis¹⁵ and tumor suppression,²⁴ we examined the miRNA signature of HNSCC cells exposed to CIS. First, we tested whether the exposure of HNSCC cells to CIS affects the expression of specific miRNAs. Second, we questioned whether p- Δ Np63 α is involved in transcriptional regulation of certain miRNAs. We suggest that any overlapping results between these two sets of experiments may indicate which miRNAs are induced by CIS through a p- Δ Np63 α -dependent mechanism.

To answer the first question, we exposed wild-type Δ Np63 α cells to control medium (Con) or 10 μ g/ml CIS for 24 h. Using total RNA from both cell lines, we performed a miRNA-chip microarray analysis using a chip containing probes to 842 human miRNAs.²⁵ We found that CIS downregulated seven miRNAs (miR-519a, miR-181a, miR-374a, miR-29c, miR-98, miR-22, and miR-18b) from –1.7- to 3.7-fold, and upregulated eight miRNAs (miR-760, miR-185, miR-574, miR-453, miR-297, miR-194, miR-885-3p, and miR-630) from +1.8- to +4.9-fold in HNSCC cells (Supplementary Table S1).

To answer the second question, we utilized the isogenic wild-type Δ Np63 α and Δ Np63 α -S385G cells.^{21,22} Both cell lines were treated with CIS, which has previously been shown to increase the p- Δ Np63 α levels.^{21,22} Using the miRNA array chip, we thus found dramatic differences in the miRNA expression levels (Supplementary Table SII). miRNAs exhibiting a threefold or greater change in expression were chosen for further study. After CIS exposure, \sim 20 miRNA species were upregulated in wild-type Δ Np63 α cells (ranging from 3.3- to 7.4-fold, Supplementary Table SII) when compared with Δ Np63 α -S385G cells. At the same time, \sim 33 miRNAs were downregulated (ranging from 3.4- to 19.2-fold, Supplementary Table SII) in wild-type Δ Np63 α cells compared with Δ Np63 α -S385G cells after CIS exposure. We found that certain miRNAs were common to both sets (e.g. miR-519a, miR-181a, miR-374a, miR-885-3p, and miR-630), suggesting that these miRNAs were regulated by a CIS-mediated/p- Δ Np63 α -dependent transcriptional mechanism. We found no significant differences in miRNA levels between wild-type Δ Np63 α cells and Δ Np63 α -S385G cells treated with Con (data not shown).

We further tested the pri-miRNA levels in wild-type Δ Np63 α cells upon CIS exposure using qPCR analysis. We thus found that the precursors for miR-630, miR-194, miR-297, miR-885-3p, miR-574, miR-185, and miR-760 were upregulated in




Figure 2 p- Δ Np63 α binds to the DICER1 promoter sequences and activates the DICER1 promoter activity upon CIS exposure. Wild-type Δ Np63 α cells were exposed to Con or 10 μ g/ml CIS for 24 h. (a) ChIP assay for the DICER1 promoter was performed using both anti- Δ Np63 and anti-p- Δ Np63 α antibodies. A rabbit immunoglobulin (IgG) was used as a negative control for ChIP. CCAAT elements 1 and 2 were amplified using primers covering the specific regions (–1892 to –1191) and (–1421 to –721), respectively, and yielding the 700-bp PCR fragments. Positive controls (Inputs) and a negative control (ChIP using primers for the DICER1 promoter nonspecific region –2639 to –2301 yielding the 340-bp PCR product) were shown. (b) Luciferase reporter activity assay. Wild-type and mutated cells were transfected with 100 ng of the promoterless pGL3 plasmid or pGL3-DCR-Luc plasmid along with 1 ng of the Renilla luciferase plasmid. Wild-type Δ Np63 α cells were also transfected with 100 ng of the Δ Np63 α -S385G-FL expression cassette, whereas Δ Np63 α -S385G cells were transfected with 100 ng of the Δ Np63 α -FL expression cassette, as indicated. Cells were exposed to Con and 10 μ g/ml CIS for 24 h. Luciferase reporter assays were conducted in triplicate (\pm S.D., $P < 0.01$). Firefly luciferase activity values were normalized against Renilla luciferase values and resulting values obtained from wild-type Δ Np63 α cells with the promoterless pGL3 plasmid and exposed to Con were designated as 1, while fold changes are shown as numerical values. Samples were also tested by immunoblotting with the indicated antibodies

wild-type Δ Np63 α cells upon CIS exposure (Figure 3a), whereas precursors for miR-29c, miR-519a, miR-181a, miR-374a, miR-98, miR-22, and miR-18b were downregulated (Figure 3b). We then found that mature miR-630, miR-194, miR-297, miR-885-3p, miR-574, miR-185, and miR-760 were upregulated (Figure 3c) to a greater extent than their pri-mRNAs (Figure 3a) in wild-type Δ Np63 α cells upon CIS exposure. However, mature miR-29c, miR-519a, miR-181a,

miR-374a, miR-98, miR-22, and miR-18b were downregulated (Figure 3d) to a lesser degree than their pri-mRNA (Figure 3b) in wild-type Δ Np63 α cells upon CIS exposure.

Interestingly, the mature miRNA/pri-miRNA ratio for individual miRNAs affected by CIS exposure varied as follows: 1.49 for miR-630, 1.42 for miR-194, 1.37 for miR-297, 1.39 for miR-453, 1.59 for miR-885-3p, 1.36 for miR-574, 1.26 for miR-185 and 1.02 for miR-760 (Figures 3a and c), and




Figure 3 CIS modulates the expression of the p- Δ Np63 α -dependent miRNAs. Wild-type Δ Np63 α cells were exposed to Con or 10 μ g/ml CIS for 24 h. Precursor pri-miRNA (a and b) and mature miRNA (c–e) levels were examined by qPCR assay using TaqMan analysis, and the data were plotted as relative units (RU), using the measurements obtained from the cells treated with Con as 1. Fold changes are presented as numerical values. Experiments were performed in triplicate with \pm S.D. as indicated (< 0.05). (a) Upregulated miRNAs (miR-630, miR-194, miR-297, miR-453, miR-885-3p, miR-574, miR-185, and miR-760). (b) Downregulated miRNAs (miR-29c, miR-519a, miR-181a, miR-374a, miR-98, miR-22, and miR-18b). (c) Upregulated miRNAs (miR-630, miR-194, miR-297, miR-453, miR-885-3p, miR-574, miR-185, and miR-760). (d) Downregulated miRNAs (miR-29c, miR-519a, miR-181a, miR-374a, miR-98, miR-22, and miR-18b). (e and f) Δ Np63 α -S385G cells were transfected with an empty pCMV-FL vector, Δ Np63 α -FL and Δ Np63 α -S385G-FL expression constructs. Resulting cells were exposed to 10 μ g/ml CIS for 24 h. (e) Immunoprecipitation (IP) of FLAG-tagged Δ Np63 α (wt) and (S385G) proteins using anti-FLAG antibody. Immunoblotting with antibodies against the FLAG epitope, Δ Np63 or p- Δ Np63 α . (f) Expression levels of indicated mature miRNA (miR-181a, miR-519a, miR-374a and miR-630) were examined by qPCR assay

1.39 for miR-29c, 1.44 for miR-22, 1.41 for miR-18b, 1.41 for miR-98, 1.31 for miR-374a, 1.35 for miR-519a, and 1.78 for miR-181a (Figures 3b and d). These data suggest that the CIS-induced DICER1 upregulation (Figure 1) is likely to contribute to these intriguing differences in expression levels of pri-miRNAs and mature miRNAs. Although the qPCR data (Figures 3a–d) supported the notion that CIS exposure leads to modulation of certain miRNAs in HNSCC cells, direct evidence of p- Δ Np63 α 's contribution to the modulation of miRNA expression was missing. Δ Np63 α -S385G cells were chosen (to avoid an endogenous p- Δ Np63 α background) for a subsequent transfection with an empty pCMV-FL vector and Δ Np63 α -FL and Δ Np63 α -S385G-FL

constructs. Cells were exposed to 10 μ g/ml CIS for 24 h to ensure the phosphorylation of exogenous Δ Np63 α -FL, and the miRNA expression was then quantified using qPCR. We showed that Δ Np63 α -FL (recognized by anti-p- Δ Np63 α antibody, Figure 3e) decreased the miR-181a, miR-519a, and miR-374a levels, while it increased the miR-630 level (Figure 3f). However, Δ Np63 α -S385G-FL (not recognized by anti-p- Δ Np63 α antibody, Figure 3e) failed to change the expression levels of these miRNAs compared with control vector (Figure 3f).

p- Δ Np63 α transcriptionally regulates miRNA expression upon CIS exposure. Using ChIP analysis, we found that




Figure 4 ChIP analysis of the p- Δ Np63 α protein binding to the miRNA promoters upon CIS exposure. Wild-type Δ Np63 α cells were exposed to Con or 10 μ g/ml CIS for 24 h. Samples were incubated with the anti-p- Δ Np63 α antibody. Precipitated DNAs were amplified with primers for the specific region (a, upper panel, 250-bp PCR product) and nonspecific region (a, lower panel, 130-bp PCR product) for the indicated miRNA promoters. (b) qPCR assay. Using the same primers for specific regions as referred to in Supplementary Figures S2–S6, we assessed the quantitative changes in the p- Δ Np63 α protein binding to the indicated miRNA promoter sequences after exposure of cells to Con or 10 μ g/ml CIS for 24 h. qPCR experiments were performed in triplicate with \pm S.D. as indicated (<0.05). Values obtained with Con were designated as 1, and fold changes are presented as numerical values

p- Δ Np63 α binds to the promoter sequences of certain miRNAs (miR-181a, miR-519a, miR-374a, miR-630, and miR-885-3p) containing the p63RE and CCAAT elements (Figure 4a, upper panel and Supplementary Figures S2–S6), whereas no detectable binding was observed in nonspecific regions of the miRNA promoters (Figure 4a, lower panel). By qPCR assay, we further showed that the CIS exposure induced binding of p- Δ Np63 α to specific regions of the miRNA promoters (miR-181a, miR-519a, miR-374a, miR-630, and miR-885-3p) to various extents (Figure 4b). We then examined whether global Δ Np63 α or p- Δ Np63 α was responsible for the transcriptional regulation of miRNA expression in wild-type Δ Np63 α cells and Δ Np63 α -S385G cells. Using qPCR assay, we quantified the ChIP-PCR data obtained after precipitation of the specific promoter regions with antibodies against both Δ Np63 and p- Δ Np63 (Supplementary Figure S7). ChIP assays performed on wild-type Δ Np63 α cells with both antibodies showed that the CIS treatment dramatically induced Δ Np63 α binding to the promoters of DICER1, miR-630, and miR-885-3p (Supplementary Figure 7SA), and miR-181a, miR-519a, and miR-374a (Supplementary Figure 7B). However, ChIP assays performed on Δ Np63 α -S385G cells showed no such increase in binding of Δ Np63 α -S385G to the same promoter sequences (Supplementary Figures S7C and D), suggesting that the altered ability of Δ Np63 α to be phosphorylated by ATM kinase substantially impaired the capacity of Δ Np63 α to bind specific promoter sequences.

Thus, these data support the notion that p- Δ Np63 α is likely to regulate transcription of specific miRNAs in HNSCC cells upon CIS exposure.

DICER1 knockdown attenuates the CIS-mediated response. Using DICER1 siRNA, we next substantially knocked down the DICER1 expression (Figure 5a) in wild-type Δ Np63 α cells exposed to the Con or 10 μ g/ml of CIS for 24 h.

We further examined the effect of the DICER1 silencing on the cell survival assessed by MTT assay and showed that the CIS-mediated DICER1 increase led to a dramatic cell death (Figure 5b), whereas DICER1 siRNA almost entirely rescued wild-type Δ Np63 α cells from CIS-induced cell death (Figure 5b). FACS analysis further showed that the CIS exposure led to a marked increase in G1 cell population and a corresponding decrease of G2/M cells at 48 h after transfection with scrambled siRNA (Figure 5c). However, when cells were transfected with DICER1 siRNA, G1 cell population decreased, whereas G2/M cell population increased, suggesting the role for DICER1 in cell cycle arrest regulation (Figure 5c).

We next showed that DICER1 siRNA dramatically upregulated miR-630 and miR-885-3p compared with a scrambled siRNA (Figure 5d). This finding suggests a cumulative or synergistic effect of the DICER1-dependent miRNA maturation after the p- Δ Np63 α -dependent miRNA transcriptional activation. However, the expression levels of miR-181a, miR-519a, or miR-374a appeared to be minimally affected by siRNA against DICER1 (Figure 5e), suggesting an opposing (antagonistic) effect of DICER1-dependent miRNA maturation after the p- Δ Np63 α -dependent miRNA transcriptional repression.

Potential mRNA targets of the p- Δ Np63 α -dependent miRNAs under CIS treatment. Taking into consideration that miRNAs bind to the 3'-untranslated region (UTR) 'seed' sequences of mRNA, leading to changes in their expression levels, we examined the potential mRNA targets of tested miRNAs.²⁶ Using the pMiRTarget-luciferase reporter assay, we further found that several mRNA 3'-UTRs acted as direct targets of tested miRNAs, including those for DICER1 (miR-519a, miR-374a), DDIT1 (miR-374a) and DDIT4 (miR-181a), YES1 (miR-519a), homeodomain-interacting protein kinase-2 (HIPK2) (miR-181a), ATM (miR-181a, miR-374a), BCL2 (miR-630), BCL2L2 (miR-630), and YAP1 (miR-630). Corresponding miRNA mimics were shown to repress the luciferase reporter activities driven by the specific pMiR-Target plasmids (Supplementary Figures S8–S16). Moreover, CIS exposure led to an increase in the luciferase reporter activities of the pMiR-Target plasmids modulated by miR-181a, miR-519a, and miR-374a. Meanwhile, there was a decrease in the luciferase activities of the pMiR-Target plasmids, which were modulated by miR-630 (Supplementary Figures S8–S16). Using qPCR assay, we further observed that the CIS exposure and inhibitors for miR-181a, miR-519a, miR-374a increased the DDIT4, YES1 and DDIT1 mRNA levels, whereas CIS treatment and the miR-630 mimic decreased BCL2 mRNA levels (Supplementary Figure S17).




Figure 5 DICER1 silencing modulates the expression of the p- Δ Np63 α -dependent miRNAs upregulated after CIS exposure. Wild-type Δ Np63 α cells were transfected with scrambled siRNA (Scr) and DICER1 (DCR) siRNA for 24 h and exposed to Con (–, Con) or 10 μ g/ml CIS (+, CIS) for an additional 24 h. (a) DICER1 expression was analyzed by immunoblotting with indicated antibodies. Borders between images indicate that the data were obtained from separate experiments. Loading levels were assessed with anti- β -actin antibody. (b) Cell survival assay. (c) FACS analysis of cells. (d) Expression of mature miR-630 and miR-885-3p. (e) Expression of mature miR-181a, miR-519a and miR-374a. miRNA levels were examined by qPCR assay using TaqMan analysis and the data were plotted as relative units (RU). Values obtained with Con (–, Con) were designated as 1, with fold changes presented as numerical values. Experiments were performed in triplicate with \pm S.D. as indicated (<0.05)

CIS induces p53-dependent and caspase-dependent apoptosis through deregulation of p- Δ Np63 α -dependent miRNAs. To further examine the miRNA-dependent mRNA targets that were potentially affected by CIS, we tested the expression of these targets at the protein level. Wild-type Δ Np63 α cells were transfected with control (mock), mimic, or inhibitor for miR-181a, miR-519a, and miR-374a for 24 h and exposed to Con (–) and 10 μ g/ml CIS (+) for an additional 24 h and examined by immunoblotting. We found that the CIS exposure induced the accumulation of total TP53, S46-p-TP53, PARP2, CDKN1B, and DDIT4 (Figure 6a), CDKN1A, PARP1, YES1, and TP73 (Figure 7a), and DDIT1 and 4, YES1, and CASP3 (Figure 8a). In contrast to miR-181a, miR-519a, and miR-374a inhibitors and the mock control, mimics for miR-181a, miR-519a, and miR-374a decreased the levels of these cell cycle arrest and pro-apoptotic markers in wild-type Δ Np63 α cells (Figures 6a, 7a and 8a).

We further assessed the ability of specific miRNAs to modulate cell survival. Wild-type Δ Np63 α cells transfected with mimic or inhibitors for specific miRNAs were exclusively exposed to Con. Cells transfected with the mock vector were incubated with Con and 10 μ g/ml CIS. We found that CIS (Figures 6b–8b, curve 2) and inhibitors for miR-181a, miR-519a and miR-374a (Figures 6b, 7b and 8b, curve 4) dramatically decreased cell survival, whereas mimics for miR-181a (Figure 6b, curve 3), miR-519a (Figure 7b, curve 3), and miR-374a (Figure 8b, curve 3) minimally decreased the survival of cells compared with incubation of cells in the Con (Figures 6b, 7b and 8b, curve 1) or CIS (Figures 6b, 7b and 8b, curve 2). This finding suggests that the p- Δ Np63 α -dependent miRNAs are critically involved in the CIS-induced apoptotic process.

CIS induces apoptosis through deregulation of p- Δ Np63 α -dependent miRNAs targeting BCL2 and BCL2L2. Of the p- Δ Np63 α -dependent miRNAs upregulated




Figure 6 CIS-induced modulation of miR-181a expression leads to the activation of cell cycle arrest and apoptotic markers. Wild-type Δ Np63 α cells were transfected with control (mock), mimic, or inhibitor for miR-181a for 24 h and then exposed to Con (–) or 10 μ g/ml CIS (+) for an additional 24 h. (a) Protein levels of TP53, p-TP53, PARP2, CDKN1B, and DDIT4 were examined by immunoblotting with the indicated antibodies and loading levels were tested with an anti- β -actin antibody. (b) CIS-induced modulation of cell survival by miR-181a. Wild-type Δ Np63 α cells were transfected with control (mock, curves 1 and 2), miR-181a mimic (curve 3), or miR-181a inhibitor (curve 4) and then exposed to Con (curves 1, 3, and 4) or 10 μ g/ml CIS (curve 2) for 0–120 h. Cell survival was assessed at 24, 48, 72, 96, and 120 h by MTT cell proliferation assay. Absorbance readings were taken using a SpectraMax M2e Microplate fluorescence reader (Molecular Devices) at 570 and 650 nm wavelengths. All samples were run in triplicate. Experiments were performed in triplicate with \pm S.D. as indicated (<0.05).

in HNSCC cells upon CIS exposure, miR-630 stood out. After inspection of the mRNA targets for miR-630 on <http://www.microrna.org> and monitoring of their miRNA-binding activities (Supplementary Figures S14, S15 and S16), we focused on critical apoptotic activators (YAP1) and apoptotic inhibitors (BCL2 and BCL2L2). Wild-type Δ Np63 α cells were




Figure 7 CIS-induced modulation of miR-519a expression leads to the activation of cell cycle arrest and apoptotic markers. Wild-type Δ Np63 α cells were transfected with control (mock), mimic, or inhibitor for miR-519a for 24 h and then exposed to Con (–) or 10 μ g/ml CIS (+) for an additional 24 h. (a) Protein levels of PARP1, CDKN1A, YES1, and TP73 were examined by immunoblotting with the indicated antibodies, and loading levels were tested with an anti- β -actin antibody. (b) CIS-induced modulation of cell survival by miR-519a. Wild-type Δ Np63 α cells were transfected with control (mock, curves 1 and 2), miR-519a mimic (curve 3), or miR-519a inhibitor (curve 4) and then exposed to Con (curves 1, 3, and 4) or 10 μ g/ml CIS (curve 2) for 0–120 h. Cell survival was assessed at 24, 48, 72, 96, and 120 h by MTT assay. Experiments were performed in triplicate with \pm S.D. as indicated (<0.05).

transfected with control (mock), mimic, or inhibitor for miR-630 for 24 h and exposed to Con (–) and 10 μ g/ml CIS (+) for an additional 24 h. We found that CIS and the miR-630 mimic decreased the expression of YAP1, BCL2, and BCL2L2, whereas the miR-630 inhibitor increased the protein levels of these apoptotic regulators (Figure 9a). For cell survival study, cells transfected with mimic or inhibitors for specific miRNAs were exposed to Con, while cells transfected with the mock vector were incubated with Con and 10 μ g/ml CIS. We then found that CIS (Figure 9b, curve 2) and miR-630 mimic (Figure 9b, curve 4) dramatically decreased cell survival, whereas the miR-630 inhibitor




Figure 8 CIS-induced modulation of miR-374a expression leads to the activation of apoptotic markers. Wild-type Δ Np63 α cells were transfected with control (mock), mimic, or inhibitor for miR-374a for 24 h and then exposed to Con (-) or 10 μ g/ml CIS (+) for an additional 24 h. (a) Protein levels of DDIT4, CASP3, and NF-YB were examined by immunoblotting with the indicated antibodies and loading levels were tested with an anti- β -actin antibody. (b) CIS-induced modulation of cell survival by miR-374a. Wild-type Δ Np63 α cells were transfected with control (mock, curves 1 and 2), miR-374a mimic (curve 3), or miR-374a inhibitor (curve 4) and then exposed to Con (curves 1, 3, and 4), or 10 μ g/ml CIS (curve 2) for 0–120 h. Cell survival was assessed at 24, 48, 72, 96, and 120 h by MTT assay. Experiments were performed in triplicate with \pm S.D. as indicated (<0.05)

(Figure 9b, curve 3) increased cell survival compared with the Con (Figure 9b, curve 1).

Discussion

Given that the miRNA expression is maintained by RNA polymerase II and III transcription machinery, it is likely that the potential regulatory role of p- Δ Np63 α is intimately intertwined with the role of other transcription factors, co-activators/co-repressors, histone acetyl-transferases/deacetylases, histone and DNA methyltransferases, and other chromatin-accessory proteins.^{27–32} Moreover, because miRNAs are processed from capped, polyadenylated transcripts,




Figure 9 CIS-induced activation of miR-630 expression leads to the modulation of apoptotic markers. Wild-type Δ Np63 α cells were transfected with control (mock), mimic, or inhibitor for miR-630 for 24 h and then exposed to Con (-) or 10 μ g/ml CIS (+) for an additional 24 h. (a) Protein levels of BCL2, BCL2L2, and YAP1 were examined by immunoblotting with the indicated antibodies, and loading levels were tested with an anti- β -actin antibody. (b) CIS-induced modulation of cell survival by miR-630. Wild-type Δ Np63 α cells were transfected with control (mock, curves 1 and 2), miR-630 inhibitor (curve 3),³⁷ or miR-630 mimic (curve 4) and then exposed to Con (curve 1, 3, and 4) or 10 μ g/ml CIS (curve 2) for 0–120 h. Cell survival was assessed at 24, 48, 72, 96, and 120 h by MTT assay. Experiments were performed in triplicate with \pm S.D. as indicated (<0.05)

a complex RNA-processing machinery^{26,33} may also be involved in the regulation of miRNA expression and may be potentially affected by CIS exposure. However, this study was exclusively focused on the CIS/p- Δ Np63 α functional relationship regarding the miRNA expression in HNSCC cells.

In the current study, we attempted to understand the role for p- Δ Np63 α in the regulation of the miRNA signature of HNSCC cells exposed to CIS treatment. (1) Which miRNAs are affected by CIS exposure and (2) which miRNAs affected by CIS exposure are potentially regulated by the p- Δ Np63 α transcription factor? We found that the exposure of wild-type Δ Np63 α cells to CIS induced the expression of DICER1, which is a critical component of the RNA-induced silencing micro-processor complex (RISC) implicated in miRNA maturation.¹⁵ Recently, Elsa Flores' research team elegantly showed that

p63 coordinately regulates DICER1 and miR-130b transcription, while suppressing the tumor cell metastatic potential.²⁴ We, however, showed that p- Δ Np63 α induced DICER1 transcription in wild-type Δ Np63 α cells upon CIS exposure. We found that p- Δ Np63 α deregulated a number of miRNAs in wild-type Δ Np63 α cells upon CIS exposure, ultimately changing the landscape of mRNA expression controlled by miRNAs. Among many miRNAs differentially regulated by p- Δ Np63 α , a few (181a, 519a, and 374a) showed the highest degree of inhibition, whereas miR-630 showed the highest degree of activation. We further found that although p- Δ Np63 α induces upregulation and downregulation of miRNA expression, the profound effect of DICER1 is seen in miRNA expression upregulated*** by CIS-mediated and p- Δ Np63 α -dependent mechanism. This observation supports the cumulative effect of the DICER1 on the p- Δ Np63 α -dependent upregulated miRNAs and antagonistic effect of DICER1 on the p- Δ Np63 α -dependent downregulated miRNAs. We then showed that p- Δ Np63 α directly affects miRNA transcription by binding to specific miRNA promoters and regulating miRNA expression levels accordingly. Finally, we showed that the downstream targets affected by the p- Δ Np63 α -dependent modulation of miRNA expression in wild-type Δ Np63 α cells include critical regulators of TP53-dependent and TP53-independent apoptotic genes.^{2,34–36}

Our data strongly suggest that following CIS exposure, the p- Δ Np63 α transcription factor is likely to play a decisive role in the regulation of certain miRNAs leading to the activation of pro-apoptotic pathways in HNSCC cells. There are two potential mechanisms by which p- Δ Np63 α can regulate miRNA expression: through the DICER1 transcriptional upregulation and subsequent maturation of miRNAs, and/or through direct transcriptional regulation of miRNA gene promoters via formation of protein complexes with other transcriptional and chromatin-associated factors. On one hand, the expression of mRNA target genes is maintained through a coupling mechanism that includes transcription factors and miRNA-mediated post-transcriptional machinery.^{14,32,37} On the other hand, the miRNA expression levels in cells under various experimental conditions are controlled by dual regulation through transcription factors and by post-transcriptional regulation by RISC components, as suggested by the data from this study. It is likely that both mechanisms play specific roles in the CIS-induced microRNAome deregulation in HNSCC cells. Certain miRNAs whose expression is altered after exposure of wild-type Δ Np63 α cells to CIS can induce pro-apoptotic signaling pathways (e.g. ATM-HIPK2-TP53-HMGA1, SIRT1-PARP, caspase cascade, TP73-YES1-YAP1, DDIT1 and 4, and BCL2, etc.), while activating cell cycle regulators (CDKN1A, CDKN1B, CDKN2C, and CDKN3).

Numerous reports have shown that TP53 protein accumulates rapidly upon DNA damage through a post-translational phosphorylation mechanism, leading to its activation as a transcriptional factor and subsequently to programmed cell death.^{5–12} CIS has been shown to increase the expression and activity of HIPK2, whose activation subsequently leads to CIS-dependent apoptosis through TP53 phosphorylation at Serine-46 (S46-p-TP53), which in turn activates certain apoptotic gene promoters and modulates the TP53-dependent

protein–protein interactions.^{34–36} Forced HIPK2 expression in U2OS cells increased the apoptotic response to CIS and potentiated the tumor suppressive activity of TP53 and TP73 through their protein–protein interactions.^{34–36} Moreover, HMGA1 overexpression promoted HIPK2 import from the nucleus to the cytoplasm with the subsequent inhibition of TP53 apoptotic function.³⁶ A recent report by other investigators also showed that specific miRNAs regulate TP53, PARP, and CDKN1 upon DNA damage through deregulation of miR-181a and miR-630 in non-small cell lung cancer A549 cells upon CIS exposure.³⁸

We found here that mRNAs for many proteins involved in cell cycle arrest and cell death are direct targets for miRNA whose expression is altered by the CIS/p- Δ Np63 α -dependent regulation in HNSCC cells (Supplementary Figures S8–S17). We showed that p- Δ Np63 α is implicated in the tight regulation of the miRNA landscape, which in turn modulates downstream mRNA targets that contribute to critical cellular decisions about cell survival or cell death. Our study establishes a new functional link between p- Δ Np63 α and the deregulated miRNAome during CIS-induced tumor cell death, suggesting novel mechanisms (transcriptional regulation and DICER1-dependent pri-miRNA maturation) through which p- Δ Np63 α could potentially act as a regulator of death or survival for HNSCC cells during chemotherapy.

Many fundamental questions remain regarding the transcriptional function of p53 family members in general and particularly in p- Δ Np63 α . How does p63 activate or repress gene transcription, specifically the miRNA gene transcription? What molecular regulators (e.g. other transcription factors, co-activators/co-repressors, histone acetyltransferases/deacetylases, histone and DNA methyltransferases) determine the binding of p- Δ Np63 α to specific gene promoters and its transactivational activity towards the RNA polymerase II initiation complex and define the outcome leading to gene activation *versus* gene repression? Although p- Δ Np63 α is supposed to control miRNA expression through the CCAAT box in both positive (upregulated miRNAs) and negative (downregulated miRNAs) manners, we can only speculate why p- Δ Np63 α activated some miRNA promoters and repressed others. Potential interactions of p- Δ Np63 α with other transcription factors (e.g. TP53, NF- κ B, E2F, STAT, C/EBP β , Oct-1, and Sox-5) and chromatin modifiers (e.g. p300, PCAF, HDAC, and DNMT) might provide the necessary support (Supplementary Figures S1–S6).^{27–32} Recent report showed that HDAC1/2 are present at the promoter regions of Δ Np63-repressed targets in keratinocytes, indicating a direct requirement for HDAC1/2 in Δ Np63-mediated repression.³⁹

Materials and Methods

Cells and reagents. The HNSCC cell line JHU-029 (expressing wild-type p53 and p63) was isolated from primary tissue at the Department of Otolaryngology/Head and Neck Surgery of the Johns Hopkins University School of Medicine.^{21,22} The Flp-In approach used to generate the JHU-029 clones allowed us to introduce the S385G mutation into the genomic DNA.^{21,22} Accordingly, these cells exclusively expressed the Δ Np63 α -S385G protein, which was confirmed by sequencing the individual mRNA-amplified clones obtained from wild-type and mutated cells. These stable clones were designated as wild-type Δ Np63 α cells and Δ Np63 α -S385G cells. Cells were maintained in RPMI medium 1640 and 10% fetal bovine serum and incubated with Con or 10 μ g/ml CIS (Sigma, St. Louis, MO, USA) for the indicated time periods. Cells were lysed with buffer A (50 mM Tris, pH 7.5, 100 mM NaCl,

2 mM EDTA, 0.5% Triton X-100, 0.5% Brij-50, 1 mM PMSF, 0.5 mM NaF, 0.1 mM Na_3VO_4 , $2 \times$ complete protease inhibitor cocktail), sonicated five times for 10-s intervals, and centrifuged for 30 min at $15\,000 \times g$. Supernatants were analyzed by immunoblotting.^{21,22} Immunoblots were scanned using a PhosphorImager and quantified using Image Quant software version 3.3 (Molecular Dynamics, Sunnyvale, CA, USA), and the levels of tested proteins were normalized against β -actin levels. Values were expressed as fold change compared with a control sample (defined as 1).

Antibodies. We used a mouse anti-YAP1 polyclonal antibody (ab22144), a mouse anti-p73 α/β monoclonal antibody (clone ER-15, ab17230), a mouse anti-YES1 monoclonal antibody (ab61206), a mouse anti-BCL2 polyclonal antibody (clone 124, ab694), a rabbit monoclonal anti-CDKN1B antibody (Ab32034), a rat monoclonal anti-BCL2L2 antibody (clone 16H12, ab54313), and a rabbit polyclonal anti-NF-YB antibody (ab6559), which were all purchased from Abcam (Cambridge, MA, USA). A rabbit anti-DICER1 polyclonal antibody (NPB1-06521) was purchased from Novus Biologicals (Littleton, CO, USA), whereas a rabbit anti-CASP3 polyclonal antibody (H00000836-D01P) and a mouse anti-CDKN1A monoclonal (H00001026-MO2) were obtained from Abnova Corporation (Taipei, Taiwan). We used a rabbit anti-PARP1 monoclonal antibody (LS-C38096) and a rabbit anti-PARP2 polyclonal antibody (LS-C40697) obtained from Lifespan Biosciences (Seattle, WA, USA). A rabbit anti-human TP53-S46 (p-Ser46) polyclonal antibody (sc-101764) was purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). A rabbit anti-human polyclonal GADD45A (DDIT1) antibody (#3518) and a rabbit anti-human DDIT4 polyclonal antibody (#2516) were obtained from Cell Signaling Technology Inc. (Danvers, MA, USA). We also used a monoclonal antibody against human β -actin (Sigma, St. Louis, MO, USA), a monoclonal antibody anti-human wild-type TP53 antibody (clone DO-1, #554293, BD Pharmingen, San Diego, CA, USA), and a mouse monoclonal antibody against the FLAG epitope (Clone 29E4.G7, #200-301-B13, Rockland Immunochemicals, Inc., Rockville, MD, USA). A rabbit polyclonal antibody against Δ Np63 (anti-p40, PC373, residues 5–17 epitope) was purchased from EMD/Calbiochem (San Diego, CA, USA). A custom rabbit polyclonal antibody against a phosphorylated peptide encompassing the Δ Np63 α protein sequence (ATM motif, NKLPSTV-pS-QLINPQQ, residues 379–392) was prepared and purified against the phosphorylated peptide *versus* non-phosphorylated peptide.²¹

Reverse transcription-PCR. The First Strand cDNA Synthesis kit (Invitrogen, Carlsbad, CA, USA) was used for reverse transcription. PCR was performed with recombinant Taq polymerase (Invitrogen) as follows: 24–30 cycles consisting of 94°C for 1 min, 55°C for 1 min, and 72°C for 30 s, as described elsewhere.²² As an internal loading control, we amplified a region from GAPDH mRNA using the following primers: sense primer, 5'-CTACATGGTTTACATGT-3' (121) and antisense primer, 5'-TGCCTCAACGACCACT-3' (920). These primers give rise to an 800-bp PCR product. For the DICER1 amplification, we used the following primers: sense primer, 5'-GGTCTTAGACAGGTACTTCT-3' (961) and antisense primer, 5'-TGCCCTACTTGACCTTG-3' (1336). These primers produce a 375-bp PCR product. PCR products were resolved by 2% agarose electrophoresis. For each set of primers the number of amplification cycles was predetermined to achieve the exponential phase of the PCR reaction. PCR amplification results were analyzed digitally by Kodak 1D 3.5 software (Kodak Scientific Imaging, Eastman-Kodak, Rochester, NY, USA). The net intensity of PCR bands was measured and normalized against the net intensity of GAPDH bands.²² The same primers were used for qPCR amplification of DICER1 and GAPDH mRNA transcripts.

ChIP. First, 5×10^6 cell equivalents of chromatin (2–2.5 kb in size) were immunoprecipitated (IP) with 10 μg of anti-p- Δ Np63 antibody or anti- Δ Np63 antibody, as essentially described elsewhere.^{22,23} After reversal of formaldehyde crosslinking, RNase A and proteinase K treatments, IP-enriched DNAs were used for PCR amplification. PCR consisted of 40 cycles of 94°C for 30 s, 60°C for 30 s, and 72°C for 30 s using Taq DNA polymerase (Invitrogen). The relative enrichment (Bound) measurements for the indicated regions of tested promoters were shown as percentage of Input (Supplementary Figure S7). PCR primers used for amplification of the DICER1 promoter and miRNA promoters are listed in Supplementary Figures S1–S6.

Luciferase reporter assay. We used the pGL3-DICER1 (S109038) promoter-luciferase reporter plasmid (encompassing –870 to +50 bp of the DICER1 promoter) and 3'-UTR luciferase reporter constructs for DICER1 (S214311),

DDIT1 (S204414), DDIT4 (S206522), and YES1 (S211786) obtained from SwitchGear Genomics (Menlo Park, CA, USA), whereas 3'-UTR luciferase reporter plasmids for HIPK1 (SC221730) and BCL2 (SC222289) were purchased from Origene Technologies (Rockville, MD, USA). A total of 5×10^4 cells/well in a 24-well plate were transfected with 100 ng of the pGL3 luciferase reporter constructs plus 1 ng of the Renilla luciferase plasmid pRL-SV40 (Promega, Madison, WI, USA) using FuGENE 6 (Roche-Diagnostics, Indianapolis, IN, USA), as previously described.^{22,23} Cells were also transfected with 100 ng of the mimics or inhibitors of tested miRNAs. At 24 h, cells were also treated with 10 $\mu\text{g}/\text{ml}$ CIS or Con and then, after an additional 24 h, luciferase assays were performed using the Dual luciferase reporter assay kit (Promega) for the promoter plasmid, and the Luc-Pair miR Luciferase Assay kit (Gene Copoeia, Inc., Rockville, MD, USA) for the 3'-UTR plasmids. For each experiment, the wells were transfected in triplicate and each well was assayed in triplicate. To 50 μl of Firefly luciferase substrate was added 10 μl of cleared cell lysate, and light units were measured in a luminometer. Renilla luciferase activity was measured in the same tube after addition of 50 μl of Stop and Glo reagent. The values for Firefly luciferase activity were normalized against the Renilla luciferase activity values for each transfected well.

miRNA microarray analysis. miRNA microarray chips for 874 human miRNAs were synthesized by Combimatrix, and miRNA array analysis was performed as described elsewhere.¹³ Probes containing two mismatches were included for all miRNAs. Arrays were prehybridized at 37°C for 1 h in $3 \times$ SSC, 0.1% SDS, and 0.2% bovine serum albumin. Ten micrograms of total RNA was isolated using Trizol (Invitrogen), labeled with Cy3 and hybridized to arrays at 37°C overnight in 400 mM Na_2HPO_4 (pH 7.0), 0.8% BSA, 5% SDS, and 12% formamide. Arrays were washed once at room temperature in $2 \times$ SSC, 0.25% SDS, three times at room temperature in $1.6 \times$ SSC, and twice in ice-cold $0.8 \times$ SSC. Hybridized arrays were then scanned using a GenePix 4000B microarray scanner (Axon, Inc., Darien, CT, USA), and signal intensities were extracted using the Combimatrix Microarray Imager software (Combimatrix Diagnostics, Irvine, CA, USA). The background value was determined by calculating the median signal from the mismatch probes, and this value was subtracted from all perfect match probes. Signals that were less than 1.5 times stronger than the background were removed, and data sets were median centered prior to calculation of fold-change values. No significant differences in miRNA levels between wild-type Δ Np63 α cells and Δ Np63 α -S385G cells treated with Con were found, with these levels being significantly lower than in cells exposed to CIS.

Validation of miRNA array data by qPCR. To validate the differential expression of miRNAs, we isolated total small RNAs using a miRvana miRNA isolation kit (#AM1560, Applied Biosystems/Life Technologies, Carlsbad, CA, USA). We then used the High Capacity cDNA Reverse Transcription kit (#4374966, Applied Biosystems) to produce single-stranded cDNA probes. Next, we performed a qPCR using the TaqMan MicroRNA Assay Kit TaqMan U47 (#4380911) and TaqMan Gene Expression Master Mix, 1-Pack (#4369016), both obtained from Applied Biosystems.¹⁹ For precursor pri-miRNAs, we used the following individual kits: pri-hsa-miR-181a (Hs03302966_pri), pri-hsa-miR-517a (Hs03302632_pri), pri-hsa-miR-374a (Hs03304235_pri), pri-hsa-miR-630 (Hs03304713_pri), and pri-hsa-miR-885-3p (Hs03305150_pri). For mature miRNAs, we also used the following: hsa-miR-181a (#000480), hsa-miR-519a (#002415), hsa-miR-374a (#000563), hsa-miR-630 (#001563), hsa-miR-194 (#000493), and hsa-miR-885-3p (#002372). The reaction conditions were 50°C for 2 min, 95°C for 10 min, 40 cycles of 95°C for 15 s and 60°C for 1 min with a sample volume of 20 μl . The independent biological experiments were performed twice. Each RNA sample was amplified in triplicate, and obtained values used for statistical analysis. Expression was normalized to the U47 expression (gene ID 26802), and expression levels were determined as the average Ct of the U47 control. This averaged value was then used to normalize the sample's Ct. The average miRNA expression was determined using the Mann–Whitney *U*-test.¹⁹

miRNA mimics and inhibitors and transfection. Individual miRNA mimics (precursors) (hsa-miR-181a (PM10381), hsa-miR-519a (PM12922), hsa-miR-374a (PM12702), and hsa-miR-630 (PM11552)), and inhibitors: (hsa-miR-181a (AM10381), hsa-miR-519a (AM12922), hsa-miR-374a (AM12702), and hsa-miR-630 (AM11552)) were purchased from Ambion/Applied Biosystems (Austin, TX, USA). Cells were transfected for 24 h in a six-well plate with 100 pmol of the mimic, inhibitor or control in 500 μl serum-free medium with 5 μl Lipofectamine 2000 reagent (Invitrogen). Each experiment was performed independently at least three times and

in triplicate. To test the effect of DICER1 on miRNA expression in HNSCC cells after CIS exposure, we used scrambled siRNA (5'-AATTCTCCGAACGTGTCACGT-3') and DICER1 siRNA (5'-AAGGCTTACCTTCTCCAGCT-3') obtained from Dharmacon (Lafayette, CO, USA), as previously described.⁴⁰

To maximally inhibit DICER1 expression and keep the cells at optimal density, Δ Np63 α cells were transfected with two rounds of DICER1 siRNA or scrambled siRNA (each at 100 nM). Cells were split 48 h after the first round of DICER1 siRNAs transfection and were transfected again with the same siRNAs for another 24 h along with exposure to Con or 10 μ g/ml CIS. After the last treatment, cells were collected for PCR or immunoblotting with antibodies to DICER1 or β -actin.

Cell survival assay. Cells were plated at 20–30% confluency in each well of a six-well plate. Growth was assayed at 24, 48, 72, 96 and 120 h by MTT cell proliferation assay (American Tissue Culture Collection, Manassas, VA, USA). Absorbance readings were taken using a SpectraMax M2e Microplate fluorescence reader (Molecular Devices, Sunnyvale, CA, USA) at 570 and 650 nm wavelengths. All samples were run in triplicate.^{21,22}

FACS cell cycle analysis. Cells were harvested, washed in PBS, and fixed in MetOH:acetic acid solution (4 : 1) for 60 min at +4 °C. Cells were then incubated in 500 μ l of staining solution (50 μ g/ μ l of propidium iodide, 50 μ g/ μ l of RNase, 0.1% Triton X-100 in 1 \times PBS for 1 h at +4 °C and analyzed by flow cytometry performed on the FACS Calibur instrument (BD Biosciences, Sparks, MD, USA).

Statistical analysis. Statistical analysis was performed using the unpaired Student's *t*-test. *P*-values of <0.05 was considered to indicate statistical significance.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements. This work was supported in part by NIH grant 5R01DE13561 (DS, ER and BT).

- Kelland L. The resurgence of platinum-based cancer chemotherapy. *Nat Rev Cancer* 2007; **7**: 573–584.
- Zangen R, Ratovitski E, Sidransky D. Np63 α levels correlate with clinical tumor response to cisplatin. *Cell Cycle* 2005; **4**: 1313–1315.
- Sancar A, Lindsey-Boltz L, Unsal-Kacmaz K, Linn S. Molecular mechanisms of mammalian DNA repair and the DNA damage checkpoints. *Annu Rev Biochem* 2004; **73**: 39–85.
- Banin S, Moyal L, Shieh S, Taya Y, Anderson CW, Chessa L *et al*. Enhanced phosphorylation of p53 by ATM in response to DNA damage. *Science* 1998; **281**: 1677–1679.
- Flores E, Tsai K, Crowley D, Sengupta S, Yang A, McKeon F *et al*. p63 and p73 are required for p53-dependent apoptosis in response to DNA damage. *Nature* 2002; **416**: 560–564.
- Rocco J, Leong C, Kuperwasser N, DeYoung M, Ellisen L. p63 mediates survival in squamous cell carcinoma by suppression of p73-dependent apoptosis. *Cancer Cell* 2006; **9**: 45–56.
- Oren M. Decision making by p53: life, death and cancer. *Cell Death Differ* 2003; **10**: 431–442.
- Walensky L. BCL-2 in the crosshairs: tipping the balance of life and death. *Cell Death Differ* 2006; **13**: 1339–1350.
- Kumar S. Caspase function in programmed cell death. *Cell Death Differ* 2007; **14**: 32–43.
- Toh W, Nam S, Sabapathy K. An essential role for p73 in regulating mitotic cell death. *Cell Death Differ* 2010; **17**: 787–800.
- Levy D, Adamovich Y, Reuven N, Shaul Y. The Yes-associated protein 1 stabilizes p73 by preventing Itch-mediated ubiquitination of p73. *Cell Death Differ* 2007; **14**: 743–751.
- Yang C, Jausdal V, Haun R, Seth R, Shah S, Kaushal G. Transcriptional activation of caspase-6 and -7 genes by cisplatin-induced p53 and its functional significance in cisplatin nephrotoxicity. *Cell Death Differ* 2008; **15**: 530–544.
- Lee M, Kim J, Han K, Yeom S, Lee S, Baek SH *et al*. MicroRNA genes are transcribed by RNA polymerase II. *EMBO J* 2004; **23**: 4051–4060.
- Chen K, Rajewsky N. The evolution of gene regulation by transcription factors and micro-RNAs. *Nat Rev Genet* 2007; **8**: 93–103.
- Pratt A, MacRae I. The RNA-induced silencing complex: a versatile gene-silencing machine. *J Biol Chem* 2009; **284**: 17897–17901.
- Volinia S, Calin G, Liu C, Ambs S, Cimmino A, Petrocca F *et al*. A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci USA* 2006; **103**: 2257–2261.
- Chang S, Jiang W, Smith I, Poeta L, Begum S, Glazer C *et al*. MicroRNA alterations in head and neck squamous cell carcinoma. *Int J Cancer* 2008; **123**: 2791–2797.
- Melino G, Knight R. MicroRNAs meet cell death. *Cell Death Differ* 2010; **17**: 189–190.
- Chang T, Wentzel E, Kent O, Ramachandran K, Mullendore M, Lee KH *et al*. Transactivation of miR-34a by p53 broadly influences gene expression and promotes apoptosis. *Mol Cell* 2007; **26**: 745–752.
- Lena A, Shalom-Feuerstein R, Rivetti di Val Cervo P, Aberdam D, Knight RA, Melino G *et al*. miR-203 represses 'stemness' by repressing DeltaNp63. *Cell Death Differ* 2008; **15**: 1187–1195.
- Huang Y, Sen T, Nagpal J, Kim M, Upadhyay S, Trink B *et al*. ATM kinase is a master switch for the Δ Np63 α protein phosphorylation/degradation in human head and neck squamous cell carcinoma cells upon DNA damage. *Cell Cycle* 2008; **7**: 2846–2855.
- Huang Y, Chuang A, Romano R, Wu G, Sinha S, Trink B *et al*. Phospho- Δ Np63 α -NF- κ B protein complex transcriptionally regulates DDIT3 expression in squamous cell carcinoma cells upon cisplatin exposure. *Cell Cycle* 2010; **9**: 332–342.
- Osada M, Yamashita K, Park H, Fomenkov A, Kim M, Wu G *et al*. Differential recognition of response elements determines target gene specificity for p53 and p63. *Mol Cell Biol* 2005; **25**: 6077–6089.
- Su X, Chakravarti D, Cho M, Liu L, Gi Y, Lin YL *et al*. TP63 suppresses metastasis through coordinate regulation of Dicer and miRNAs. *Nature* 2010; **467**: 986–990.
- Liu C, Calin G, Volinia S, Croce C. MicroRNA expression profiling using microarrays. *Nat Protoc* 2008; **3**: 563–578.
- Lai E. Micro RNAs are complementary to 3'-UTR sequence motifs that mediate negative post-transcriptional regulation. *Nat Genet* 2002; **30**: 363–364.
- Reynold N, Schwartz B, Delvecchio M, Sadoul K, Meyers D, Mukherjee C *et al*. Oncogenesis by sequestration of CBP/p300 in transcriptionally inactive hyperacetylated chromatin domains. *EMBO J* 2010; **29**: 2943–2952.
- Strano S, Monti O, Pediconi N, Baccarini A, Fontemaggi G, Lapi E *et al*. The transcriptional coactivator Yes-associated protein drives p73 gene-target specificity in response to DNA Damage. *Mol Cell* 2005; **18**: 447–459.
- Perissi V, Aggarwal A, Glass C, Rose D, Rosenfeld M. A corepressor/coactivator exchange complex required for transcriptional activation by nuclear receptors and other regulated transcription factors. *Cell* 2004; **116**: 511–526.
- Pediconi N, Guerrieri F, Vossio S, Bruno T, Belloni L, Schinzari V *et al*. SIRT1-dependent regulation of the PCAF-E2F1-p73 apoptotic pathway in response to DNA damage. *Mol Cell Biol* 2009; **29**: 1989–1998.
- Cho H, Orphanides G, Sun X, Yang X, Ogryzko V, Lees E *et al*. A human RNA polymerase II complex containing factors that modify chromatin structure. *Genes Dev* 1998; **12**: 2153–2163.
- Kaesser M, Iggo R. Promoter-specific p53-dependent histone acetylation following DNA damage. *Oncogene* 2004; **23**: 4007–4013.
- Cai X, Hagedorn C, Cullen B. Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA* 2004; **10**: 1957–1966.
- Leong C, Vidnovic N, DeYoung M, Sgroi D, Ellisen L. The p63/p73 network mediates chemosensitivity to cisplatin in a biologically defined subset of primary breast cancers. *J Clin Invest* 2007; **117**: 1370–1380.
- D'Orazi G, Cecchinelli B, Bruno T, Manni I, Higashimoto Y, Saito S *et al*. Homeodomain-interacting protein kinase-2 phosphorylates p53 at Ser 46 and mediates apoptosis. *Nat Cell Biol* 2002; **4**: 11–19.
- Pierantoni G, Rinaldo C, Mottolise M, Di Benedetto A, Esposito F, Soddu S *et al*. High-mobility group A1 inhibits p53 by cytoplasmic relocalization of its proapoptotic activator HIPK2. *J Clin Invest* 2007; **117**: 693–702.
- Shalgi R, Brosh R, Oren M, Pilpel Y, Rotter V. Coupling transcriptional and post-transcriptional miRNA regulation in the control of cell fate. *Aging* 2009; **1**: 762–770.
- Galluzzi L, Morselli E, Vitale I, Kepp O, Senovilla L, Criollo A *et al*. miR-181a and miR-630 regulate cisplatin-induced cancer cell death. *Cancer Res* 2010; **70**: 1793–1803.
- Leboeuf M, Terrell A, Trivedi S, Sinha S, Epstein J, Olson EN *et al*. Hdac1 and Hdac2 act redundantly to control p63 and p53 functions in epidermal progenitor cells. *Dev Cell* 2010; **19**: 807–818.
- Tang K, Ren H, Cao J, Zeng G, Xie J, Chen M *et al*. Decreased Dicer expression elicits DNA damage and up-regulation of MICA and MICB. *J Cell Biol* 2008; **182**: 233–239.

Supplementary Information accompanies the paper on Cell Death and Differentiation website (<http://www.nature.com/cdd>)