

Characterization of a caspase-3-substrate kinome using an N- and C-terminally tagged protein kinase library produced by a cell-free system

D Tadokoro¹, S Takahama², K Shimizu¹, S Hayashi¹, Y Endo^{*,1,2,3} and T Sawasaki^{*,1,2,3}

Caspase-3 (CASP3) cleaves many proteins including protein kinases (PKs). Understanding the relationship(s) between CASP3 and its PK substrates is necessary to delineate the apoptosis signaling cascades that are controlled by CASP3 activity. We report herein the characterization of a CASP3-substrate kinome using a simple cell-free system to synthesize a library that contained 304 PKs tagged at their N- and C-termini (N/C-tagged PKs) and a luminescence assay to report CASP3 cleavage events. Forty-three PKs, including 30 newly identified PKs, were found to be CASP3 substrates, and 28 cleavage sites in 23 PKs were determined. Interestingly, 16 out of the 23 PKs have cleavage sites within 60 residues of their N- or C-termini. Furthermore, 29 of the PKs were cleaved in apoptotic cells, including five that were cleaved near their termini *in vitro*. In total, approximately 14% of the PKs tested were CASP3 substrates, suggesting that CASP3 cleavage of PKs may be a signature event in apoptotic-signaling cascades. This proteolytic assay method would identify other protease substrates.

Cell Death and Disease (2010) 1, e89; doi:10.1038/cddis.2010.65; published online 28 October 2010

Subject Category: Immunity

On the basis of the corresponding genetic sequences, > 500 human and mouse proteolytic enzymes have been predicted.¹ This number is comparable with that found for protein kinases (PKs), which are the main signal-transduction enzymes.^{2,3} Proteases are involved in the maturation, localization, stabilization, and complex formation of proteins, and in many biological processes, for example, normal development,^{4,5} cancer,^{6,7} infectious diseases,⁸ and cell death.⁹ Therefore, it is important to be able to identify protease substrates using simple assays.

Apoptosis requires the action of many different proteins that participate in apoptotic cell-signaling pathways.¹⁰ Caspases and PKs are critical components of growth and apoptosis signaling pathways.^{2,10} Large-scale analyses of the biological networks involving PKs and caspases are vital for the elucidation of apoptosis signaling pathways. Recent whole-cell proteomic studies that used mass spectrometry attempted to identify substrates of caspases that are involved in apoptosis and have shown that the percentage of PKs found as caspase substrates during apoptosis is 3–6% of ~300.^{11,12} However, cellular protein expression levels may have biased the results.¹³ Furthermore, it is difficult to identify specific pairs of proteases and substrates because numerous cleavage events occur simultaneously in cells. Therefore, an *in vitro* approach that could identify specific proteases and their corresponding substrates would complement cell-based approaches. A diagram, derived from a comprehensive *in vitro* study, that illustrates the relationships between

caspases and their PK substrates would help clarify the signal-transduction events that occur during apoptosis.

A collection of recombinant proteins, that is, a protein library, is needed to screen a large number of protein substrates. In addition, to screen a protein library comprehensively two *in vitro* high-throughput methods – one for protein synthesis and one for the detection of the targeted biochemical reaction – are required. Recently, we developed an automated protein synthesis system that uses a wheat cell-free system.^{14–16} Using this system, we were able to synthesize many human and Arabidopsis PKs.^{17,18} Recent work by others suggested that the wheat cell-free system could produce 13364 human proteins, which, because of the large number of proteins involved, represents an *in vitro*-expressed proteome.¹⁹ We also recently developed a method to label monobiotin proteins that had been synthesized in the wheat cell-free system.²⁰ These monobiotin-labeled proteins were then used directly – without purification – to detect protein ubiquitination²¹ and an autoantibody in the serum.²² As the procedures used with many commercially available detection kits depend on biotin–streptavidin interactions, our purification-free, synthesis/biotin-labeling method provides a simple and highly specific system that can be used for biochemical analyses.

Caspase-3 (CASP3) cleaves many different proteins,^{23,24} and its action *in vivo* irreversibly induces apoptosis. For the study reported herein, we delineated a CASP3-substrate kinome using a simple luminescent-based detection method

¹Cell-Free Science and Technology Research Center, Ehime University, Matsuyama, Ehime, Japan; ²The Venture Business Laboratory, Ehime University, Matsuyama, Ehime, Japan and ³RIKEN Genomic Sciences Center, Tsurumi, Yokohama, Japan

*Corresponding authors: T Sawasaki or Y Endo, Cell-Free Science and Technology Research Center, Ehime University, 3 Bunkyo-cho, Matsuyama, Ehime 790-8577, Japan. Tel: +81 89 927 8530; Fax: +81 89 927 9941; E-mail: sawasaki@eng.ehime-u.ac.jp or Tel: +81 89 927 9936; Fax: +81 89 927 9941; E-mail: yendo@eng.ehime-u.ac.jp

Keywords: caspase; protein kinases; apoptosis; cell-free protein synthesis; protein library

Abbreviations: CASP3, caspase 3; PK, protein kinase; N/C-tagged, N- and C-terminally tagged; TD, terminal detection

Received 10.6.10; revised 31.8.10; accepted 15.9.10; Edited by A Finazzi-Agró

to screen an N- and C-terminally tagged (Ntagged) PK library produced in the wheat cell-free system. This comprehensive characterization of a CASP3-substrate kinome is a resource that can be used to understand the roles of PKs in apoptosis.

Results

Generation of an Ntagged PK library used to identify CASP3 PK substrates. To identify PKs that are substrates of CASP3, we first made a library consisting of 248 human and 56 mouse PKs (Supplementary Table S1). The nucleotide sequences for the Flag-tag and the biotin ligation site (bls) were added upstream and downstream, respectively, of the PK open-reading frame by PCR incorporation of Gateway recombination tags. Each PCR product (attB1-Flag-PK-blb-attB2) was inserted into a pDONR221 vector using the Gateway BP Clonase II system (upper panel, Figure 1). The Flag-PK-blb nucleotide sequences from the *Escherichia coli* cultures were used without purification to construct, by split-primer PCR, the DNA templates for protein synthesis.¹⁴ The Ntagged PK library (304 PKs) was produced using an automated protein synthesizer (GenDecoder 1000; CellFree Sciences Co., Ltd., Matsuyama, Japan), with biotin and biotin ligase added into the synthesis mixtures for monobiotin labeling at the bls.^{20,21} That the members of the protein library were Ntagged was confirmed by immunoblotting with anti-Flag antibodies and Alexa488-labeled streptavidin.

To assess the suitability of the designed PKs to act as CASP3 substrates, we used Ntagged p21-activated kinase 2 (PAK2), which is a known CASP3 substrate,²⁵ as the test case. The biotinylated Ntagged-PAK2 (Flag-PAK2-blb~biotin) was treated with CASP3 and cleavage of PAK2 was confirmed by immunoblotting with Alexa488-conjugated streptavidin (Figure 2a). In addition, the cleavage site (₃₁₉DELD↓S₃₂₃), determined by amino-acid sequencing, was found to be the same as that reported previously.²⁵ (The arrow indicates the hydrolytic bond.)

A luminescent assay to detect PK substrates of CASP3. A schematic of the assay used to monitor cleavage of the Ntagged PKs by CASP3 is shown in Figure 1. The PK construct is first incubated with CASP3. If the construct contains a sequence that can be cleaved by CASP3, cleavage occurs. Acceptor and donor beads are then added. The Flag-tag binds a protein A-conjugated acceptor bead via an anti-Flag antibody, and the biotin bound to the C-terminus of the PK construct binds a streptavidin-conjugated donor bead. If an acceptor bead is in close contact with the donor bead, as is the case when the construct is not a CASP3 substrate and both beads are therefore bound intramolecularly, the system luminesces. However, if CASP3 had cleaved the Ntagged PK, luminescence is suppressed because the beads are no longer in close contact. As a proof-of-concept experiment, cleavage of the test PK, Ntagged PAK2, was assessed




Figure 1 Schematics of the DNA template construction and the CASP3-substrate-screening assay. *Protein kinase (PK)* genes were obtained from the human MGC and mouse FANTOM libraries, and from a library of *PK* genes that we had cloned. The *PK* genes were PCR amplified with the Flag and the biotin ligation site (bls) tags added to the upstream and downstream ends, respectively. The modified genes were each inserted into a Gateway pDONR221 vector (pDONR-Flag-PK-blb) and DNA templates (SP6-E02-Flag-PK-blb) were constructed by split-primer PCR and then expressed in the wheat cell-free protein synthesis system that included biotin ligase and D-biotin to give Flag-PK-blb~biotin constructs. The Flag and biotin tags were bound to protein A-conjugated acceptor beads via an anti-Flag antibody and streptavidin-conjugated donor beads, respectively. An intact complex luminesced strongly, whereas after CASP3 cleavage and dissociation of the protein fragments, the luminescence was abolished or reduced




Figure 2 Screening of CASP3-cleaved PK substrates from the NCTagged PK library. (a) Immunoblot of NCTagged PAK2 that had been incubated in the presence (+) or absence (–) of CASP3. Alexa488-labeled streptavidin (STA(C)) was used for detection. (b) Detection of CASP3-cleaved NCTagged PAK2 using the AlphaScreen system. The luminescence for the control (no CASP3) was set to 100%. The value of 19% indicated that most of the NCTagged PAK2 was cleaved by CASP3. Each value is the mean of three independent experiments, and the uncertainty is reported as the standard deviation. (c) Luminescent signals remaining after *in vitro* CASP3 treatment of NCTagged PKs that had been synthesized in the wheat cell-free system. The x axis lists the NCTagged PKs in ascending order of their luminescent signals after CASP3 treatment. (d) NCTagged PKs that returned luminescent signals of < 78% of the control values. The plot contains the data of (c) within the green rectangle. Red bars are for PKs that were known to be substrates of CASP3 before this report

using this system. CASP3 treatment decreased the luminescent signal to $19 \pm 7\%$ that of the control (no CASP3; Figure 2b). Therefore, the system could detect CASP3 cleavage and can replace conventional immunoblotting procedures.

Screening of the CASP3-substrate kinome. Using the luminescent system, 304 NCTagged PKs were screened. The level of luminescence after CASP3 treatment is reported as the percentage of the corresponding control (no CASP3; Figure 2c and d). Thirteen of the NCTagged PKs for which luminescence was low after CASP3 treatment are known CASP3 substrates.^{23,24,26} The smallest and largest luminescent values were for STK4 (1%) and BMX (73%), respectively; we therefore examined the physical states of the PKs that had been treated with CASP3 and had associated luminescence values of ~80% by immunoblotting with anti-Flag antibodies and Alexa488-streptavidin to detect the N- and C-termini of the NCTagged PKs, respectively. This ‘terminal detection’ (TD) immunoblot assay identified 43 NCTagged PKs that had been cleaved (Supplementary Table S1). In addition to the 13 PKs that were known to be CASP3 substrates, 30 previously unidentified PK that were substrates of CASP3 were found (Figure 3 and Table 1). In addition, because the apparent molecular weights of the N- and C-terminal fragments could be estimated from their positions in the TD immunoblot, the

CASP3 cleavage sites could be predicted (red arrowheads, Figure 3). For MASTL, the signal on the immunoblot with Alexa488-conjugated streptavidin was not detectable, probably indicating that the efficiency of biotinylation in MASTL proteins might be too low to detect for the immunoblot. Luminescent signal of this clone was also very low (see Supplementary Table S1).

A comparison of the luminescent and immunoblot data correlated a luminescent signal of < 78% with a positive immunoblot result. Forty-eight PK constructs with luminescent signals > 78% were tested and returned negative immunoblot results (Supplementary Table S1). Therefore, a luminescent signal of ~78% is the apparent divisor between PKs that can be cleaved by CASP3 and those that cannot be cleaved.

***In vivo* identification of the PKs that were identified as CASP3 substrates by the luminescent assay.** We investigated whether the newly identified PKs that were substrates of CASP3 were cleaved in HeLa cells that had been induced to undergo apoptosis by TNF α plus cycloheximide (TNF α)²⁷ or anti-Fas antibody (anti-Fas).²⁸ The genes encoding these PKs were each inserted into the transfection vector, pDEST26, using the Gateway system and expressed as (His)₆-PK-Flag constructs. We were able to detect all expressed PK constructs, except DYRK3, by immunoblotting with anti-(His)₆ or anti-Flag antibodies.



Figure 3 *In vitro* cleavage of Ntagged PKs by CASP3. The Ntagged PKs that had been incubated in the presence (+) or absence (–) of CASP3 and their cleavage products were detected using anti-Flag antibodies (FLAG(N)) and Alexa488-conjugated streptavidin (STA(C)), which bound to the N- and C-termini of the PK constructs, respectively. The cartoons of the proteins that are under the lanes show the locations of the conserved domains (colored boxes) and the predicted cleavage sites (red arrowheads). The conserved domains that are found in the Conserved Domains Database (<http://www.ncbi.nlm.nih.gov/cdd>) are: ACD, alpha-crystallin domain; activin, conserved domain for activin members; ADF, actin depolymerization factor/cofilin-like domain; α -kinase, conserved kinase domain for the α -kinase family; C1, phorbol esters/diacylglycerol binding domain; DCX, doublecortin domain; death, death domain; GS, GS motif; KA1, kinase-associated domain; kinase, catalytic domain of protein kinase; PB1, Phox and Bem1p domain; PH, pleckstrin homology domain; RIO, catalytic domain of eukaryotic RIO kinase family; SH2, src homology 2 domain; SH3, src homology 3 domain; UBL, ubiquitin-like domain

Notably, they were detected as cleavage products and/or were found in smaller amounts when the cells had been induced to undergo apoptosis than when apoptosis had been inhibited by z-VAD-FMK (Figure 4a and b). Furthermore, apoptosis-induced cleavage of four endogenous PKs was found by immunoblotting with commercially available antibodies against the endogenous PKs (Figure 4c). These *in vivo* experiments validated the underlying concept of our *in vitro* cell-free system as the *in vivo* system found all of the PKs identified by the *in vitro* system.

Characterization of the CASP3 cleavage sites in the newly identified PK substrates. We characterized the CASP3 cleavage sites in the newly identified PK substrates. As the positions of the cleaved PK fragments in the TD immunoblot could be used to estimate the size of the cleaved fragments and because the antibodies could be

used to identify whether the fragments were derived from the N- or C-terminal regions of the PKs, we could predict the approximate positions of the CASP3 cleavage sites (red arrowheads, Figure 3). Each Ntagged PK that was a substrate for CASP3 was synthesized in the cell-free system and purified using Streptavidin Magnesphere Paramagnetic beads. Their C-terminal fragments that bound to the beads were recovered after CASP3 cleavage and their sequences were determined. Using this approach, the cleavage sites of ACVR1, AKT2, BMPR1B, CaMKK1, HSPB8, MAPK12, MKNK2(D58), PDPK1, PRKCI, PRKX, RIOK1, RIOK3, and RPS6KA5 were determined. We then attempted to determine the cleavage sites of the remaining PKs by other methods.

The Ntagged PKs that had low biotin-labeling efficiencies and were cleaved near their C-termini were genetically modified by the addition of a glutathione-S-transferase (GST) fragment at their C-termini to facilitate recovery with

Table 1 Characteristics of the newly identified CASP3 PK substrates

Symbols	Kinome names	Groups	Clone origin	AA ^a	Cleavage sequence	Cleavage sites	Methods ^b	Conservation ^c	Smallest frag. ^d	<i>In vivo</i> cleavages ^e
ACVR1	ALK2	TKL	Hs	509	IASD↓M	269	NT	Yes	C240	Yes
AKT2	AKT2	AGC	Mm	481	DAMD↓Y	121	NT	Yes	N121	Yes
BMPR1B	ALK6	TKL	Hs	502	CSTD↓G	50	NT	Yes	N50	Yes
					DFVD↓G	120	NT	Yes		
CaMKK1	CaMKK1	Other	Hs	520	EEAD↓G	32	NT	Yes	N32	Yes
CSK	CSK	TK	Mm	450	DAPD↓G	409	MS	Yes	C41	Yes
CSNK1G1	CK1g1	CK1	Mm	459	VHVD↓S	343	MU	Yes	C116	Yes
eEF2K	eEF2K	Atypical	Hs	725	EGVD↓G	14	MU	Yes	N14	Yes
					DHLD↓N	430	MU	Yes		
HSPB8	H11	Atypical	Hs	196	MAD↓G	3	NT	Yes	N3	Yes
MAP2K6	MAP2K6	STE	Mm	334	DFVD↓F	289	MU	Yes	C45	Yes
MAPK12	p38g	CMGC	Hs	367	SAVD↓G	46	NT	Yes	N46	Yes
MARK1	MARK1	CAMK	Hs	795	SATD↓E	52	MU	Yes	N52	Yes
MKNK2	MNK2	CAMK	Hs	414	DQPD↓H	32	MU	No	N32	Yes
					DIPD↓A	58	NT	Yes		
PDPK1	PDK1	AGC	Hs	556	SHPD↓A	552	NT	Yes	C4	Yes
PIM2	PIM2	CAMK	Hs	334	TDFD↓G	198	MU	Yes	C113	Yes
PRKAA1	AMPKa1	CAMK	Hs	550	TSLD↓S	520	MS	Yes	C30	Yes
PRKCI	aPKCi	PKC	Hs	596	TQRD↓S	6	NT	Yes	N6	Yes
PRKX	PRKX	AGC	Hs	358	ETPD↓G	25	NT	No	N25	Yes
RIOK1	RIOK1	Atypical	Mm	568	EKDD↓I	37	NT	Yes	N37	Yes
RIOK3	RIOK3	Atypical	Hs	516	DTRD↓D	139	NT	Yes	N139	Yes
RPS6KA5	MSK1	AGC	Hs	549	DGGD↓G	20	NT	Yes	N20	Yes
					DELD↓V	344	NT	Yes		
					TEMD↓P	356	NT	Yes		
SNARK	NuaK2	CAMK	Hs	628	VSED↓S	546	MU	Yes	C82	Yes
TRIB3	TRB3	CAMK	Hs	358	VVPD↓G	338	NT	Yes	C20	Yes
ULK4	ULK4	Other	Hs	580	SQID↓S	473	MU	Yes	C107	Yes
DCAMKL2	DCAMKL2	CAMK	Hs	695	—	—	—	—	—	Yes
DYRK3	DYRK3	CMGC	Hs	568	—	—	—	—	—	ND
IRAK2	IRAK2	TKL	Mm	622	—	—	—	—	—	Yes
MASTL	MASTL	AGC	Hs	879	—	—	—	—	—	Yes
PCTK1	PCTAIRE1	CMGC	Hs	496	—	—	—	—	—	Yes
PTK9	A6	Atypical	Hs	384	—	—	—	—	—	Yes
TGFBR1	TGFBR1	TKL	Hs	426	—	—	—	—	—	Yes

Abbreviations: Hs, human clone; Mm, mouse clone; MS, mass spectroscopy; MU, mutation; ND, not determined; NT, N-terminal sequencing. ^aLength of amino acids. ^bMethods for determination of cleavage site. ^cVery similar site conserving the Asp (D) of the hydrolytic bond was found between human and mouse PKs (Yes), whereas no similar sites was done (No). ^dThe smallest N (N) or C (C) fragment in the cleaved PKs. Number is the length of amino acids of the fragment. ^eData from Figure 4.

glutathione Sepharose 4B beads after CASP3 cleavage. CASP3 cleavage of the PK-GSTs produced the same size N-terminal fragments as those of the corresponding CASP3-cleaved NCTagged PKs, indicating that the GST tags did not alter the positions of the cleavage sites. In addition, the sequences of the cleaved c-src tyrosine kinase (CSK) and AMP-activated kinase-α1 (AMPKa1) fragments were determined using MALDI/TOF-MS. Other PK constructs that were synthesized in small amounts were subjected to D→A mutagenesis to determine their cleavage sites. In total, 28 cleavage sites in 23 PKs were identified (Table 1). Identical or similar cleavage sites were found in the corresponding human and mouse PKs, except for those of PRKX (Supplementary Table S2). (Sequence analysis showed that mouse PRKX does not have the N-terminal region that is found in human PRKX.) Therefore, the CASP3-substrate kinome may be highly conserved in mammals.

We also analyzed the common sequence attributes among the 28 cleavage sites and found that CASP3 prefers the sequence, DXXD↓G (Figure 5a). The consensus PK cleavage site for CASP3 in the MEROPS database is DXXD↓X. In the NCTagged PK library, 208 of the 304 PKs contain a DXXDX sequence. However, only 33 PKs were

cleaved by CASP3; therefore, to be cleaved by CASP3, the DXXDX sequence and a structural element – probably accessibility – are required.

Characterization of the newly identified PKs that were cleaved near their N- or C-termini. Interestingly, 16 out of the 23 PKs, for which cleavage sites were characterized, have cleavage sites within 60 residues of their N- or C-termini. We investigated whether these sites were also cleaved *in vivo* when apoptosis was induced by TNFα. For these experiments, CaMKK1, eEF2K, MNK2, AMPKa1, and TRIB3, which were cleaved *in vitro* at D32, D14, D32/D58, D520 (30 residues away from the C-terminus), and D338 (20 residues away from the C-terminus), respectively, were used (Figure 5b and Table 1). Their genes (wild type, WT) were each reconstructed with a V5 tag added at the end opposite the cleavage site. The genes for their D→A mutants (DA), and for the sequences of their longer CASP3-cleaved fragments (C3M), were also constructed and all were expressed in control and in apoptotic cells (Figure 5c). Cleavage of the WT PKs produced long fragments corresponding to C3M in apoptotic cells, whereas z-VAD-FMK blocked cleavage. These cleavages near the N- and




Figure 4 *In vivo* caspase cleavage of the newly identified PK substrates and of endogenous PK substrates. **(a)** *In vivo* cleavage of the (His)₆-PK-Flag constructs expressed in apoptotic HeLa cells. The cells were treated with DMSO (control) or with TNF α and cycloheximide (TNF α) in the presence and absence of z-VAD-FMK (a CASP3 inhibitor) for 6 h and then lysed. The cell extracts were immunoblotted and the PK constructs were detected with anti-Flag antibodies, except for HSPB8 and MAP2K6. Anti-His tag antibody was used for the two PKs. **(b)** The cells were transfected with a plasmid of (His)₆-PK-Flag constructs, and treated with DMSO (control), or with anti-Fas antibody (anti-Fas) in the presence and absence of z-VAD-FMK for 6 h and then lysed. Immunoblotting was carried out as in **(a)**. **(c)** HeLa cells were treated as in **(a)**, but were not transfected with a (His)₆-PK-Flag gene. Each endogenous PK was detected using an antibody specific for it. α -Tubulin was used as an internal marker




Figure 5 The cleavage site logo and *in vivo* cleavage of five PKs that are cleaved by CASP3 near their N- or C-termini. **(a)** The 20 residues surrounding the D of the hydrolytic bond in 28 PKs were analyzed using WebLogo, version 3.0.³⁶ **(b)** Cartoons of five PK sequences that have cleavage sites near their N- or C-termini. The corresponding PKs were used for the experiment shown in **(c)**. The positions of the alanines in the D \rightarrow A mutants (DA) are shown, as are the long fragments (C3M) produced by CASP3 cleavage. V5 tags were fused at the ends farther away from the cleavage sites. The first M in C3M of CaMKK1, eEF2K, and MNK2 indicates a methionine as a start amino acid. **(c)** Immunoblots of PK-V5s and V5-PKs that had been expressed in apoptotic HeLa cells. The cells were treated with DMSO (control) or with TNF α and cycloheximide in the presence or absence of z-VAD-FMK for 6 h and then lysed. The proteins were blotted and then detected with anti-V5 antibodies. WT indicates a wild-type protein

C-termini of the PKs suggest that CASP3 cleavage may regulate the activity level and/or cellular localization of the PKs, rather than simply inactivate the kinases.

Discussion

In 1995, PITSLRE,²⁹ PKC δ ,³⁰ and DNA-PKcs³¹ were reported as the first PK-type substrates of CASP3. During the next 15 years, 36 additional PKs that can be cleaved by CASP3 were found.^{23,24,26} Notably, these authors showed that CASP3-cleaved PKs abrogate survival signals and accelerate apoptosis. In this study, we identified an additional 30 PKs that can be cleaved by CASP3. In addition, many of the cleavage sites were found in regulatory elements or in the regions near the N- and C-termini, rather than the kinase domain itself. Some of the newly identified CASP3-substrate PKs may be involved in apoptotic signal cascades. Sixteen PKs were shown to be cleaved *in vitro* near their N- or C-termini and at least five of them were also cleaved near their N- or C-termini in apoptotic cells (Figure 5). Using standard immunoblotting, proteins that are cleaved into a large and a small fragment may be overlooked because the mobilities of the large fragment and the intact protein will be nearly identical. Most of the PKs that had been previously reported to be cleaved by CASP3 were identified because the cleaved fragments had very different molecular weights than did the intact PK and were therefore easily detected by SDS-PAGE. Consequently, cleavages near the termini may have been overlooked. Taken together, our results suggest that CASP3 cleavage of some of the members of the CASP3-substrate kinome alters the function of the PKs and thereby signals apoptosis.

For the study reported herein, 304 out of 518 known PKs, synthesized as Ntagged PKs, were subjected to the *in vitro* cleavage assay (Supplementary Table S1). The relative number of PKs that were cleaved was ~14%. A total of 69 PKs that are CASP3 substrates are now known, which suggests that at least ~13% of the PKs in the human kinome are targets of CASP3. As ~200 PKs have yet to be tested as CASP3 substrates, an additional 26 PKs (13% of the 200) may be CASP3 substrates. The human genome contains 518 annotated PKs, which have been divided into 10 groups on the basis of their sequence homologies.³ Interestingly, the groups differ in terms of their susceptibilities to CASP3 cleavage (Table 2). Approximately 30% of the PKs in the AGC group are known CASP3 substrates, for example, AKT2, S6K, MSK, PKC, and PDK1. Many of the AGC-type PKs are commonly found in mammalian tissues,³² and their cleavage sites are located in their regulatory domains (Figure 4 and Table 1). Therefore, these abundant PKs may be activated when CASP3 cleaves them and then act as intracellular apoptosis signals. Conversely, CASP3 cleaved only a relatively small number (6~8%) of the PKs in the CMGC group, which includes the kinases of the CDK and CDKL families, and the tyrosine kinase groups. Therefore, most members of these groups may only act indirectly as apoptosis signals after CASP3 activation.

Such ROCK1 and MST1, certain caspase cleavage products, work as apoptosis signaling.^{23,24} In this study, we found at least six new CASP3 cleavage products, derived from AKT2, CaMKK1, eEF2K, MARK1, MNK2, and TRB3,

Table 2 Characteristics of the protein kinases used in this study

Groups	Total ^a	Tested clones	Cleaved clones (new ^b)	Cleaved/test clones (%)
AGC	63	33	10 (6)	30
CAMK	74	52	8 (7)	15
CK1	12	8	1 (1)	13
CMGC	61	39	3 (3)	8
Other	83	46	3 (2)	7
STE	47	25	4 (1)	16
TK	90	51	3 (1)	6
TKL	43	27	6 (4)	22
RGC	5	2	0	0
Atypicals	40	21	5 (5)	24
Total	518	304	43 (30)	14

^aEach number is corresponding to human kinome. ^bNewly PK numbers found in this study.

after 6 h from apoptosis induction (Figures 4 and 5). These cleavage products retain kinase domain, as in the case of ROCK1 and MST1. On the other hand, we could not detect any cleavage products from the other kinases *in vivo*. The reasons are not yet understood. However, recent proteomics approach has shown that the cleaved proteins displayed transient fragments in the apoptotic cells.¹² Further analysis at multiple time points during the apoptotic cascade would be required for detection of the cleavage products from the remaining PKs.

For TRB3, full-length TRB3 (D338A) mutant was decreased in apoptotic condition (compared TNF α lane with TNF α plus z-VAD-FMK lane in Figure 5c). However, the mutant could not produce the CASP3 cleavage product found as the shorter form in TNF α lane of WT, indicating that the mutant was not cleaved by CASP3. The mutant was also not cleaved by CASP3 *in vitro* (data not shown). As TRB3 has been known to receive proteasomal degradation,³³ this unexpected reduction of the mutant TRB3 in the apoptotic cells may be the effects of cycloheximide and/or caspase-inhibitor treatment on TRB3 degradation.

Proteases often modify the activities of their targeted protein substrates. Identification of the specific substrate that is cleaved by a protease is necessary if the functions of both the protease and its substrate are to be understood. Proteomic studies have used mass spectrometry to exhaustively identify cellular proteins that have been cleaved by proteases.^{11,12} However, it has been difficult to correlate specific proteases with their substrates because many proteases act at the same time *in vivo*.

Many full-length cDNAs derived from the genes of higher eukaryotes are available from many different sources. These cDNAs are potentially a great DNA template resource for *in vitro* syntheses of proteins. As a protein production system and for the functional analysis of proteins, the wheat cell-free system has many advantages: It can effectively use PCR-generated DNA templates.¹⁴ It is easily adapted to an automated system.¹⁵ It can be used to incorporate a single label into target proteins.²⁰ Its synthesized proteins do not require purification before being assayed, and it has no detectable proteasome activity.²¹ In addition, the screening cost is very low (~US\$1/assay), which for our study translated to 10 cents to produce each Ntagged protein and

20 cents for the beads, CASP3, and disposable hardware used in one assay.

In summary, we showed that an NTagged PK library synthesized in a cell-free system could be used to characterize a CASP3-substrate kinome. Analysis of the CASP3-cleavage sites indicated that CASP3 cleavage of PKs depends on both primary and tertiary structure. Almost all of the PK substrates that we identified *in vitro* were also identified *in vivo*. Systems similar to that used herein could be used to screen other protease substrates.

Materials and Methods

General. The following procedures have been described:^{14–16,20–22,34} wheat cell-free protein production; split-primer PCR synthesis of the DNA templates; parallel syntheses of mRNAs and their translated proteins; and measurements of the amounts of protein synthesized using densitometer scans of Coomassie brilliant blue-stained proteins or of radiolabeled proteins.

Construction of DNA templates for the expression of a PK protein library. The cloned genes encoding the PKs used in this study are listed in Supplementary Table S1. Their open-reading frames (without stop codons) were modified in two steps using PCR and the primers S1 (5'-CCACCCACCACACCAatg(n)₁₆-3') and T1 (5'-TCCAGCACTAGCTCCAGA(n)₁₉-3') (lowercase letters indicate nucleotides of the gene) for the first step, and the primers attB1-Flag-S1 (5'-GGGGACAAGTTTGTACAAAAAGCAGGCTTCATGGACTACAAGGATGACGATGACAAGCTCCACCCACCACCACCAATG-3') and T1-bis-STOP-attB2-anti (5'-GGGGACCACCTTTGTACAAGAAAGCTGGGTTTATTCGTGCCACTC GATCTTCTGGGCCGTCGAAGATGCTGTTCCAGGCCGTTCCAGCACTAGCTCCA GA-3') for the second step. The PCR-modified genes were each inserted into a pDONR221 vector using the Gateway BP Clonase II enzyme mix (Invitrogen, Carlsbad, CA, USA) to give pDONR-Flag-PK-bis vectors. *Escherichia coli* cells were transformed with these vectors and then cultured in wells of a 96-well plate that contained GYT medium (10% (v/v) glycerol, 0.125% (w/v) yeast extract, and 0.25% (w/v) tryptone) for 48 h without shaking. DNA templates for mRNA and protein expression were constructed using split-primer PCR¹⁴ in two steps. For the first step, the pDONR221-Flag-PK-bis plasmids that had not been isolated from the *E. coli* cells, and the primers pDONR221-1st_4080 (5'-ATCTTTTCTACGGGCTCTGA-3') and deSP6E02-Flag-S1 (5'-GGTGACACTATAGAAGCACTACCTATCTCTACACAAAACATTTCCCTACATACAACCTTCAACTTCTATTATGGACTACAA GGATGACGATGACAAGCTCCACCCACCACCACCAATG-3') were used, and for the second step, the amplified sequences of the first step and the primers SPu (5'-GGGTAGCATTTAGGTGACACT-3') and pDONR221-2nd_4035 (5'-ACGTAA GGGATTTTGGTCA-3') were used to give SP6-E02-Flag-PK-bis DNA templates. (The E02 sequence is a translational enhancer,³⁵ and the SP6 sequence is an SP6 RNA polymerase promoter.)

Cell-free protein synthesis. Cell-free protein synthesis used the reagents of an ENDEXT Wheat Germ Expression S Kit according to the manufacturer's instructions (CellFree Sciences Co., Ltd.), the bilayer translation method,^{15,16,34} and a robotic synthesizer (GenDecorder 1000; CellFree Sciences). Each DNA template was transcribed by SP6 RNA polymerase, then precipitated with ethanol, and collected by centrifugation (15 000 r.p.m. for 5 min., R10H rotor; Hitachi). Each mRNA (~30–35 µg) was washed with 75% ethanol, added into a translation mixture, and translated in the bilayer mode³¹ with the following modifications. The translation mixture (25 µl) (bottom layer) contained 60 A₂₆₀/ml of ENDEXT wheat germ extract, 1 × SUB-AMIX (24 mM Hepes-KOH, pH 7.8, 1.2 mM ATP, 0.25 mM GTP, 16 mM creatine phosphate, 2.5 mM DTT, 0.4 mM spermidine, 0.3 mM each of the 20 amino acids, 2.8 mM magnesium acetate, 100 mM potassium acetate), 2 µg creatine kinase (Roche Applied Science, Indianapolis, IN, USA), 500 nM D-biotin (Nacal Tesque, Kyoto, Japan), and 1 µl of the wheat cell-free translational mixture that expressed BirA biotin ligase (~50 ng/µl, BirA: GenBank Accession No. NP_0312927). A 1 × SUB-AMIX solution (125 µl) was placed over the translation mixture. The bilayer was incubated at 26°C for 17 h to allow for protein synthesis. All steps including construction of the DNA templates were performed in the wells of a 96-well plate.

Cleavage assay. The cell-free-synthesized PKs that had luminescent signals > 500 units (in the absence of CASP3) were studied. For each PK, 10 µl of the

CASP3 cleavage buffer (20 mM Tris-HCl, pH 7.5, 0.2 mM DTT, 5 mM MgCl₂, 3 mM ATP, 1 mg/ml BSA, 1 mU CASP3 (Sigma-Aldrich, St. Louis, MO, USA)) was mixed with 1 µl of the translation mixture that contained a Flag-PK-bis ~ biotin construct, and the mixture was incubated at 30°C for 2 h in a well of a 384-well Optiplate (Perkin Elmer, Foster City, CA, USA). Using the reagents of an AlphaScreen IgG (protein A) detection kit (Perkin Elmer) according to the manufacturer's instructions, 15 µl of 20 mM Tris-HCl, pH 7.5, 0.2 mM DTT, 5 mM MgCl₂, 5 µg/ml anti-FLAG M2 antibody (Sigma-Aldrich), 1 mg/ml BSA, 0.1 µl of streptavidin-coated donor beads and 0.1 µl of anti-IgG acceptor beads were added to the well. The solution was incubated at 23°C for 1 h. Luminescence was analyzed using the AlphaScreen detection program (Perkin Elmer). All repetitive mechanical procedures were performed by a Biomek FX robotic workstation (Beckman Coulter, Fullerton, CA, USA). The value of a luminescent signal is reported as the mean of three independent measurements.

TD immunoblotting. A mixture of each Flag-PK-bis ~ biotin construct (3 µl of a translation mixture) and 7 µl of the CASP3 cleavage solution was incubated at 30°C for 1 h in a well of a 384-well Optiplate (Perkin Elmer). Then, the proteins were separated in SDS-PAGE gels and transferred to PVDF membranes (Millipore Bedford, MA, USA). The blotted proteins were prepared for detection using the reagents of an ECL-Plus Western Blotting Detection System kit (GE Healthcare, Piscataway, NJ, USA), anti-Flag M2 antibodies (Sigma-Aldrich) for N-TD, and Alexa488-streptavidin (Invitrogen) for C-TD. The labeled proteins were visualized using a Typhoon Imager (GE Healthcare) with a 532-nm laser and a 526-nm emission filter or an ImageQuant LAS-4000 mini CCD camera system (Fujifilm).

Sequencing and other purification procedures. When possible, long biotinylated C-terminal fragments produced by CASP3 cleavage were recovered attached to streptavidin beads, and then sequenced directly. When a PK construct had a low biotin-labeling efficiency and was cleaved near its C-terminus, a new construct was made by fusing the GST nucleotide sequence encoded in the pEU-E01-Gateway-GST vector to the C-terminal codon of the corresponding PK open-reading frame using the Gateway system and the pEU-E01-Gateway-GST vector. For purification, synthesized PKs (1.2 ml) were purified using Streptavidin Magnesphere Paramagnetic beads (Promega Corp., Madison, WI, USA) for the Flag-PK-bis ~ biotin constructs or glutathione Sepharose 4B (GE Healthcare) for the PK-GST constructs. After washing the beads with PBS, the bound PKs were incubated with CASP3 (15 µl of total volume) as described above. The samples were boiled and the proteins separated by SDS-PAGE. After blotting and visualization (ProBlott, Applied Biosystems, Foster City, CA, USA), the membrane areas that contained the cleaved fragments were cut out and the fragments were sequenced (Applied Biosystems ABI 473A). CSK kinase (Carna Biosciences Inc., Kobe, Japan) and AMPKα1 (Cell Signaling Technology, Beverly, MA, USA) were cleaved with CASP3 (10 µl of total volume), and the cleavage products subjected to MALDI/TOF-MS (Shimazu Techno-Research Inc., Kyoto, Japan) for sequencing. D → A mutagenesis was carried out using the reagents of a PrimeSTAR Mutagenesis Basal kit (TakaraBio, Otsu, Japan) according to the manufacturer's instructions. The mutated genes were sequenced using an ABI PRISM 310 DNA sequencer (Applied Biosystems).

Construction of PK expression plasmids for the cell-based assay. Expression plasmids were produced using the Gateway method. To obtain the attB1-PK-Flag-(stop codon)-attB2 for Gateway BP Clonase II recombination, the open-reading frame products of the 30 newly identified PK substrates of CASP3 that had been produced by PCR using the S1 and T1 primers as described above were PCR amplified using the primers, attB1-S1 (5'-GGGGACAAGTTTGTACAAAAAGCAGGCTTCACCCACCACCACCA-3') and T1-Flag-stop-attB2 (5'-GGGGACCACCTTTGTACAAGAAAGCTGGGTTTACTGTGCATCGTCATCCTTG TAGTCGCTTCCAGCACTAGCTCCAGA-3'). These PCR products were each inserted into a pDEST26 vector (Invitrogen) using the Gateway system for construction of the His-PK-Flag nucleotide sequences. All sequences were confirmed by DNA sequencing as described above.

Cell-based assay. HeLa cells were cultured in Dulbecco's modified Eagle's medium, 10% fetal bovine serum, penicillin (100 mg/ml), and streptomycin (50 µg/ml). Transient transfections were carried out using Lipofectamine 2000 Transfection Reagent (Invitrogen) according to the manufacturer's instructions. At 24 h after transfection, cells were harvested after apoptosis was induced. Control cells were treated with DMSO and, for apoptosis induction or inhibition, with 20 ng/ml TNF α

(Calbiochem, La Jolla, CA, USA) and 100 μ M cycloheximide (Chemicon, Temecula, CA, USA) or 125 ng/ml anti-Fas antibody (Medical & Biological Laboratories Co., Ltd., Nagoya, Japan) in the presence (inhibition) or absence (induction) of 100 μ M Z-VAD-FMK (Peptide Institute Inc., Osaka, Japan) for 6 h. Cells were washed with PBS and then lysed directly by adding one volume of 2 \times SDS-PAGE sample buffer (125 mM Tris-HCl, pH 6.8, 20% glycerol, 4% SDS, 10% 2-mercaptoethanol, 0.001% bromophenol blue) before subjecting the cell extracts to SDS-PAGE and immunoblotting, which used anti-His antibodies (GE Healthcare) or anti-Flag M2 antibodies (Sigma-Aldrich). The following antibodies were employed to detect endogenous proteins: anti- α -tubulin (Sigma-Aldrich); anti-AKT2, anti-eEF2K, anti-AMPKa1, and anti-TGF β R1 (Cell Signaling Technology). Chemiluminescent signals, generated by ECL-Plus reagents (GE Healthcare), or Immobilon Western HRP substrate Luminol Reagent (Millipore), were detected using an LAS-4000 mini biomolecular imager (GE Healthcare).

Conflict of interest

Dr. Endo is a founder of CellFree Sciences Co., Ltd. and a member of its scientific advisory board. Other authors declare no conflict of interest.

Acknowledgements. We thank Professor Akihide Ryo (Yokohama City University, Japan) for his useful comments and suggestions concerning the cell analysis, Mr. Tatsuya Akagi for technical assistance, and Dr. Morishita (CellFree Sciences) for assistance with the robotic operations. This work was partially supported by the Special Coordination Funds for Promoting Science and Technology by the Ministry of Education, Culture, Sports, Science, and Technology, Japan (Nos. 19657041 and 22310127 to TS).

- Puente XS, Sanchez LM, Overall CM, Lopez-Otin C. Human and mouse proteases: a comparative genomic approach. *Nat Rev Genet* 2003; **4**: 544–558.
- Hunter T. Signaling – 2000 and beyond. *Cell* 2000; **100**: 113–127.
- Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S. The protein kinase complement of the human genome. *Science* 2002; **298**: 1912–1934.
- Matrisian LM, Hogan BL. Growth factor-regulated proteases and extracellular matrix remodeling during mammalian development. *Curr Top Dev Biol* 1990; **24**: 219–259.
- Turgeon VL, Houenou LJ. The role of thrombin-like (serine) proteases in the development, plasticity and pathology of the nervous system. *Brain Res Brain Res Rev* 1997; **25**: 85–95.
- van Kempen LC, de Visser KE, Coussens LM. Inflammation, proteases and cancer. *Eur J Cancer* 2006; **42**: 728–734.
- Friedl P, Wolf K. Tube travel: the role of proteases in individual and collective cancer cell invasion. *Cancer Res* 2008; **68**: 7247–7249.
- Abdel-Rahman HM, Kimura T, Hidaka K, Kiso A, Nezami A, Freire E *et al*. Design of inhibitors against HIV, HTLV-I, and *Plasmodium falciparum* aspartic proteases. *Biol Chem* 2004; **385**: 1035–1039.
- Alnemri ES. Mammalian cell death proteases: a family of highly conserved aspartate specific cysteine proteases. *J Cell Biochem* 1997; **64**: 33–42.
- Hengartner MO. The biochemistry of apoptosis. *Nature* 2000; **407**: 770–776.
- Mahrus S, Trinidad JC, Barkan DT, Sali A, Burlingame AL, Wells JA. Global sequencing of proteolytic cleavage sites in apoptosis by specific labeling of protein N termini. *Cell* 2008; **134**: 866–876.
- Dix MM, Simon GM, Cravatt BF. Global mapping of the topography and magnitude of proteolytic events in apoptosis. *Cell* 2008; **134**: 679–691.
- Righetti PG, Boschetti E. Sherlock Holmes and the proteome – a detective story. *FEBS J* 2007; **274**: 897–905.
- Sawasaki T, Ogasawara T, Morishita R, Endo Y. A cell-free protein synthesis system for high-throughput proteomics. *Proc Natl Acad Sci USA* 2002; **99**: 14652–14657.

- Sawasaki T, Morishita R, Gouda MD, Endo Y. Methods for high-throughput materialization of genetic information based on wheat germ cell-free expression system. *Methods Mol Biol* 2007; **375**: 95–106.
- Takai K, Sawasaki T, Endo Y. Practical cell-free protein synthesis system using purified wheat embryos. *Nat Protoc* 2010; **5**: 227–238.
- Endo Y, Sawasaki T. Cell-free expression systems for eukaryotic protein production. *Curr Opin Biotechnol* 2006; **17**: 373–380.
- Sawasaki T, Hasegawa Y, Morishita R, Seki M, Shinozaki K, Endo Y. Genome-scale, biochemical annotation method based on the wheat germ cell-free protein synthesis system. *Phytochemistry* 2004; **65**: 1549–1555.
- Goshima N, Kawamura Y, Fukumoto A, Miura A, Honma R, Satoh R *et al*. Human protein factory for converting the transcriptome into an *in vitro*-expressed proteome. *Nat Methods* 2008; **5**: 1011–1017.
- Sawasaki T, Kamura N, Matsunaga S, Saeki M, Tsuchimochi M, Morishita R *et al*. Arabidopsis HY5 protein functions as a DNA-binding tag for purification and functional immobilization of proteins on agarose/DNA microplate. *FEBS Lett* 2008; **582**: 221–228.
- Takahashi H, Nozawa A, Seki M, Shinozaki K, Endo Y, Sawasaki T. A simple and high-sensitivity method for analysis of ubiquitination and polyubiquitination based on wheat cell-free protein synthesis. *BMC Plant Biol* 2009; **9**: 39.
- Matsuoka K, Komori H, Nose M, Endo Y, Sawasaki T. Simple screening method for autoantigen proteins using the N-terminal biotinylated protein library produced by wheat cell-free synthesis. *J Proteome Res* 2010; **9**: 4264–4273.
- Fischer U, Janicke RU, Schulze-Osthoff K. Many cuts to ruin: a comprehensive update of caspase substrates. *Cell Death Differ* 2003; **10**: 76–100.
- Kurokawa M, Kornbluth S. Caspases and kinases in a death grip. *Cell* 2009; **138**: 838–854.
- Lee KK, Murakawa M, Nishida E, Tsubuki S, Kawashima S, Sakamaki K *et al*. Proteolytic activation of MST/Krs, STE20-related protein kinase, by caspase during apoptosis. *Oncogene* 1998; **16**: 3029–3037.
- Tomiyoshi G, Horita Y, Nishita M, Ohashi K, Mizuno K. Caspase-mediated cleavage and activation of LIM-kinase 1 and its role in apoptotic membrane blebbing. *Genes Cells* 2004; **9**: 591–600.
- Tatsuta T, Shiraishi A, Mountz JD. The prodomain of caspase-1 enhances Fas-mediated apoptosis through facilitation of caspase-8 activation. *J Biol Chem* 2000; **275**: 14248–14254.
- Packard BZ, Komoriya A, Brotz TM, Henkart PA. Caspase 8 activity in membrane blebs after anti-Fas ligation. *J Immunol* 2001; **167**: 5061–5066.
- Lahti JM, Xiang J, Heath LS, Campana D, Kidd VJ. PITSLRE protein kinase activity is associated with apoptosis. *Mol Cell Biol* 1995; **15**: 1–11.
- Emoto Y, Manome Y, Meinhardt G, Kasaki H, Kharbanda S, Robertson M *et al*. Proteolytic activation of protein kinase C delta by an ICE-like protease in apoptotic cells. *EMBO J* 1995; **14**: 6148–6156.
- Casciola-Rosen LA, Anhalt GJ, Rosen A. DNA-dependent protein kinase is one of a subset of autoantigens specifically cleaved early during apoptosis. *J Exp Med* 1995; **182**: 1625–1634.
- Pearce LR, Komander D, Alessi DR. The nuts and bolts of AGC protein kinases. *Nat Rev Mol Cell Biol* 2010; **11**: 9–22.
- Ohoka N, Sakai S, Onozaki K, Nakanishi M, Hayashi H. Anaphase-promoting complex/cyclosome-cdh1 mediates the ubiquitination and degradation of TRB3. *Biochem Biophys Res Commun* 2010; **392**: 289–294.
- Sawasaki T, Hasegawa Y, Tsuchimochi M, Kamura N, Ogasawara T, Kuroita T *et al*. A bilayer cell-free protein synthesis system for high-throughput screening of gene products. *FEBS Lett* 2002; **514**: 102–105.
- Kamura N, Sawasaki T, Kasahara Y, Takai K, Endo Y. Selection of 5'-untranslated sequences that enhance initiation of translation in a cell-free protein synthesis system from wheat embryos. *Bioorg Med Chem Lett* 2005; **15**: 5402–5406.
- Crooks GE, Hon G, Chandonia JM, Brenner SE. WebLogo: a sequence logo generator. *Genome Res* 2004; **14**: 1188–1190.



Cell Death and Disease is an open-access journal published by Nature Publishing Group. This work is licensed under the Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Supplementary Information accompanies the paper on Cell Death and Disease website (<http://www.nature.com/cddis>)