

TTC5 is required to prevent apoptosis of acute myeloid leukemia stem cells

JT Lynch¹, TDD Somerville¹, GJ Spencer¹, X Huang¹ and TCP Somerville^{*1}

Using a screening strategy, we identified the tetratricopeptide repeat (TPR) motif protein, Tetratricopeptide repeat domain 5 (TTC5, also known as stress responsive activator of p300 or Strap) as required for the survival of human acute myeloid leukemia (AML) cells. TTC5 is a stress-inducible transcription cofactor known to interact directly with the histone acetyltransferase EP300 to augment the TP53 response. Knockdown (KD) of *TTC5* induced apoptosis of both murine and human AML cells, with concomitant loss of clonogenic and leukemia-initiating potential; KD of *EP300* elicited a similar phenotype. Consistent with the physical interaction of TTC5 and EP300, the onset of apoptosis following KD of either gene was preceded by reduced expression of *BCL2* and increased expression of pro-apoptotic genes. Forced expression of *BCL2* blocked apoptosis and partially rescued the clonogenic potential of AML cells following *TTC5* KD. KD of both genes also led to the accumulation of MYC, an acetylation target of EP300, and the form of MYC that accumulated exhibited relative hypoacetylation at K148 and K157, residues targeted by EP300. In view of the ability of excess cellular MYC to sensitize cells to apoptosis, our data suggest a model whereby TTC5 and EP300 cooperate to prevent excessive accumulation of MYC in AML cells and their sensitization to cell death. They further reveal a hitherto unappreciated role for TTC5 in leukemic hematopoiesis.

Cell Death and Disease (2013) 4, e573; doi:10.1038/cddis.2013.107; published online 4 April 2013

Subject Category: Cancer

Acute myeloid leukemia (AML) is a genetically heterogeneous clonal disorder characterized by the accumulation of poorly differentiated myeloid precursor cells in the bone marrow (BM) and blood.¹ Epigenetic dysfunction is central to the molecular pathology of the disease, as evidenced by the recurrent mutations in genes coding for proteins that directly or indirectly regulate the structure and function of chromatin.² Commonly occurring exemplar mutations in myeloid cancers include those targeting the gene coding for the histone H3K4 methyltransferase MLL, either by chromosomal translocation or partial tandem duplication; those inactivating the enzymatic activities of DNA methyltransferase 3A or the methylcytosine dioxygenase TET2; or those targeting isocitrate dehydrogenase 1 or 2, which result in the production by the mutant enzyme of a neometabolite, 2-hydroxyglutarate, which inhibits the activities of TET-family proteins and Jumonji-domain histone demethylases.² Further understanding of the structure and function of chromatin in myeloid leukemia cells may lead to the identification of novel therapeutic targets and strategies.

In the course of our studies, we performed a lentiviral short hairpin RNA (shRNA) genetic knockdown screen to identify regulators of chromatin structure and function required for the proliferation or survival of human AML cells (Huang *et al.*, 2013,

manuscript submitted). The screen identified tetratricopeptide repeat domain 5 (*TTC5*; also known as stress responsive activator of p300, Strap) as required for leukemia cell survival. *TTC5* is a tetratricopeptide repeat (TPR) motif protein known to have important roles in DNA damage and heat-shock responses. It was originally discovered in complex with an EP300-interacting protein called junction mediating and regulatory protein (JMY), and the histone acetyltransferase (HAT) EP300 (E1A-binding protein p300; also known as p300) itself.³ *TTC5* augments the interaction of JMY and EP300, increasing the HAT activity of EP300. This leads to enhanced TP53 acetylation, stability and transcriptional activation during the DNA damage response.³ *TTC5* is phosphorylated by ataxia telangiectasia mutated (ATM) at S203 and checkpoint kinase 2 (CHK2) at S221. These phosphorylations promote its stability and nuclear accumulation, also augmenting TP53 activity and promoting apoptosis in etoposide-treated cells.^{4,5} In the heat shock response, *TTC5* associates with heat shock factor 1 (HSF1) in a chromatin-bound complex that includes EP300. The complex is located at the promoters of heat shock genes and is required to prevent apoptosis.⁶ A physical interaction of *TTC5* with Protein Methyltransferase 5 (PRMT5), which arginine methylates TP53, has been reported,⁷ as has a regulatory role for

¹Cancer Research UK Leukaemia Biology Laboratory, Paterson Institute for Cancer Research, The University of Manchester, Manchester, UK

*Corresponding author: TCP Somerville, Cancer Research UK Leukaemia Biology Laboratory, Paterson Institute for Cancer Research, Manchester M20 4BX, UK. Tel: +44 161 918 7160; Fax: +44 161 446 3109; E-mail: tsomerville@picr.man.ac.uk

Keywords: TTC5; EP300; MYC; leukemia

Abbreviations: 7-AAD, 7-aminoactinomycin; AML, acute myeloid leukemia; ATM, ataxia telangiectasia mutated; BM, bone marrow; CFC, colony forming cell; CHEK2, checkpoint kinase 2; CHX, cycloheximide; EP300, E1A-binding protein p300; EV, empty vector; FACS, fluorescence-activated cell sorter; HSF1, heat-shock factor 1; IP, immunoprecipitation; JMY, junction mediating and regulatory protein; KD, knockdown; LSC, leukemia stem cell; NS, not significant; NTC, non-targeting control; PTMs, post-translational modifications; Q-PCR, quantitative reverse transcriptase PCR; SEM, standard error of the mean; shRNA, short hairpin RNA; TPR, tetratricopeptide repeat; TTC5, tetratricopeptide repeat domain 5; TP53, tumor protein 53; MDM2, murine double minute 2

Received 06.12.12; revised 10.2.13; accepted 26.2.13; Edited by P Salomoni

TTC5 in steroid hormone signaling where it acts as a cofactor for the glucocorticoid and estrogen receptors.⁸

While TTC5 is ubiquitously expressed, including in hematopoietic cells (data not shown), to date there is no information as to any functional role for TTC5 in normal or leukemic hematopoiesis. Here, we present evidence demonstrating a critical role for TTC5 in prevention of apoptosis of human AML cells.

Results

TTC5 is required to prevent apoptosis of AML cells. To confirm the preliminary results from our knockdown screen (data not shown), human THP1 AML cells were infected with lentiviral vectors targeting *TTC5* for KD. THP1 cells exhibit a t(9;11) translocation, which is the cytogenetic hallmark of *MLL-AF9*, a fusion oncogene found in 4–5% of patients with AML.¹ Successfully transduced cells, as indicated by puromycin drug resistance, were then cultured in semisolid conditions to assess their clonogenic potential. Untransduced THP1 cells treated with puromycin are killed by drug within 48 h (data not shown). While control cells that were infected with a vector expressing a non-targeting hairpin formed plentiful colonies, cells infected with vectors expressing hairpins targeting *TTC5* for KD formed significantly fewer colonies (Figure 1a). *TTC5* KD was confirmed at both the mRNA and protein levels, and the extent of *TTC5* KD

correlated with the observed reduction in clonogenic potential (Figures 1b and c). To confirm that the phenotype was an on-target effect of *TTC5* KD, we performed repeat experiments using THP1 cells that stably expressed a version of human *TTC5* that lacked the 3'-UTR sequences targeted by KD constructs #1 and #4 (Figure 1d). A partial (construct #1) or complete (construct #4) rescue of clonogenic potential was observed (Figure 1e), providing strong additional evidence that the observed phenotype was specific for *TTC5*. The loss of clonogenic potential following *TTC5* KD was in large part due to induction of apoptosis (Figure 1f), although a modest but statistically significant G1 arrest with reduction in the proportion of cells in S phase was also observed (Figure 1g). Thus, *TTC5* is required to prevent apoptosis and maintain the proliferative and clonogenic potential of human AML cells.

To determine whether *TTC5* is required for the leukemia-initiating potential of human AML cells, we transplanted control or KD THP1 AML cells into sublethally irradiated immune-deficient mice. All mice injected with control cells died of short latency AML, whereas mice injected with *TTC5* KD cells did not succumb to leukemia (Figures 2a and b). There was no evidence of leukemic engraftment in any of the mice euthanized at the termination of the experiment (Figure 2b and data not shown). To determine whether the observed dependency on *TTC5* was specific to the *MLL* molecular subtype of AML or a more general feature of myeloid leukemia cells, *TTC5* KD was initiated in additional human AML cell lines. Significant loss of clonogenic potential of AML cells was




Figure 1 *TTC5* prevents apoptosis of THP1 *MLL-AF9* AML cells. THP1 cells were infected with lentiviruses expressing shRNAs targeting *TTC5* for KD, or a non-targeting control shRNA (NTC), with puromycin drug resistance as the selectable marker. Next day, successfully infected cells were drug-selected in puromycin (3 μ g/ml) for 48 h. Bar charts show (a) mean \pm S.E.M. colony forming cell (CFC) frequencies of drug-resistant cells enumerated after 10 days in semisolid culture ($n = 3$); and (b) mean \pm S.E.M. *TTC5* mRNA levels in KD cells relative to control cells, 72 h after lentiviral infection. Representative western blots show expression of (c) endogenous *TTC5* and ACTB in KD and control cells, 7 days after lentiviral infection or (d) endogenous or ectopically expressed *TTC5* in THP1 cell lines stably infected with the indicated vectors. Vertical dashed line indicates intervening gel lanes excised for clarity of presentation. Bar charts show (e) mean \pm S.E.M. CFC frequencies in control or KD cells with or without ectopic expression of human *TTC5* ($n = 3$); (f) mean \pm S.E.M. percentage of apoptotic cells, as determined by annexin V binding, 7 days after lentiviral infection of THP1 cells ($n = 3$) and (g) mean \pm S.E.M. percentage of control or KD cells in the indicated phase of the cell cycle ($n = 5$). S.E.M. = standard error of the mean; EV = Empty Vector; *TTC5* O/E = force-expressed *TTC5*; * indicates $P \leq 0.05$ for comparison of KD versus control conditions, or for the indicated comparisons, using one-way ANOVA and Fisher's least significant difference *post hoc* test for (a), (b), (f) and (g) and an unpaired *t*-test for (e)




Figure 2 *TTC5* is required for the leukemia-initiating potential of AML cells. THP1 human AML cells were infected with lentiviruses expressing shRNAs targeting *TTC5* for KD, or a non-targeting control (NTC) shRNA, with puromycin drug resistance as the selectable marker. (a) Survival curves of sublethally irradiated immune-deficient mice transplanted with 5000 viable drug resistant control or *TTC5* KD THP1 cells ($n = 5$ per cohort). (b) Representative FACS plots show percentage human CD45⁺ THP1 AML cells in the BM of recipient mice either at death (NTC) or at termination of the experiment on day 125 (*TTC5* KD). (c) Bar chart shows mean \pm S.E.M. colony-forming cell (CFC) frequencies of drug-resistant *TTC5* KD cells relative to control cells enumerated after 10 days in semisolid culture for the indicated AML cell lines. * Indicates $P \leq 0.05$ for comparison of KD versus control conditions using one-way ANOVA and Fisher's least significant difference *post hoc* test

observed following *TTC5* KD in both MLL mutated lines (MonoMac1 and MV4;11) and lines representative of other AML molecular subtypes (NB4 and K562) (Figure 2c). There was also a reduction in the clonogenic potential of *TTC5* KD HL60 cells, although this did not reach statistical significance. These data demonstrate that *TTC5* is required for the leukemia-initiating potential of human AML cells and that the requirement for *TTC5* is not specific to the MLL mutated molecular subtype.

To confirm and expand these observations in a separate species, we performed similar experiments using a mouse model of human MLL-AF9 leukemia, which faithfully recapitulates many of the clinical and pathological features of the human disease.⁹ Three separate lentivirally expressed shRNAs were used to target murine *Ttc5* in murine MLL-AF9 AML cells, this time with GFP as the selectable

marker. The extent of transcript KD, as determined by Q-PCR, correlated once more with significant loss of clonogenic potential (Figures 3a and b). To determine whether the leukemia stem cell (LSC) compartment of the leukemia clone was targeted by *Ttc5* KD, we performed secondary transplantation of FACS-purified control or *Ttc5* KD MLL-AF9 AML cells 48 h following lentiviral infection.¹⁰ As for human THP1 cells, all mice receiving control AML cells died of short latency disease, whereas mice transplanted with *Ttc5* KD AML cells survived until the termination of the experiment (Figure 3c), and at autopsy exhibited no evidence of leukemic engraftment, as determined by flow cytometry of bone marrow cells. By contrast, engraftment of donor GFP⁺ AML cells in the BM of leukemic mice was readily confirmed (Figure 3d). These data confirm that, as for human AML cells, *Ttc5* is essential for murine MLL-AF9 leukemia-initiating cells.




Figure 3 *Ttc5* is required to maintain the leukemia-initiating potential of murine MLL-AF9 AML cells. MLL-AF9 cells were infected with lentiviruses expressing shRNAs targeting *Ttc5* for KD or a non-targeting control (NTC) shRNA, with GFP as the selectable marker. After 48 h, cells were FACS-purified for GFP expression. Bar charts show (a) mean \pm S.E.M. CFC frequencies of control and *Ttc5* KD cells after 5 days in semisolid culture ($n=3$) and (b) mean \pm S.E.M. *Ttc5* mRNA levels in KD cells relative to control cells 48 h following lentiviral infection ($n=3$). (c) Survival curves of sublethally irradiated syngeneic mice secondarily transplanted with 1000 control or *Ttc5* KD AML cells ($n=4$ mice per cohort). (d) Representative FACS plots show the presence of GFP⁺ donor AML cells in the BM of a leukemic mouse transplanted with control cells (Day 60) and absence of AML cells in a mouse transplanted with *Ttc5* KD cells at termination of the experiment (Day 120). * Indicates $P \leq 0.05$ for comparison of KD versus control conditions using one-way ANOVA and Fisher's least significant difference *post hoc* test

TTC5 sustains expression of *BCL2* in AML cells. To address the mechanism by which *TTC5* KD cells undergo apoptosis, we determined transcript levels of a panel of genes coding for proteins that regulate cell death 96 and 120 h following lentiviral infection of THP1 AML cells. Relative to that of *ACTB*, expression of the anti-apoptotic gene *BCL2* decreased by $>50\%$ and expression of pro-apoptotic genes such as *BAK1*, *BOK*, *BIK*, *BAD*, *BID*, *PUMA* (also known as *BBC3*) and *BMF* increased by $>50\%$ in *TTC5* KD cells by comparison with control cells at one or the other time point (Figure 4a). The change in expression of other genes was less marked. Reduced *BCL2* expression was confirmed in repeat experiments with two separate *TTC5* KD constructs at both protein (Figure 4b) and transcript levels (Figures 4c and d). To determine whether reduced expression of *BCL2* contributes to apoptosis of AML cells following *TTC5* KD, we force-expressed *BCL2* in THP1 cells and repeated KD. Interestingly, we observed an increase in both *BCL2* transcripts and protein in *BCL2*-overexpressing THP1 cells following *TTC5* KD (Figures 4b and d), suggesting that *TTC5* may directly or indirectly contribute to repression of the integrated retroviral long terminal repeat promoter. Forced expression of *BCL2* (Figures 4b–d) blocked apoptosis and partially rescued the loss of clonogenic potential consequent upon *TTC5* KD (Figures 4e and f). However, forced expression of *BCL2* did not rescue the modest but significant G1 arrest

and reduction of the proportion of cells in S phase, likely explaining why only a partial rescue was observed (Figure 4g). Thus, *TTC5* selectively and positively regulates expression of *BCL2* in human AML cells, either directly or indirectly, and downregulation of *BCL2* following *TTC5* KD contributes to apoptosis and loss of clonogenic potential.

EP300 sustains the expression of *BCL2* in AML cells. *TTC5* was originally identified in complex with EP300 and an EP300-interacting protein JMY,³ and was also found to augment EP300 histone acetyltransferase activity in U2OS osteosarcoma cells.⁶ We therefore hypothesized that *TTC5* might similarly regulate EP300 activity in human THP1 AML cells to enhance cell survival. Consistent with this, *EP300* KD to $51 \pm 8\%$ (mean \pm S.E.M.; $n=3$) of control levels induced a similar phenotype to *TTC5* KD: *EP300* KD cells exhibited reduced *BCL2* expression (Figures 5a and b), a $>50\%$ increase in expression of the pro-apoptotic genes *BID*, *PUMA* and *BMF* (which were also induced by *TTC5* KD) (Figure 5c), and a failure to form colonies in semisolid culture (Figure 5d) due to induction of apoptosis (Figure 5e). These data suggest that *TTC5* and EP300 cooperate, directly or indirectly, to maintain expression of *BCL2* in AML cells and repress pro-apoptotic genes such as *BID*, *PUMA* and *BMF*, thus preventing leukemia cell apoptosis.




Figure 4 TTC5 prevents apoptosis of THP1 cells through regulation of *BCL2* expression. THP1 cells were infected with lentiviruses expressing shRNAs targeting *TTC5* for KD, or a non-targeting control (NTC) shRNA, with puromycin drug resistance as the selectable marker. Bar chart shows (a) mean \pm S.E.M. transcript expression levels of the indicated genes in KD cells relative to control cells at the indicated time points following lentiviral infection ($n=2$). (b) Representative western blots show expression of the indicated proteins, 7 days after lentiviral infection. Bar charts show (c) mean \pm S.E.M. *TTC5* or (d) *BCL2* transcript levels in control (EV, empty vector) or *BCL2* overexpressing *TTC5* KD THP1 cells relative to control cells, 72 h after lentiviral infection ($n=3$). Bar charts show (e) mean \pm S.E.M. percentage of apoptotic cells, as determined by annexin V binding, 7 days after lentiviral infection ($n=3$) and (f) mean \pm S.E.M. colony-forming cell (CFC) frequencies of drug-resistant cells enumerated following 10 days in semisolid culture ($n=3$). * Indicates $P \leq 0.05$ (Mann-Whitney *U*-test) for comparison of KD versus control conditions, or for the indicated comparisons. NS = not significant. (g) Bar chart shows mean \pm S.E.M. percentage of *TTC5* KD cells in G1 or S phase of cell cycle relative to that observed in control cells (for *BCL2* overexpressing and control THP1 lines) 96 h after initiation of KD ($n=3$). * Indicates $P \leq 0.05$ for comparison of KD versus control conditions using one-way ANOVA and Fisher's least significant difference *post hoc* test

TTC5 and EP300 regulate turnover of MYC in AML cells.

The pattern of change in expression of *BCL2* family genes following *TTC5* KD (Figure 4a), with reduced expression of *BCL2* and increased expression of pro-apoptotic genes, was reminiscent of the pattern we have previously observed in AML cells in which *Myc* was force-expressed (Huang *et al*, 2013, manuscript submitted). Furthermore, EP300 regulates the stability of MYC through both acetylation-dependent and acetylation-independent mechanisms.¹¹ These data prompted us to investigate whether *TTC5* or *EP300* KD altered the expression of MYC. Neither *TTC5* KD nor *EP300* KD in THP1 AML cells led to a significant change in the expression of MYC transcripts (Figures 6a and b). By contrast, both *TTC5* KD and *EP300* KD led to a robust accumulation of MYC protein (Figure 6c), which was explained by an increase in its cellular half-life from <30 min

(control cells) to >60 min (*EP300* KD) (Figures 6d and e). Given that EP300 directly acetylates MYC at K148, K157 and K323,¹¹ we next investigated whether acetylation of MYC at these residues was perturbed in *TTC5* or *EP300* KD cells. In contrast to the robust accumulation of total MYC protein, levels of acetyl-K157 MYC did not change following either *TTC5* or *EP300* KD, and levels of acetyl-K148 MYC increased only modestly (Figure 7a); this was confirmed by semiquantitative assessment of western blot bands using ImageJ analysis (Figure 7b). By contrast, acetylation of MYC at K323 was not readily detected in control or *TTC5* KD THP1 cells, although it was detected in *EP300* KD cells. Thus, following *TTC5* or *EP300* KD, the form of MYC that accumulated in human AML cells is relatively hypoacetylated at K148 and K157. These data are consistent with previously described roles for EP300 in MYC




Figure 5 EP300 KD phenocopies TTC5 KD. THP1 cells were infected with lentiviruses expressing shRNAs targeting EP300 for KD, or a non-targeting control shRNA (NTC), with puromycin drug resistance as the selectable marker. (a) Bar chart shows mean \pm S.E.M. *BCL2* mRNA levels in EP300 KD cells relative to control cells, 72 h after lentiviral infection ($n=3$). (b) Representative western blot shows expression of the indicated proteins, 72 h after lentiviral infection ($n=3$). (c) Bar chart shows mean \pm S.E.M. transcript levels of the indicated genes in EP300 KD cells relative to control cells, 72 h after lentiviral infection ($n=3$). Bar charts show (d) mean \pm S.E.M. colony-forming cell (CFC) frequencies of drug-resistant cells enumerated following 10 days in semisolid culture ($n=3$) and (e) mean \pm S.E.M. percentage of apoptotic cells, as determined by annexin V binding, 7 days after lentiviral infection ($n=3$). * indicates $P \leq 0.05$ (Mann-Whitney *U*-test) for comparison of KD versus control conditions

acetylation and turnover¹¹ and TTC5 in regulation of the HAT activity of EP300.^{3,6}

To determine whether TTC5 and MYC interact, co-immunoprecipitation experiments were performed in 293FT cells: an HA-tagged version of TTC5 readily pulled down co-expressed MYC in (Figure 7c). Finally, to determine whether MYC accumulation might contribute to induction of apoptosis, we performed TTC5 KD in THP1 cells where murine *Myc* was force-expressed. Expression of murine *Myc* transcripts was detected in *Myc*-expressing human THP1 cells, but was absent in control cells (data not shown). Consistent with a role for MYC in sensitizing cells to apoptosis,¹² THP1 cells force-expressing *Myc* exhibited

significantly greater levels of apoptosis by comparison with control cells, both when infected with non-targeting control vectors (albeit modestly) and when infected with vectors targeting *TTC5* for KD (Figures 7d and e). The substantially greater proportion of *TTC5* KD versus control cells undergoing apoptosis in the context of *Myc* forced expression furthermore suggested a quantitative link between the amount of cellular MYC and the rapidity of onset of apoptosis. There was no significant difference in the clonogenic potential of control or *Myc* force-expressing *TTC5* KD cells (Figure 7f). Together, these data indicate a role for TTC5 in the regulation of MYC protein turnover in leukemia cells and suggest that MYC accumulation might contribute to the induction of apoptosis through altering the balance of expression of anti-versus pro-apoptotic proteins.

Discussion

TTC5 was originally described as an EP300-interacting transcription cofactor that accumulates following etoposide treatment of U2OS osteosarcoma cells, and which augments the TP53 response.³ Subsequent work has identified more general roles for TTC5 in the regulation of the cellular response to stress, for example, following heat shock.⁶ Our studies using cells from two different species now reveal roles for TTC5 in the prevention of apoptosis of leukemia cells and maintenance of their clonogenic capacity, through regulation of MYC and *BCL2* expression. Syngeneic and xenogeneic transplantation experiments confirm an *in vivo* requirement for TTC5 in leukemia initiation, as well as its *in vitro* requirement. Importantly, our observations are not explained by an effect on TP53 because THP1 cells exhibit a frame-shift deletion of TP53 from R174¹³ and, as expected, we were unable to detect TP53 expression in these cells (data not shown). While the dependency of AML cells on TTC5 was discovered in the context of the *MLL* mutated molecular subtype of AML, the *in vitro* clonogenic activity of AML cell lines representative of other molecular subtypes was likewise dependent on TTC5, suggesting a wide role for the protein in leukemic hematopoiesis.

Knockdown of either *TTC5* or *EP300* in AML cells similarly alters the balance of expression of pro- versus anti-apoptotic genes, consistent with the physical and functional interaction of their gene products. The decrease in expression of the anti-apoptotic gene *BCL2* and increase in expression of pro-apoptotic genes such as *BID*, *PUMA* and *BMF* likely alters the cellular apoptotic rheostat in favor of apoptosis. Such changes might be explained by localized, specific and direct effects of a TTC5/EP300 complex at gene promoters to repress or activate expression respectively. Alternatively, changes in gene expression might be consequent upon altered expression of a protein target of the histone acetyltransferase activity of the TTC5/EP300 complex.

In support of the latter interpretation, we have previously found that forced expression of *Myc* in AML cells induces expression of pro-apoptotic genes (Huang *et al*, 2013, manuscript submitted). In contrast, forced expression of *Myc* in THP1 cells reduces expression of *BCL2* relative to *ACTB* to $44 \pm 12\%$ of control levels (mean \pm S.E.M., $n=3$). The similarity in the change in expression of pro- versus




Figure 6 TTC5 and EP300 promote MYC turnover. THP1 cells were infected with lentiviruses expressing shRNAs targeting *TTC5* or *EP300* for KD, or a non-targeting control shRNA (NTC), with puromycin drug resistance as the selectable marker. Bar charts show (a) mean \pm S.E.M. mRNA levels of *MYC* and *TTC5* in *TTC5* KD cells and (b) mean \pm S.E.M. mRNA levels of *MYC* and *EP300* in *EP300* KD cells relative to control cells, 72 h after lentiviral infection ($n=3$). Representative western blots show (c) expression of *MYC* after *EP300* or *TTC5* KD ($n=3$) and (d) *MYC* half-life, as determined by incubation of control or KD cells with 10 μ g/ml cycloheximide for the indicated times, 72 h after lentiviral infection ($n=2$). Vertical dashed line demarcates separate gels, although exposure time was identical. (e) *MYC* half-life was semiquantified by densitometric analysis using ImageJ software




Figure 7 *Myc* accumulation and apoptosis after *TTC5* or *EP300* KD. THP1 cells were infected with lentiviruses expressing shRNAs targeting *TTC5* or *EP300* for KD, or a non-targeting control shRNA (NTC), with puromycin drug resistance as the selectable marker. (a) Representative western blots show expression of *MYC*, acetylated *MYC* and *ACTB* after *EP300* or *TTC5* KD, 72 h after lentiviral infection. (b) Bar chart shows semiquantitative analysis of acetyl-K148 *MYC* or acetyl-K157 *MYC* signal as a percentage of total *MYC* signal, as determined by ImageJ analysis of the bands shown in (a). (c) 293FT cells were transiently transfected with the indicated constructs. After 48 h, cells were harvested and lysates were immunoprecipitated with an HA-tag antibody. The immunoprecipitated complex was analyzed by western blotting using *MYC* or *TTC5* antibodies. IP = immunoprecipitation. (d–f) Control and *Myc*-overexpressing THP1 cells were infected with lentiviruses expressing shRNAs targeting *TTC5* for KD or a NTC shRNA, with puromycin drug resistance as the selectable marker. (d) Representative FACS plots show annexin V and 7-aminoactinomycin (7-AAD) binding to control and KD cells, 7 days after lentiviral infection. Bar charts show (e) mean \pm S.E.M. percentage of apoptotic cells, as determined by annexin V binding, 7 days after lentiviral infection ($n=3$); and (f) mean \pm S.E.M. colony-forming cell (CFC) frequencies of drug-resistant cells enumerated following 10 days in semisolid culture ($n=3$). EV = empty vector. * indicates $P \leq 0.05$ (paired *t*-test) for comparison of KD versus control conditions

Table 1 Primer and probe sets used for Q-PCR

Gene	Q-PCR primer sequences	Roche probe library number
<i>TTC5 F</i>	ACCGAATTCAGCACAAAGGA	47
<i>TTC5 R</i>	CTAGCAGGAGGGGCGTCT	—
<i>ACTB F</i>	ATTGGCAATGAGCGGTTCC	11
<i>ACTB R</i>	GGATGCCACAGGACTCCAT	—
<i>EP300 F</i>	GATCTGTGTCCCTCACCATGAG	40
<i>EP300 R</i>	AAACAGCCATCACAGACGAA	—
<i>MYC F</i>	GCTGCTTAGACGCTGGATTT	66
<i>MYC R</i>	TAACGTTGAGGGGCATCG	—
<i>BCL2 F</i>	AGTACCTGAACCGGCACCT	75
<i>BCL2 R</i>	GCCGTACAGTTCACAAAGG	—
<i>BCL2L1 (BCL-XL) F</i>	AGCCTTGGATCCAGGAGAA	66
<i>BCL2L1 (BCL-XL) R</i>	AGCGGTTGAAGCGTTCCT	—
<i>MCL1 F</i>	TGGATGGTGGCCTACCTG	28
<i>MCL1 R</i>	CGTCCCGTATAGAGCTGT	—
<i>BCL2A1 (A1) F</i>	CAGGAGAATGGATAAGGCAAA	75
<i>BCL2A1 (A1) R</i>	CCAGCCAGATTTAGGTTCAAA	—
<i>BCL2L2 (BCL-W) F</i>	TGGATGGTGGCCTACCTG	28
<i>BCL2L2 (BCL-W) R</i>	CGTCCCGTATAGAGCTGTG	—
<i>BAX F</i>	AGCAAAGTGGTGTCAAGG	69
<i>BAX R</i>	TCTTGGATCCAGCCCAAC	—
<i>BAK F</i>	TGGTACCTTACCTCTGCAAC	43
<i>BAK R</i>	ATGTCGTCCCGATGATG	—
<i>BIK F</i>	CCCTATGGAGGACTTCGATTC	28
<i>BIK R</i>	GGCTCACGTCCATCTCGT	—
<i>BCL2L11 (BIM) F</i>	CATCGCGGTATTCCGGTTC	70
<i>BCL2L11 (BIM) R</i>	GCTTTGCCATTTGGTCTTTTT	—
<i>BAD F</i>	CGAGTTTGTGGACTCCTTTAAGA	78
<i>BAD R</i>	CACCAGGACTGGAAGACTCG	—
<i>BID F</i>	GTGCTGGGGTCATGATGG	19
<i>BID R</i>	CGACTCACTCCTGGTTCACA	—
<i>BBC3 (PUMA) F</i>	GACCTCAACGCACAGTACGA	51
<i>BBC3 (PUMA) R</i>	CACCTAATTGGGCTCCATCTC	—
<i>PMAIP1 (NOXA) F</i>	GGAGATGCCTGGGAAGAAG	67
<i>PMAIP1 (NOXA) R</i>	CCTGAGTTGAGTAGCACACTCG	—
<i>BMF F</i>	GAGACTCTCCTGGAGTCACC	25
<i>BMF R</i>	CTGGTTGGAACACATCATCCT	—
<i>Ttc5 F</i>	TGCAGCAGATGGAGGAAGTA	98
<i>Ttc5 R</i>	GAGTCACATTCAGTGCCTTCC	—
<i>Actb F</i>	TGACAGGATGCAGAAGGAGA	106
<i>Actb R</i>	CGCTCAGGAGGAGCAATG	—
<i>Myc F</i>	CCTAGTGTGCATGAGGAGA	77
<i>Myc R</i>	TCCACAGACACCACATCAATTT	—

anti-apoptotic genes observed following *Myc*-overexpression or *TTC5/EP300* KD prompted our investigation of MYC as an intermediary of the observed gene expression changes. MYC associates directly with and is acetylated by EP300 on multiple residues, including K148, K157 and K323.¹¹ Acetylation of MYC by EP300 promotes its turnover.¹¹ In keeping with these observations, we observed that *EP300* KD led to robust accumulation of a form of MYC that was relatively hypoacetylated at both K148 and K157; *TTC5* KD gave similar results. Together, these data support a role for *TTC5* in promoting the turnover of MYC though its regulation of the acetyltransferase activity of EP300. Given that TPR-motif proteins function as scaffolding proteins, it is possible that *TTC5* could stabilize the interaction between EP300 and MYC.¹⁴

While physiologic levels of MYC are required for normal cell cycle progression, accumulation of MYC above a certain threshold is associated with induction of apoptosis,¹² through a transcriptional mechanism that results in leak of cytochrome *c* from the mitochondrion into the cytoplasm.¹⁵ We found a modest but significant increase in apoptosis in THP1 cells

where *Myc* was force-expressed. However, following *TTC5* or *EP300* KD, the extent of apoptosis was much greater, consistent with the observed significant and substantial accumulation of endogenous MYC. Nevertheless, our observations of a role for *TTC5* in the regulation of MYC turnover do not rule out other important roles for *TTC5* in leukemic hematopoiesis. For example, because *TTC5* is an EP300-interacting protein, and given the varied and critical roles of EP300 in hematopoiesis, it is possible that *TTC5* regulates other hematopoietic transcriptional complexes.^{16,17}

Materials and Methods

Cell lines, culture and clonogenic assays. THP1 and MonoMac-1 cells were from DMSZ (Braunschweig, Germany). MV(4;11), NB4, HL60 and K562 cells were gifts from Dr. Vaskar Saha and were cultured as recommended (www.dmsz.de). Clonogenic assays of cell lines were performed in methylcellulose medium (H4320, Stem Cell Technologies, Vancouver, BC, Canada) with no supplemental growth factors. Liquid culture of murine AML cells was in RPMI 1640 containing 20% fetal calf serum and 5% X63 supernatant.¹⁸ Clonogenic assays of murine AML cells were performed in methylcellulose medium (M3231, Stem Cell Technologies) containing 20 ng/ml SCF, 10 ng/ml IL6, 10 ng/ml GM-CSF and 10 ng/ml IL3 (PeproTech, London, UK).

Reagents, plasmids and lentiviral production. Puromycin, cycloheximide and lentiviral vectors were purchased from Sigma (Poole, UK). Vectors were as follows: non-targeting control pLKO.1 (SHC002); *TTC5* KD pLKO.1 (TRCN0000156809 (#1), TRCN0000156289 (#2) and TRCN0000152678 (#4)); *Ttc5* KD pLKO.1 (TRCN0000176549 (#1), TRCN0000178101 (#2) and TRCN0000181231 (#3)) and *EP300* KD pLKO.1 (TRCN0000039886). For the murine *Ttc5* KD constructs, the PGK-puromycin cassette was exchanged for an SFFV-eGFP cassette (amplified from pHR'SIN.cPPT-SEW, a gift from Dr. Adrian Thrasher), to generate eGFP versions. To generate a lentiviral vector for expression of *TTC5*, *TTC5* cDNA was PCR amplified using pcDNA3-HA-TTC5 as a template (a gift from Marija Krstic-Demonacos), incorporating Cla1 and Xba1 restriction sites, using the following primers:

- 5' ATAATATCGATATGATGGCTGATGAAGAGGAAGAAGTC,
- 3'ATAATTCTAGATCATTACACTGTGGTCGCGATG.

The amplified fragment was sequence-verified and cloned in to the Cla1 and Xba1 sites of pLentiGS downstream of an EF1 α promoter (Huang *et al*, 2013, manuscript submitted). pMSCV-BCL2-IRES-GFP (#8972) and pMSCV-Myc-IRES-GFP (#18770) were purchased from Addgene (Cambridge, MA, USA). Lentiviral and retroviral supernatants were prepared and leukemic human and murine cells infected with viral particles as described.¹⁹

Western blotting and immunoprecipitation. For western blotting, cells were lysed in High Salt Lysis Buffer (45 mM HEPES (pH 7.5), 400 mM NaCl, 1 mM EDTA, 10% Glycerol, 0.5% NP40, 6.25 mM NaF, 20 mM β -glycerophosphate, 1 mM DTT, 20 mM sodium butyrate and 1 \times Protease Inhibitor cocktail (Roche, Burgess Hill, UK)) and equal amounts of protein were loaded and separated by SDS-PAGE. For co-immunoprecipitation, cells were lysed in NP40 Lysis Buffer (50 mM Tris/HCl (pH 8.0), 50 mM KCl, 10 mM EDTA, 1% NP40, 6.25 mM NaF, 20 mM β -glycerophosphate, 1 mM DTT and 1 \times Protease Inhibitor cocktail (Roche)). Lysate was incubated overnight with the appropriate antibody and immune complexes were pulled down with Protein G-Sepharose (Sigma). Samples were washed four times with IP wash buffer (50 mM Tris/HCl (pH 7.5), 150 mM NaCl, 5 mM EDTA and 0.1% Tween 20), eluted and separated by SDS-PAGE. Horseradish peroxidase-linked secondary antibodies (GE Healthcare, Little Chalfont, UK) and ECL (enhanced chemiluminescence; GE Healthcare) or Supersignal (Pierce, Rockford, IL, USA) were used to detect immune complexes. Protein expression levels were quantified using ImageJ software (NIH, USA). Antibodies used for western blotting or immunoprecipitations were as follows: anti-TTC5 (PAB12888, Abnova, Cambridge, UK), anti-ACTB (C4, Millipore, Watford, UK), anti-BCL2 (#124, Dako, Ely, UK), anti-MYC (Y69, Abcam, Cambridge, UK), anti-acetyl MYC K148 (ABE25, Millipore), anti-acetyl MYC K157 (ABE27, Millipore), anti-acetyl MYC K323 (ABE26, Millipore) and anti-HA (12CA5, Abcam).

RNA isolation and analysis. RNA was extracted using QIAshredder spin columns and an RNeasy Plus Micro kit (Qiagen, Manchester, UK). For quantitative PCR, first strand synthesis was performed using a High Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Paisley, UK). Quantitative PCR assays were performed in 384-well MicroAmp optical reaction plates using Taqman Fast Universal PCR Mastermix (Applied Biosystems) and Universal Probe Library System (Roche) designed primers and probes, which are listed in Table 1.

Flow cytometry, apoptosis and cell cycle analysis. FACS analyses were performed using a LSR Model II flow cytometer (BD Biosciences, Oxford, UK). Cell sorting experiments were performed using either an Influx or a FACSAria II flow cytometer (both from BD Biosciences). Anti-CD45-PE-Cy7 (HI30) was from eBioscience (Hatfield, UK). Apoptosis was assessed using a BD Pharmingen APC Annexin V Kit (Oxford, UK), according to the manufacturer's instructions. Propidium iodide cell cycle analyses were performed as described.²⁰

Mice and murine experiments. Experiments were approved by the Paterson Institute's Animal Ethics Committee and performed under a project license issued by the United Kingdom Home Office, in keeping with the Home Office Animal Scientific Procedures Act, 1986. C57BL/6 (CD45.2⁺) mice were purchased from Harlan (Shardlow, UK). NOD.Cg-Prkdc^{scid} Il2rg^{tm1Wjl}/SzJ (NSG) mice were purchased from Jackson Laboratories (Bar Harbor, ME, USA) and bred

in-house. The cohort of mice with syngeneic MLL-AF9 AML, initiated using a retroviral transduction and transplantation protocol, was generated as described.^{10,19}

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements. We thank Morgan Blaylock, Jeff Barry, Mike Hughes, Gail Bruder and Angela Cooke for technical support; and Marija Krstic-Demonacos and Costas Demonacos for plasmids and helpful discussions. This work was supported by Cancer Research UK grant number C5759/A12328.

1. Estey E, Döhner H. Acute myeloid leukaemia. *Lancet* 2006; **368**: 1894–1907.
2. Shih AH, Abdel-Wahab O, Patel JP, Levine RL. The role of mutations in epigenetic regulators in myeloid malignancies. *Nat Rev Cancer* 2012; **12**: 599–612.
3. Demonacos C, Krstic-Demonacos M, La Thangue NB. A TPR motif cofactor contributes to p300 activity in the p53 response. *Mol Cell* 2001; **8**: 71–84.
4. Adams CJ, Graham AL, Jansson M, Coutts AS, Edelmann M, Smith L *et al*. ATM and Chk2 kinase target the p53 cofactor Strap. *EMBO Rep* 2008; **9**: 1222–1229.
5. Demonacos C, Krstic-Demonacos M, Smith L, Xu D, O'Connor DP, Jansson M *et al*. A new effector pathway links ATM kinase with the DNA damage response. *Nat Cell Biol* 2004; **6**: 968–976.
6. Xu D, Zalmas LP, La Thangue NB. A transcription cofactor required for the heat-shock response. *EMBO Rep* 2008; **9**: 662–669.
7. Jansson M, Durant ST, Cho EC, Sheahan S, Edelmann M, Kessler B *et al*. Arginine methylation regulates the p53 response. *Nat Cell Biol* 2008; **10**: 1431–1439.
8. Davies L, Paraskevopoulou E, Sadeq M, Symeou C, Pantelidou C, Demonacos C *et al*. Regulation of glucocorticoid receptor activity by a stress responsive transcriptional cofactor. *Mol Endocrinol* 2011; **25**: 58–71.
9. Somerville TCP, Matheny CJ, Spencer GJ, Iwasaki M, Rinn JL, Witten DM *et al*. Hierarchical maintenance of MLL myeloid leukemia stem cells employs a transcriptional program shared with embryonic rather than adult stem cells. *Cell Stem Cell* 2009; **4**: 129–140.
10. Somerville TCP, Cleary ML. Identification and characterization of leukemia stem cells in murine MLL-AF9 acute myeloid leukemia. *Cancer Cell* 2006; **10**: 257–268.
11. Faiola F, Liu X, Lo S, Pan S, Zhang K, Lyman E *et al*. Dual regulation of c-Myc by p300 via acetylation-dependent control of Myc protein turnover and coactivation of Myc-induced transcription. *Mol Cell Biol* 2005; **25**: 10220–10234.
12. Evan GI, Wyllie AH, Gilbert CS, Littlewood TD, Land H, Brooks M *et al*. Induction of apoptosis in fibroblasts by c-myc protein. *Cell* 1992; **69**: 119–128.
13. Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S *et al*. The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. *Nature* 2012; **483**: 603–307.
14. Allan R, Ratajczak T. Versatile TPR domains accommodate different modes of target protein recognition and function. *Cell Stress Chaperones* 2011; **16**: 353–367.
15. Juin P, Hueber AO, Littlewood T, Evan G. c-Myc-induced sensitization to apoptosis is mediated through cytochrome c release. *Genes Dev* 1999; **13**: 1367–1381.
16. Blobel GA. CREB-binding protein and p300: molecular integrators of hematopoietic transcription. *Blood* 2000; **95**: 745–755.
17. Rebel VI, Kung AL, Tanner EA, Yang H, Bronson RT, Livingston DM. Distinct roles for CREB-binding protein and p300 in hematopoietic stem cell self-renewal. *Proc Natl Acad Sci USA* 2002; **99**: 14789–14794.
18. Karasuyama H, Melchers F. Establishment of mouse cell lines which constitutively secrete large quantities of interleukin 2, 3, 4 or 5, using modified cDNA expression vectors. *Eur J Immunol* 1988; **18**: 97–104.
19. Harris WJ, Huang X, Lynch JT, Spencer GJ, Hitchin JR, Li Y *et al*. The histone demethylase KDM1A sustains the oncogenic potential of MLL-AF9 leukemia stem cells. *Cancer Cell* 2012; **21**: 473–487.
20. Somerville TCP, Lynch DC, Khwaja A. Different levels of p38 MAP kinase activity mediate distinct biological effects in primary human erythroid progenitors. *Br J Haematol* 2003; **120**: 876–886.



Cell Death and Disease is an open-access journal published by Nature Publishing Group. This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>