

News and Commentary

Heme-albumin: an honorary enzyme

P Ascenzi^{*1}, A di Masi^{1,2}, G Fanali³ and M Fasano^{3,4}*Cell Death and Disease* (2015) 6, e1895; doi:10.1038/cddis.2015.287; published online 8 October 2015

Human serum albumin displays time-dependent heme-based catalytic properties,¹ representing a case for 'chronosteric effects'.² In fact, HSA has a pivotal role in heme transfer from high- and low-density lipoproteins to hemopexin. After endocytosis of the hemopexin-heme complex into the hepatic parenchymal cells through the CD91 receptor, hemopexin releases the heme, which undergoes degradation. Then, hemopexin is released intact into the bloodstream.^{3,4}

The three-domain organization of HSA is at the root of its extraordinary ligand-binding capacity and allosteric control. The most relevant clefts hosting ligands are the so-called fatty acid (FA) binding sites (named FA1 to FA9). Bacterial protein-recognition cleft(s), thyroxine-binding pockets and metal ion-recognition sites also participate to HSA actions.⁴ Noteworthy, the HSA structure and reactivity is affected not only reversibly by pH and ligands (e.g., heme, FAs and drugs), but also irreversibly by chemical modifications, which in turn confer antigenicity properties.⁴

Ferrous human serum heme-albumin (HSA-heme-Fe(II)) binds reversibly to NO and CO. Although the heme-Fe atom of HSA-heme-Fe(II) is rapidly oxidized by O₂, HSA-heme-Fe(II) mutants bearing residues pivotal for O₂ recognition have been proposed not only as red blood cell substitutes, but also as O₂-therapeutic agents.^{1,4,5} Moreover, HSA-heme-Fe(II) catalyzes the nitrite conversion to nitrogen monoxide under acidosis and anaerobic conditions,⁶ HSA-heme-Fe(II)-NO reacts with O₂ and peroxy-nitryte leading to the formation of NO₃⁻,^{7,8} and ferric HSA-heme-Fe (HSA-heme-Fe(III)) catalyzes the conversion of peroxy-nitryte to NO₃⁻,⁹ and displays weak catalase and peroxidase activities.¹⁰

The heme-based catalytic properties of HSA are allosterically modulated by drugs (Figure 1).¹ Domains I and II have a major role in the allosteric modulation of ligand-binding and reactivity properties of HSA, the FA1, FA2, FA6 and FA7 sites being functionally linked. Allosteric modulators (e.g., drugs) of the heme-based catalytic properties of HSA-heme affect the coordination state of the heme-Fe atom. In ligand-free active HSA-heme, the heme-Fe atom displays a four- or five-coordinated heme-Fe atom, whereas inactive HSA-heme shows a six-coordinated heme-Fe atom. Upon drug binding to HSA-heme (most probably to the FA2 site), the re-orientation of the Glu131-Arg145 α -helix and the axial coordination of the heme-Fe atom by His146 and Tyr161 occur; as a consequence, the unreactive six-coordinated HSA-heme species becomes predominant.^{1,11,12}




Figure 1 Human serum albumin displays time-dependent heme-based catalytic properties, which are allosterically modulated by drugs

As a whole, the allosteric modulation of heme-based reactivity properties of HSA-heme by drugs represents a pivotal issue in the pharmacological therapy management, heme-binding switching HSA from a plasmatic carrier to a transient metal-enzyme.

Conflict of Interest

The authors declare no conflict of interest.

1. Ascenzi P *et al.* *Cell Death Discov* 2015; 1: 15025.
2. Ascenzi P *et al.* *IUBMB Life* 2013; 65: 836–844.
3. Miller YI *et al.* *Biochim Biophys Acta* 1999; 1454: 153–164.
4. Fanali G *et al.* *Mol Aspects Med* 2012; 33: 209–290.
5. Tsuchida E *et al.* *Bioconjugate Chem* 2009; 20: 1419–1440.
6. Ascenzi P *et al.* *J Biol Inorg Chem* 2013; 18: 939–946.
7. Ascenzi P *et al.* *Biochem Biophys Res Commun* 2007; 353: 469–474.
8. Ascenzi P *et al.* *Biochem Biophys Res Commun* 2011; 406: 112–116.
9. Ascenzi P *et al.* *J Biol Chem* 2009; 284: 31006–31017.
10. Monzani E *et al.* *Biochim Biophys Acta* 2001; 1547: 302–312.
11. Nicoletti FP *et al.* *J Am Chem Soc* 2008; 130: 11677–11688.
12. Meneghini C *et al.* *PLoS One* 2014; 9: e104231.



Cell Death and Disease is an open-access journal published by Nature Publishing Group. This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

¹Interdepartmental Laboratory for Electron Microscopy, Roma Tre University, Roma I-00146, Italy; ²Department of Sciences, Roma Tre University, Roma I-00146, Italy; ³Department of Theoretical and Applied Sciences, Biomedical Research Division, University of Insubria, Busto Arsizio I-21052, Italy and ⁴Center of Neuroscience, University of Insubria, Busto Arsizio I-21052, Italy

*Corresponding author: P Ascenzi, Interdepartmental Laboratory for Electron Microscopy, Roma Tre University, Viale G Marconi 446, Roma I-00146, Italy. Tel: +39 339 295 5115; Fax: +39 06 5733 6321; E-mail: ascenzi@uniroma3.it