

## Review

# Long noncoding RNAs in the progression, metastasis, and prognosis of osteosarcoma

Zuozhang Yang<sup>\*1,2</sup>, Xiaojuan Li<sup>1,2</sup>, Yihao Yang<sup>1,2</sup>, Zewei He<sup>1</sup>, Xin Qu<sup>1</sup> and Ya Zhang<sup>1</sup>

Long noncoding RNAs (lncRNAs) are a class of non-protein-coding molecules longer than 200 nucleotides that are involved in the development and progression of many types of tumors. Numerous lncRNAs regulate cell proliferation, metastasis, and chemotherapeutic drug resistance. Osteosarcoma is one of the main bone tumor subtypes that poses a serious threat to adolescent health. We summarized how lncRNAs regulate osteosarcoma progression, invasion, and drug resistance, as well as how lncRNAs can function as biomarkers or independent prognostic indicators with respect to osteosarcoma therapy.

*Cell Death and Disease* (2016) 7, e2389; doi:10.1038/cddis.2016.272; published online 29 September 2016

## Facts

- Long noncoding RNAs (lncRNAs) regulate cell proliferation in osteosarcoma.
- lncRNAs regulate cell invasion and chemotherapeutic drug resistance in human osteosarcoma.
- lncRNAs function as biomarkers and independent prognostic indicators with respect to osteosarcoma therapy.

## Open Questions

- How do lncRNAs regulate osteosarcoma progression and invasion?
- How do lncRNAs regulate osteosarcoma chemotherapeutic drug resistance?
- Can lncRNAs be used as biomarkers or prognostic indicators with respect to human osteosarcoma treatment?

Osteosarcoma is the most common malignant bone tumor in children and adolescents. It is a genetically unstable and highly malignant mesenchymal tumor of bone characterized by structural chromosomal alterations.<sup>1,2</sup> Malignant osteosarcoma cells produce osteoid matrix and fibrillary stroma.<sup>3</sup> The most common osteosarcoma subtypes are osteoblastic osteosarcoma, chondroblastic osteosarcoma, and fibroblastic sarcoma.<sup>4,5</sup> Osteosarcoma occurs predominantly in adolescents and young adults and accounts for ~5% of childhood cancers. Most osteosarcoma patients are diagnosed under

the age of 25 years, and the disease occurs more often in males than in females.<sup>6</sup> Osteosarcoma often exhibits locally invasive growth. Pulmonary metastases are often seen in patients with aggressive tumors. Both biopsy findings and classic X-ray findings contribute to the diagnosis of osteosarcoma and yield important information that can be used to select appropriate therapies.<sup>7,8</sup> In most osteosarcoma patients, chemotherapy and/or radiation therapy are usually administered before or after surgery to prevent tumors from spreading throughout the body. However, patients with distant metastases still fare poorly, as the 5-year survival rate in these patients is ~20%.<sup>9,10</sup> Thus, developing comprehensive and multidimensional treatments for osteosarcoma is necessary, and gene therapies using viral vectors, immune therapies, antiangiogenic therapies, and proapoptotic therapies have been investigated regarding their application in patients with osteosarcoma.

To date, the molecular mechanism underlying osteosarcoma development remains unclear. The majority of previous studies have focused on protein-coding genes as crucial components involved in the progression and metastasis of osteosarcoma and have overlooked the vast landscape of noncoding genes.

Since the invention of DNA sequencing methods and the completion of the draft human genome sequence, researchers have found that only 1.5% of 3.2 billion nucleotide pairs code for proteins and that the other 98.5% of DNA sequences do not code for proteins. These sequences are recognized as junk

<sup>1</sup>Bone and Soft Tissue Tumors Research Center of Yunnan Province, Department of Orthopaedics, The Third Affiliated Hospital of Kunming Medical University (Tumor Hospital of Yunnan Province), Kunming 650118, China

\*Corresponding author: Z Yang, Department of Orthopaedics, The Third Affiliated Hospital of Kunming Medical University (Tumor Hospital of Yunnan Province), Kunming 650118, China. Tel: +86 871 68221157; Fax: +86 871 68221157; E-mail: yangzuozhang@163.com

<sup>2</sup>These authors contributed equally to this work and should be considered as co-first authors.

**Abbreviations:** siRNA, small interfering RNA; microRNA, micro RNA; lncRNA, long noncoding RNA; Hh signaling pathway, hedgehog signaling pathway; BMP, bone morphogenetic protein; PI3K, phosphatidylinositol 3-kinase; PACER, P50-associated COX-2 extragenic RNA; MALAT1, metastasis-associated lung adenocarcinoma transcript 1; MMP-9, matrix metalloproteinase 9; Yap1, yes-associated protein 1; ECM, extracellular matrix; TIMP, inhibitors of metalloproteinase; HOTAIR, HOX antisense intergenic RNA; SNHG12, small nucleolar RNA host gene 12; AMOT, angiomin; FGFR3-AS1, FGFR3 antisense transcript 1; HIF2PUT, hypoxia-inducible factor-2 $\alpha$  (HIF2 $\alpha$ ) promoter upstream transcript; TUSC7, tumor suppressor candidate 7


Received 10.6.16; revised 27.7.16; accepted 01.8.16; Edited by E Candi

sequences that have accumulated because of the process of evolution.<sup>11,12</sup> ENCODE (Encyclopedia of DNA Elements) projects postulate that 80% of genome sequences are transcribed into primary transcripts and have biochemical functions.<sup>13,14</sup> The concept of 'junk DNA' has rapidly attracted the attention of researchers. According to their biological functions, noncoding RNAs can be divided into housekeeping noncoding RNAs and regulatory noncoding RNAs.<sup>15–17</sup> Housekeeping noncoding RNAs comprise ribosomal RNAs,<sup>18–20</sup> transfer RNAs,<sup>21,22</sup> small nuclear RNAs,<sup>23–25</sup> small nucleolar RNAs,<sup>26–28</sup> guide RNAs,<sup>29–32</sup> and telomerase RNAs.<sup>33,34</sup> Regulatory noncoding RNAs comprise small interfering RNAs (siRNAs),<sup>35–37</sup> micro RNAs (microRNA),<sup>38–40</sup> piwi-interacting RNAs,<sup>41–43</sup> and long noncoding RNAs (lncRNAs).<sup>44,45</sup>

lncRNAs are a large and diverse class of non-protein-coding transcripts longer than 200 nucleotides.<sup>46,47</sup> lncRNAs have recently gained widespread attention and have been shown to have crucial roles in various biological regulatory processes. lncRNA sequences are conserved, and lncRNA expression profiles in adult tissues are broad. lncRNA messenger RNAs (mRNAs) are generally less abundant than protein-coding mRNAs but exhibit stronger tissue- and cell-specific lncRNA expression patterns.<sup>48,49</sup> Most lncRNAs are transcribed by RNA polymerase II enzymes that lack open-reading frames and are expressed in specific tissues and/or during specific developmental stages, demonstrating that the genes encoding these molecules are strictly regulated with respect to tissue development. Previous research has mainly focused on microRNAs and siRNAs. lncRNAs have recently been found to be involved in development, differentiation, and proliferation, as well as cell cycle regulation and programmed cell death.<sup>50–53</sup> They also have important roles in the progression and metastasis of various tumors, such as colon cancer, liver cancer, breast cancer, bladder cancer, and cervical cancer<sup>51,54–59</sup> (Figure 1). In this paper, we reviewed the biological functions of lncRNAs and the molecular mechanisms underlying these functions with respect to osteosarcoma progression. Chemotherapy drug resistance remains an obstacle affecting osteosarcoma treatment. We therefore also summarized the lncRNAs that are correlated with chemotherapy drug resistance in osteosarcoma therapy. Furthermore, we summarized several lncRNAs that can function as independent prognostic indicators of overall survival and can serve as useful biomarkers of osteosarcoma progression and prognosis. An overview of the lncRNAs that are associated with osteosarcoma is shown in Table 1.

### lncRNA Regulates Signaling Pathways in Osteosarcoma

Developing effective and targeted therapies for osteosarcoma is dependent on gaining an improved understanding of the molecular mechanisms underlying osteosarcomagenesis, proliferation, invasion, and metastasis.<sup>60</sup> To date, the molecular mechanism underlying osteosarcoma development has not been elucidated. It is known that Wnt signaling is involved in osteosarcoma development, metastasis, and drug resistance. For example, inhibiting Wnt signaling by targeting c-Met, a Wnt-regulated proto-oncogene, was shown to be useful for treating osteosarcoma, suggesting that the Wnt



**Figure 1** Biological processes are regulated by lncRNAs, and several regulatory mechanisms are shown

signaling pathway is involved in osteosarcoma development and metastasis.<sup>61</sup> Chemotherapeutic drug resistance represents a major obstacle with respect to osteosarcoma treatment, due in part to phenotypic cell transitions toward stem-like phenotypes caused by exposure to conventional chemotherapeutics.<sup>62,63</sup> However, the combination of a Wnt/ $\beta$ -catenin signaling pathway inhibitor and doxorubicin prevented the upregulation of factors linked to these types of transitions and was thus envisaged as a means of overcoming adaptive resistance.<sup>64</sup> Aberrant hedgehog (Hh) signaling pathway activity has been observed in osteosarcoma cell lines, as well as in primary human osteosarcoma tissue specimens, and exerts promigratory effects leading to the development of osteoblastic osteosarcoma.<sup>65</sup> Other studies have also demonstrated that dysregulated Hh signaling contributes to poor clinical outcomes in osteosarcoma therapy.<sup>66,67</sup> Bone morphogenetic protein (BMP) signaling pathways have been reported to induce mesenchymal stem cell osteogenic commitments and terminal differentiation, which is initiated by BMP ligand heterodimer (BMPRI and II) binding and signal transduction through the Smad pathway, as well as mitogen-activated protein kinase (MAPK) phosphorylation.<sup>68–71</sup> In particular, of the 31 different types of known BMP ligands, BMP-2, -4, -6, -7, and -9 have significant roles in osteogenesis induction in osteosarcoma.<sup>72–74</sup> Moreover, the Notch pathway has been described as an oncogene that is involved in osteosarcoma proliferation, migration, invasiveness, and oxidative stress resistance, as well as the expression of markers associated with stemness or tumor-initiating cells.<sup>75–78</sup> Moreover, this pathway has a vital role in regulating tumor angiogenesis and vasculogenesis in osteosarcoma.<sup>79</sup> The phosphatidylinositol 3-kinase (PI3K)/Akt pathway is also

Table 1 Overview of lncRNA in osteosarcoma

Name	Functions	Up/downregulation	Regulation mechanism	Reference
H19	Oncogene	+	lncRNA H19 was one of the candidates with the highest relevance in cancer development H19 is induced by Yap1 overexpression in the mouse osteosarcoma tissues and primary osteosarcoma cells.	98
MALAT1	Oncogene	+	MALAT1 might be an oncogenic lncRNA that promoted proliferation and metastasis of osteosarcoma by PI3K/AKT signaling pathway in osteosarcoma cell lines and clinical tumor samples Knockdown of the putative Myc-6 target MALAT1 significantly impaired cell growth of MG63 cells, suggesting that Myc-6 might exert its tumor-suppressive ability in part through the specific downregulation of MALAT1.	94,95,97
ANCR	Oncogene	+	ANCR worked as an oncogenic lncRNA that promoted proliferation of osteosarcoma by regulating the cell cycle progression The cell cycle of U2OS cells was arrested at the G0/G1 phase and the expression level of p21 was increased and CDK2 was decreased in ANCR knockdown cells..	108
HOTAIR	Oncogene	+	Overexpression of HOTAIR promoted tumor growth and metastasis by increasing the secretion of MMP-2 and MMP-9 in human osteosarcoma	101
PACER	Oncogene	+	High levels of HOTAIR were correlated with advanced tumor stage and poor prognosis. PACER was overexpressed in clinical osteosarcoma tissues and cell lines. Its overexpression promoted proliferation and metastasis of osteosarcoma cells by activating COX-2 gene in an NF- $\kappa$ B-dependent manner.	92
SNHG12	Oncogene	+	SNHG12 expression was upregulated in tissue samples from primary osteosarcoma and adjacent normal tissues, the osteosarcoma cell lines, SAOS-2, MG63, U-2 OS, and the human osteoblast cell line hFOB (OB3), by upregulating AMOT gene expression.	102
FGFR3-AS1	Oncogene	+	The expression of FGFR3-AS1 and FGFR3 was positively correlated in osteosarcoma tissues FGFR3-AS1 promoted osteosarcoma growth through regulating its natural antisense transcript FGFR3.	104
ENST00000563280, NR-036444	Oncogene	+	The lncRNA had a critical role in doxorubicin resistance of OS by interacting with important genes such as ABCB1, HIF1A, and FOXO2.	117
ENST00000563280	Oncogene	+	The lncRNA was significantly increased in specimens of OS patients with a poor chemoresponse compared with those with a good chemoresponse The patients with lower expression of ENST00000563280 may survived longer compared with those of higher expression.	117
TUG1	Oncogene	+	TUG1 was overexpressed in patients with osteosarcoma TUG1 was positively correlated with disease status It may serve as a molecular indicator in maintaining surveillance and forecasting prognosis.	114
HOTTIP	Oncogene	+	HOTTIP was overexpressed in OS tissues HOTTIP was correlated with advanced clinical stage and distant metastasis OS patients with high HOTTIP expression level had poorer overall survival compared with those with low HOTTIP expression	115
MEG3	Tumor suppressor	-	It could be a novel prognostic marker and potential therapeutic target in OS patients. Liu et al. found that the expression of lncRNA MEG3 was associated with clinical stage and distant metastasis	116
HIF2PUT	Suppressor gene	-	The patients with low lncRNA MEG3 expression had a shorter overall survival MEG3 was an independent predictor to overall survival of osteosarcoma patients.	110
TUSC7	Tumor suppressor	-	HIF2PUT is a novel regulatory factor of osteosarcoma stem cells HIF2PUT exerted its function partly by controlling HIF2 $\alpha$ expression. TUSC7 was significantly downregulated in osteosarcoma tissues TUSC7 regulated the cell proliferation ability and colony formation ability involved in regulating the expression of proapoptotic proteins.	111

Abbreviations: AMOT, angiominin; ANCR, anticifferentiation noncoding RNA; FGFR3-AS1, FGFR3 antisense transcript 1; HIF2 $\alpha$ , hypoxia-inducible factor-2 $\alpha$ ; HIF2PUT, HIF2 $\alpha$  promoter upstream transcript; HOTAIR, HOX antisense intergenic RNA; HOTTIP, HOXA transcript at the distal tip; lncRNA, long noncoding RNA; MALAT1, metastasis-associated lung adenocarcinoma transcript 1; MEG3, maternally expressed gene 3; MMP, matrix metalloproteinase; NF- $\kappa$ B, nuclear factor- $\kappa$ B; PACER, P50-associated COX-2 extragenic RNA; SNHG12, small nucleolar RNA host gene 12; TUG1, taurine-upregulated gene 1; TUSC7, tumor suppressor candidate 7; yap1, yes-associated protein 1

thought to be one of the most important oncogenic pathways in human osteosarcoma.<sup>80,81</sup> A large number of regulatory factors regulate osteosarcoma cell proliferation, apoptosis, angiogenesis, metastasis, and chemotherapy drug sensitivity by regulating PI3K/Akt signaling, including p53,<sup>82,83</sup> VEGF,<sup>84</sup> CXCR7,<sup>85</sup> Aurora-B,<sup>86</sup> microRNA-221,<sup>87</sup> cyclooxygenase-2,<sup>88</sup> BYL719, a PI3K inhibitor,<sup>89</sup> and LY294002.<sup>90</sup> All these signaling pathways are interconnected to regulate osteosarcoma progression and migration.

To date, few studies have reported the roles of lncRNAs in osteosarcoma osteogenesis, development, invasion, metastasis or chemotherapy resistance. Alterations in the expression of several lncRNAs have been observed in osteosarcoma. Li *et al.*<sup>91</sup> detected the expression profiles of numerous lncRNAs via microarray analysis and observed several differentially expressed lncRNAs in osteosarcoma tissues compared with paired adjacent noncancerous tissues. In particular, 25 733 lncRNAs were expressed in osteosarcoma, including 403 consistently over-regulated lncRNAs involved in 34 pathways and 798 consistently under-regulated lncRNAs involved in 32 pathways, across all samples (2.0-fold,  $P < 0.05$ ), suggesting that lncRNAs can function as therapeutic targets and serve as novel candidate biomarkers with respect to osteosarcoma diagnosis and prognosis.<sup>91</sup> P50-associated COX-2 extragenic RNA (PACER) was overexpressed in clinical osteosarcoma tissues and cell lines influenced by DNA methylation, activated the *COX-2* gene in an NF- $\kappa$ B-dependent manner and functioned as an oncogene in osteosarcoma.<sup>92</sup> Metastasis-associated lung adenocarcinoma transcript 1 (MALAT1), one of the first cancer-associated lncRNAs to be identified, is expressed in numerous tissues, is highly abundant in neurons and is involved in regulating the recruitment of SR family pre-mRNA-splicing factors to sites of transcription involved not only in nuclear processes but also in synapse function.<sup>93</sup> Aberrant MALAT1 expression has been observed in many types of tumors, including hepatocellular carcinoma, cervical cancer, breast cancer, ovarian cancer, and colorectal cancer. Dong *et al.*<sup>94</sup> found that MALAT1 was highly expressed in human osteosarcoma tissues and that its expression level was closely correlated with pulmonary metastasis. Moreover, they found that MALAT1 knockdown suppressed human osteosarcoma cell proliferation, invasion, and metastasis *in vitro* and *in vivo*. They also explored the molecular mechanisms underlying the function of MALAT1 in osteosarcoma and observed that MALAT1 inhibited tumor growth and metastasis via the PI3K/AKT signaling pathway, as the expression levels of proliferating cell nuclear antigen, matrix metalloproteinase 9 (MMP-9), phosphorylated PI3Kp85 $\alpha$ , and Akt were significantly decreased in MALAT1-knockdown cells.<sup>94</sup> Cai *et al.*<sup>95</sup> observed similar results in MALAT1 siRNA-treated osteosarcoma cells. They showed that MALAT1 knockdown inhibited osteosarcoma cell proliferation and migration, induced osteosarcoma cell cycle arrest and cell apoptosis, and delayed tumor growth in an osteosarcoma xenograft model. Specifically, they found that MALAT1 siRNA administration decreased the protein expression levels of RhoA and its downstream effectors, the Rho-associated coiled-coil containing protein kinases (ROCKs). Consistent with these studies, high-dose 17 $\beta$ -estradiol (E2) treatment markedly


downregulated MALAT1-mediated osteosarcoma cell proliferation, migration, invasion, and metastasis by upregulating miR-9 in E2-dose-dependent and ER-independent manners. In addition, MALAT1 downregulation promoted the formation of the SFPQ/PTBP2 complex.<sup>96</sup> Moreover, Taniguchi *et al.*<sup>97</sup> found that MALAT1 contains a theoretical Myc-6-target sequence that includes an E-box-like motif (at positions -258 to -251). Interestingly, knockdown of the putative Myc-6 target MALAT1 obviously impaired MG63 cell growth. In general, Myc-6 appears to exert its tumor-suppressive effects, at least in part, through the specific downregulation of MALAT1. The Hh signaling pathway has important roles in vertebrate embryonic development and growth regulation, functions as a morphogen and mitogen, and is normally deactivated after embryogenesis. However, Hh signaling is reactivated and upregulated in various cancers, including osteosarcoma, resulting in high levels of yes-associated protein 1 (Yap1) expression. Yap1, a potent oncogene expressed in both human and mouse tumor tissues, is amplified in various cancers. Hh signaling inhibition reduces Yap1 expression, and Yap1 knockdown significantly inhibits tumor progression. Chan *et al.*<sup>98</sup> found that aberrant Hh signaling in mature osteoblasts is responsible for the pathogenesis of osteoblastic osteosarcoma. Moreover, Hh signaling upregulation and Yap1 overexpression lead to aberrant lncRNA H19 expression in malignant osteosarcoma. The lncRNAs involved in osteosarcoma and osteosarcoma-related signaling pathways are shown in Figure 2.

### lncRNA Regulates Osteosarcoma Metastasis


Distant metastases are commonly observed in patients with osteosarcoma after surgery. It is estimated that metastases have been found in 85% of patients with osteosarcoma. The most common site of osteosarcoma metastasis is the lung. Metastatic osteosarcoma is difficult to control, and respiratory symptoms appear only in the setting of extensive involvement. Osteosarcoma also metastasizes to other bone and soft tissue locations. This issue is still controversial, as some authors have argued that bone metastases may actually be multifocal osteosarcomas rather than actual metastases. Death from osteosarcoma is usually a result of pulmonary metastasis and respiratory failure because of widespread progression.

Tumor invasion and metastasis is a multistep, multilink, complex process comprising invasion, intravasation, dissemination, extravasation, and colonization. Briefly, tumor cells alter cell-extracellular matrix (ECM) interactions at the primary tumor site, escape from the primary site and invade adjacent tissues, and translocate through the vasculature to migrate to other systems. Then, these metastatic cancer cells anchor to distant vessel walls and extravasate into their destination tissues (Figure 3) before finally proliferating from microscopic growths to form secondary tumors.

Adhesion molecules, angiogenic factors, proteolytic enzymes, tumor metastasis-related factors, and metastasis suppressors are involved in migration and metastasis. MMPs are a family of proteolytic enzymes and are the key proteases involved in digesting components of the ECM and surface receptors. MMPs has an important role in tumor invasion and metastasis by degrading the ECM and basement membrane



**Figure 2** Osteosarcoma cell proliferation is regulated by lncRNAs, including H19, MALAT1, ANCR, and PACER. These osteosarcoma-related lncRNAs are involved in the PI3K/Akt signaling pathway, NF-κB signaling pathway, and Hh/Yap1 signaling pathway




**Figure 3** Tumor invasion and metastasis is a multistep, multistage complex process. Tumor cells at primary tumor sites invade surrounding tissues, migrate through the blood or lymph and localize in distal targeted tissues. This process is divided into the following five stages: invasion, intravasation, dissemination, extravasation, and colonization

to remodel the tumor microenvironment and promote tumor angiogenesis. Conversely, MMP activity is suppressed by endogenous tissue inhibitors of metalloproteinases (TIMPs), specific MMP inhibitors. The levels of endogenous MMPs and TIMPs contribute to imbalances between MMPs and TIMPs and regulate ECM degradation and deposition. It has been

reported that the levels of MMP-2 and MMP-9 secretion are elevated in several types of human cancers and that these elevations are associated with a poor prognosis.<sup>99</sup> During osteosarcoma cell invasion and migration, several lncRNAs reportedly promote or inhibit cell proliferation and invasion by regulating MMP-2 and MMP-9 secretion.<sup>100</sup> Osteosarcoma cell invasion and metastasis and the lncRNAs associated with these processes are shown in Figure 4.

The HOX antisense intergenic RNA (HOTAIR), a well-known lncRNA, is involved in the pathogenesis and progression of multiple tumors. HOTAIR is commonly overexpressed in osteosarcoma, and its knockdown significantly inhibits cellular proliferation and invasion by decreasing MMP-2 and MMP-9 secretion in osteosarcoma cells. Meanwhile, high HOTAIR expression levels are significantly associated with advanced tumor stages, high histological grades, and poor prognoses. Thus, HOTAIR may be an important target in the treatment of human osteosarcoma.<sup>101</sup> It has been reported that the small nucleolar RNA host gene 12 (SNHG12) promotes cell proliferation and migration by upregulating angiomin (AMOT) gene expression in human osteosarcoma cells.<sup>102</sup> In particular, tissue samples from primary osteosarcomas ( $n=20$ ) and adjacent normal tissues ( $n=20$ ), as well as samples from the osteosarcoma cell lines SAOS-2, MG63, and U2OS and the human osteoblast cell line hFOB (OB3), were studied using quantitative real-time polymerase chain reaction to detect SNHG12 expression. They found that SNHG12 mRNA expression was upregulated in osteosarcoma tissues and cell lines compared with normal tissues and cells and that SNHG12 knockdown suppressed cell proliferation and migration but did not affect cell apoptosis. These findings suggest that SNHG12 lncRNA promotes cell



**Figure 4** Osteosarcoma invasion and metastasis is regulated by lncRNAs, such as MALAT1, SNHG12, HOTAIR, FGFR3-AS1, and HIF2PUT. MMP-2 and MMP-9 secretion is regulated by the Erk1/2, JNK1/2, P38, PI3K/Akt, and NF- $\kappa$ B signaling pathways. Osteosarcoma cell invasion is regulated by MMP-2 and MMP-9

proliferation and migration by upregulating AMOT gene expression in osteosarcoma cells *in vivo* and *in vitro* and are consistent with the findings of previous studies involving human gastric cancer patients, which showed that upregulation of SNHG15 lncRNA expression promotes cell proliferation and invasion by regulating MMP-2/MMP-9 expression.<sup>103</sup> Mammalian genomes encode numerous natural antisense transcripts that are at least partially complementary to their sense transcripts. FGFR3 antisense transcript 1 (FGFR3-AS1) increased FGFR3 mRNA stability and upregulated FGFR3 expression via antisense pairing with FGFR3 3'-UTR. Increased FGFR3-AS1 expression was correlated with large tumor size, advanced Enneking stage, metastasis and poor survival. FGFR3-AS1 knockdown inhibited xenograft tumor growth of osteosarcoma cells *in vitro* and *in vivo*. Therefore, lncRNA FGFR3-AS1 promoted osteosarcoma growth by regulating its natural antisense transcript FGFR3.<sup>104</sup>

#### lncRNA and Osteosarcoma Cell Proliferation

Cancer occurrence is characterized by uncontrolled cell cycle activity, including uncontrolled DNA replication and parental cell division.<sup>105</sup> Imbalances between programmed cell death and cell proliferation contribute to the development of various cancers. Both oncogene activation and tumor suppressor gene inactivation lead to cancer occurrence and development.<sup>106,107</sup> In osteosarcoma, lncRNAs also exhibit oncogenic

properties or act as tumor suppressors to control osteosarcoma progression by regulating cell cycle progression or cell apoptosis to regulate cell proliferation or migration. Antidifferentiation noncoding RNA (ANCR) is a newly identified oncogenic lncRNA that has an important role in the maintenance of cell undifferentiation. ANCR knockdown significantly inhibited U2OS and SAOS cell proliferation and U2OS cell colony formation and arrested the U2OS cell cycle at the G0/G1 phase. Moreover, ANCR regulated and controlled cell cycles by regulating the endogenous levels of cell cycle-related proteins, including p21, CDK2, and CDK4.<sup>108</sup> The levels of taurine-upregulated gene 1 (TUG1) and one of its transcript variants (n377360) were significantly higher in osteosarcoma tissues compared with that in matched non-tumorous tissues. Consistent with this finding, TUG1 and n377360 suppression by siRNA significantly impaired osteosarcoma cell proliferation potential and promoted osteosarcoma cell apoptosis.<sup>109</sup>

Tumor suppressor lncRNAs are involved in regulating human osteosarcoma. The levels of hypoxia-inducible factor-2 $\alpha$  (HIF2 $\alpha$ ) promoter upstream transcript (HIF2PUT), a novel lncRNA, were assessed via quantitative polymerase chain reaction in 17 osteosarcoma tissue specimens, and the data demonstrated that HIF2PUT functions as an osteosarcoma stem cell inhibitor *in vitro* partly by controlling HIF2 $\alpha$  expression. HIF2PUT overexpression markedly inhibited cell proliferation and migration, decreased the percentage of CD133-expressing cells, and impaired the osteosarcoma stem sphere-forming ability of MG63 cells.<sup>110</sup> It has been

reported that the HIF2PUT expression levels were positively correlated with HIF2 $\alpha$  expression in osteosarcoma tissues. However, HIF2PUT overexpression obviously suppressed cell proliferation and migration, decreased the percentage of CD133-expressing cells, and impaired the osteosarcoma stem sphere-forming ability of MG63 cells. However, HIF2PUT knockdown had the opposite effect. Tumor suppressor candidate 7 (TUSC7) is a potential tumor suppressor that has been shown to inhibit cell proliferation in osteosarcoma. Cong *et al.*<sup>111</sup> reported that TUSC7 expression was significantly downregulated in osteosarcoma tissues compared with paired non-tumor tissues. Low TUSC7 expression is associated with poor survival (HR = 0.313, 95% confidence interval (CI) 0.092–0.867) in osteosarcoma patients. Loss of TUSC7 copy number is also associated with a poor prognosis (HR = 3.994, 95% CI: 1.147–13.91) in osteosarcoma patients. The author of the above study used two osteosarcoma cell lines, HOS and MG63, to investigate the biological function of TUSC7. Silencing TUSC7 increased osteosarcoma cell proliferation ability and colony formation ability. The cell cycle was not affected by TUSC7 silencing; however, the percentage of apoptotic cells decreased, and the expression levels of several proapoptotic proteins were downregulated. Importantly, xenograft tumor models were established in nude mice using MG63 cells. Silencing TUSC7 significantly promoted tumor growth *in vivo* in treated mice compared with negative-control mice. Thus, TUSC7 may be a tumor suppressor in osteosarcoma. Similarly, Wang *et al.*<sup>112</sup> determined that TUSC7 is a potential biomarker for NSCLC prognosis and that TUSC7 dysregulation has an important role in NSCLC progression. In their studies, they found that the expression levels of TUSC7 were lower in NSCLC tissues and lung cancer cells compared with that in normal tissues and cells. Lower TUSC7 expression levels in NSCLC tissues were associated with larger tumor sizes and higher TNM stages. Patients with lower TUSC7 expression levels exhibited worse overall survival compared with patients with high TUSC7 expression levels. Univariate and multivariate analyses suggested that low TUSC7 expression was an independent prognostic indicator of a poor prognosis in NSCLC patients. Moreover, TUSC7 upregulation inhibited lung cancer cell proliferation *in vitro*.

### LncRNA and Osteosarcoma Prognosis

Genetic variants of HOTAIR lncRNA contribute to the risk of osteosarcoma. A two-stage, case–control study involving 900 OS patients and 900 controls was performed to evaluate the associations between HOTAIR lncRNA genetic variants and OS risk in the Chinese population, the results of which demonstrated that the C allele of rs7958904 was associated with a significantly decreased OS risk compared with the G allele (OR: 0.77; 95% CI: 0.67–0.90;  $P = 6.77 \times 10^{-4}$ ), suggesting that patients with the rs7958904 CC genotype had significantly lower HOTAIR RNA levels compared with patients with other genotypes, as well as a lower OS risk.<sup>113</sup> Ma *et al.*<sup>114</sup> found that TUG1 was significantly overexpressed in osteosarcoma tissues compared with matched adjacent normal tissues ( $P < 0.01$ ). Moreover, TUG1 levels were strongly correlated with disease status and tumor size,

postoperative chemotherapy, and Enneking surgical stage. Furthermore, TUG1 upregulation was strongly correlated with a poor prognosis and was an independent prognostic indicator for overall survival (HR = 2.78; 95% CI: 1.29–6.00;  $P = 0.009$ ) and progression-free survival (HR = 1.81; 95% CI = 1.01–3.54;  $P = 0.037$ ). HOTTIP was overexpressed in OS tissues and was correlated with advanced clinical stage and distant metastasis. High HOTTIP expression levels were associated with poor overall survival in OS patients. Moreover, HOTTIP expression was an independent prognostic factor for overall survival in OS patients and may represent a novel prognostic marker and therapeutic target in OS patients.<sup>115</sup> Liu *et al.*<sup>116</sup> demonstrated that MEG3 lncRNA levels were clearly lower in osteosarcoma tissues compared with that in adjacent non-tumor tissues. Patients with low MEG3 lncRNA expression levels exhibited shorter overall survival compared with patients with high expression levels (log-rank test,  $P < 0.05$ ). Furthermore, decreased MEG3 lncRNA expression, advanced clinical stage, and distant metastasis were all independent predictors of shorter overall survival in osteosarcoma patients.

### LncRNA and Chemotherapeutic Drug Resistance in Osteosarcoma

Surgery, radiotherapy, and chemotherapy are the three main treatments for cancer. In particular, chemotherapy has an important role in cancer therapy. However, chemotherapeutic drug resistance is the largest obstacle limiting the success of cancer therapy. Large numbers of studies have focused on chemotherapy drug resistance in human osteosarcoma, but the mechanism underlying this resistance remains to be elucidated. In osteosarcoma, chemotherapy drug efficacy is usually limited by acquired resistance to specific drugs, such as doxorubicin and cisplatin. Zhu *et al.*<sup>117</sup> studied three sets of doxorubicin-resistant MG63/DXR cells and their paired parental MG63 cells and identified 3465 lncRNAs (1761 up and 1704 down) and 3278 mRNAs (1607 up and 1671 down) that were aberrantly expressed in MG63/DXR cells (fold change  $> 2.0$ ,  $P < 0.05$  and FDR  $< 0.05$ ). Moreover, an lncRNA-mRNA coexpression network identified lncRNAs, including ENST00000563280 and NR-036444, that interact with genes such as *ABCB1*, *HIF1A*, and *FOXO2* and may have an important role in doxorubicin resistance in OS. Several lncRNAs have been found to serve as biomarkers predicting the chemoresponses and prognoses of osteosarcoma patients, including ENST00000563280, whose expression level was significantly increased in the tissue specimens of OS patients with poor chemoresponses compared with those with good chemoresponses.

### Conclusions and Perspectives

Previous studies have reported that lncRNAs regulate the transcription, stability and translation of protein-coding genes in the mammalian genome, play important roles in regulating protein-coding genes at the transcriptional and post-transcriptional levels, and participate in important biological processes, including cell differentiation, development and human diseases.<sup>118–120</sup> Human genome studies have shown

that ~18% of protein-coding genes that produce lncRNAs (10/57) are related to cancer, whereas only 9% of all human protein-coding genes (2147/23621) are related to cancer (chi-square test,  $P$ -value: 0.047; hypergeometric probability  $P$ -value = 0.018), clearly demonstrating that genes implicated in cancer development have a greater tendency to produce lncRNAs.<sup>121</sup> In this review, we have summarized how lncRNAs regulate cell proliferation, invasion and chemotherapeutic drug resistance in human osteosarcoma patients and osteosarcoma cells. We have also summarized the roles of lncRNAs as prognostic biomarkers in osteosarcoma therapy. Finding promising therapeutic targets for the treatment for human osteosarcoma, especially chemotherapeutic drug-resistant osteosarcoma, will be beneficial for patients. However, several questions regarding the involvement of lncRNAs in osteosarcoma remain unexplored and unresolved.

- (1) To date, limited studies regarding the involvement of lncRNAs in human osteosarcoma have been published. Although several lncRNAs are known to exert tumor-promoting or tumor-suppressing effects in osteosarcoma species and cancer cell lines, the exact molecular mechanisms underlying these effects remain unclear. Thus, additional investigations are required to elucidate the molecular mechanisms underlying human osteosarcoma progression, metastasis and drug resistance.
- (2) One lncRNA may be involved in several different signaling pathways associated with cancer development and may have more than one target associated with osteosarcoma proliferation and metastasis. For instance, MALAT1 plays an important role in the PI3K/AKT and RhoA/ROCKs signaling pathways. However, understanding the connections between these signaling pathways, as well as determining whether one of them plays a major role in osteosarcoma development and progression, warrants further study.
- (3) lncRNAs usually have short half-lives and exhibit low transcript abundance. They need to be transiently expressed *in vitro*. Additionally, it is necessary to determine how the secondary and tertiary structures of lncRNAs interact with specific protein targets.
- (4) lncRNAs may represent novel therapeutic targets, which are critical for developing novel strategies for the early diagnosis and treatment of human osteosarcoma. The potential clinical applications of miRNAs warrant investigation.

### Conflict of Interest

The authors declare no conflict of interest.

**Acknowledgements.** This research was supported, in part, by grants from the National Natural Science Foundation of China (No.81460440 and 81372322), the Joint Special Funds for the Department of Science and Technology of Yunnan Province-Kunming Medical University (No. 2014FB059), the Scientific Research Projects from Internal Research Institutions of Medical and Health Units in Yunnan Province (No. 2014NS013, 2014NS014, 2014NS015, and 2014NS016), Foundation of the Yunnan Provincial Innovative Team of Bone and Soft Tissue Tumor (No.2015HC026), Foundation of the Young and Middle-aged Academic and Technical Leaders of Yunnan Province (No.2014HB034).

1. He H, Ni J, Huang J. Molecular mechanisms of chemoresistance in osteosarcoma [review]. *Oncol Lett* 2014; **7**: 1352–1362.
2. Liu JJ, Liu S, Wang JG, Zhu W, Hua YQ, Sun W et al. Telangiectatic osteosarcoma: a review of literature. *Oncotargets Ther* 2013; **6**: 593–602.
3. Tsang KY, Blakemore WS. Immunologic studies in contacts of osteosarcoma in humans and animals. *Nature* 1977; **265**: 541–542.
4. Williams SA, Maecker HL, French DM, Liu J, Gregg A, Silverstein LB et al. USP1 deubiquitinates ID proteins to preserve a mesenchymal stem cell program in osteosarcoma. *Cell* 2011; **146**: 918–930.
5. Cutilli T, Scarsella S, Fabio DD, Oliva A, Cargini P. High-grade chondroblastic and fibroblastic osteosarcoma of the upper jaw. *Ann Maxillofac Surg* 2011; **1**: 176–180.
6. Fu HL, Shao L, Wang Q, Jia T, Li M, Yang DP. A systematic review of p53 as a biomarker of survival in patients with osteosarcoma. *Tumour Biol* 2013; **34**: 3817–3821.
7. Kundu ZS. Classification, imaging, biopsy and staging of osteosarcoma. *Indian J Orthop* 2014; **48**: 238–246.
8. Conde DM, Morais LC, Pacheco CF, Ferreira RB, Sousa ESEP, Nunes AR et al. Primary osteosarcoma of the breast: pathological and imaging findings. *Rev Assoc Med Brasil* 2015; **61**: 497–499.
9. Guo J, Reddick WE, Glass JO, Ji Q, Billups CA, Wu J et al. Dynamic contrast-enhanced magnetic resonance imaging as a prognostic factor in predicting event-free and overall survival in pediatric patients with osteosarcoma. *Cancer* 2012; **118**: 3776–3785.
10. Han G, Wang Y, Bi WZ, Wang DJ, Lu SB, Zhang L et al. Magnetic resonance imaging is appropriate for determining the osteotomy plane for appendicular osteosarcoma after neoadjuvant chemotherapy. *Med Oncol* 2012; **29**: 1347–1353.
11. Wright MW, Bruford EA. Naming 'junk': human non-protein coding RNA (ncRNA) gene nomenclature. *Human genomics* 2011; **5**: 90–98.
12. Biscotti MA, Olmo E, Heslop-Harrison JS. Repetitive DNA in eukaryotic genomes. *Chromosome Res* 2015; **23**: 415–420.
13. Pennisi E. Genomics. ENCODE project writes eulogy for junk DNA. *Science* 2012; **337**: 1159–1161.
14. Niu DK, Jiang L. Can ENCODE tell us how much junk DNA we carry in our genome? *Biochem Biophys Res Commun* 2013; **430**: 1340–1343.
15. Yang JX, Rastetter RH, Wilhelm D. Non-coding RNAs: an introduction. *Adv Exp Med Biol* 2016; **886**: 13–32.
16. Kunej T, Obsteter J, Pogacar Z, Horvat S, Calin GA. The decalog of long non-coding RNA involvement in cancer diagnosis and monitoring. *Crit Rev Clin Lab Sci* 2014; **51**: 344–357.
17. Goldman A, Capovano CA, Gonzalez-Lopez E, Geisinger A. Identifier (ID) elements are not preferentially located to brain-specific genes: high ID element representation in other tissue-specific- and housekeeping genes of the rat. *Gene* 2014; **533**: 72–77.
18. Calo E, Flynn RA, Martin L, Spitale RC, Chang HY, Wysocka J. RNA helicase DDX21 coordinates transcription and ribosomal RNA processing. *Nature* 2015; **518**: 249–253.
19. Zhong C, Zhang S. Clustering RNA structural motifs in ribosomal RNAs using secondary structural alignment. *Nucleic Acids Res* 2012; **40**: 1307–1317.
20. van Dijk M, Visser A, Buabeng KM, Poutsma A, van der Schors RC, Oudejans CB. Mutations within the LINC-HELLP non-coding RNA differentially bind ribosomal and RNA splicing complexes and negatively affect trophoblast differentiation. *Hum Mol Genet* 2015; **24**: 5475–5485.
21. Green D, Fraser WD, Dalmay T. Transfer RNA-derived small RNAs in the cancer transcriptome. *Pflügers Archiv* 2016; **468**: 1041–1047.
22. Casas E, Cai G, Neill JD. Characterization of circulating transfer RNA-derived RNA fragments in cattle. *Front Genet* 2015; **6**: 271.
23. O'Reilly D, Kuznetsova OV, Laitem C, Zaborowska J, Dienstbier M, Murphy S. Human snRNA genes use polyadenylation factors to promote efficient transcription termination. *Nucleic Acids Res* 2014; **42**: 264–275.
24. Penna I, Vassallo I, Nizzari M, Russo D, Costa D, Menichini P et al. A novel snRNA-like transcript affects amyloidogenesis and cell cycle progression through perturbation of Fe65L1 (APBB2) alternative splicing. *Biochim Biophys Acta* 2013; **1833**: 1511–1526.
25. Hung KH, Stumph WE. Regulation of snRNA gene expression by the *Drosophila melanogaster* small nuclear RNA activating protein complex (DmSNAPc). *Crit Rev Biochem Mol Biol* 2011; **46**: 11–26.
26. Qu G, Kruszka K, Plewka P, Yang SY, Chiou TJ, Jarmolowski A et al. Promoter-based identification of novel non-coding RNAs reveals the presence of dicistronic snoRNA-miRNA genes in *Arabidopsis thaliana*. *BMC Genomics* 2015; **16**: 1009.
27. Holley CL, Topkara VK. An introduction to small non-coding RNAs: miRNA and snoRNA. *Cardiovasc Drugs Ther* 2011; **25**: 151–159.
28. Scott MS, Avolio F, Ono M, Lamond AI, Barton GJ. Human miRNA precursors with box H/ACA snoRNA features. *PLoS Comput Biol* 2009; **5**: e1000507.
29. Wright MW. A short guide to long non-coding RNA gene nomenclature. *Hum Genomics* 2014; **8**: 7.
30. Siomi MC, Nishida KM, Siomi H. Chapter 16. How to define targets for small guide RNAs in RNA silencing: a biochemical approach. *Methods Enzymol* 2008; **449**: 345–355.
31. Wang T, Lander ES, Sabatini DM. Single guide RNA library design and construction. *Cold Spring Harbor Protocols* 2016; **2016**: pdb prot090803.
32. Randau L. Evolution of small guide RNA genes in hyperthermophilic archaea. *Ann NY Acad Sci* 2015; **1341**: 188–193.
33. Webb CJ, Zakian VA. Telomere les(i)ons from a telomerase RNA mutant. *Cell Cycle* 2015; **14**: 3769–3770.



34. Podlevsky JD, Li Y, Chen JJ. Structure and function of echinoderm telomerase RNA. *RNA* 2016; **22**: 204–215.
35. Kesharwani P, Gajbhiye V, Jain NK. A review of nanocarriers for the delivery of small interfering RNA. *Biomaterials* 2012; **33**: 7138–7150.
36. Wang XW, Li WD, Xia JR, Li Z, Cai XG. Small interfering RNA targeting receptor for advanced glycation end products suppresses the generation of proinflammatory cytokines. *Exp Ther Med* 2015; **10**: 584–590.
37. Lee SH, Kang YY, Jang HE, Mok H. Current preclinical small interfering RNA (siRNA)-based conjugate systems for RNA therapeutics. *Adv Drug Deliv Rev* 2015; **104**: 78–92.
38. Luo JW, Wang X, Yang Y, Mao Q. Role of micro-RNA (miRNA) in pathogenesis of glioblastoma. *Eur Rev Med Pharmacol Sciences* 2015; **19**: 1630–1639.
39. Montani F, Bianchi F. Circulating cancer biomarkers: the macro-revolution of the micro-RNA. *EBioMedicine* 2016; **5**: 4–6.
40. Casey MC, Sweeney KJ, Brown JA, Kerin MJ. Exploring circulating micro-RNA in the neoadjuvant treatment of breast cancer. *Int J Cancer* 2016; **139**: 12–22.
41. Fang W, Wang X, Bracht JR, Nowacki M, Landweber LF. Piwi-interacting RNAs protect DNA against loss during *Oxytricha* genome rearrangement. *Cell* 2012; **151**: 1243–1255.
42. Chu H, Xia L, Qiu X, Gu D, Zhu L, Jin J et al. Genetic variants in noncoding PIWI-interacting RNA and colorectal cancer risk. *Cancer* 2015; **121**: 2044–2052.
43. Menor MS, Baek K, Poisson G. Prediction of mature microRNA and piwi-interacting RNA without a genome reference or precursors. *Int J Mol Sci* 2015; **16**: 1466–1481.
44. He JH, Han ZP, Li YG. Association between long non-coding RNA and human rare diseases [review]. *Biomed Rep* 2014; **2**: 19–23.
45. Yu X, Li Z. Long non-coding RNA HOTAIR: a novel oncogene [review]. *Mol Med Rep* 2015; **12**: 5611–5618.
46. Ye LC, Zhu X, Qiu JJ, Xu J, Wei Y. Involvement of long non-coding RNA in colorectal cancer: from benchtop to bedside [review]. *Oncol Lett* 2015; **9**: 1039–1045.
47. Zhang A, Zhang J, Kaipainen A, Lucas JM, Yang H. Long non-coding RNA: a newly deciphered 'code' in prostate cancer. *Cancer Lett* 2016; **375**: 323–330.
48. Sun J, Lin Y, Wu J. Long non-coding RNA expression profiling of mouse testis during postnatal development. *PLoS One* 2013; **8**: e75750.
49. Rios-Barrera LD, Gutierrez-Perez I, Dominguez M, Riesgo-Escovar JR. *acal* is a long non-coding RNA in JNK signaling in epithelial shape changes during *Drosophila* dorsal closure. *PLoS Genet* 2015; **11**: e1004927.
50. Zhang Z, Zhou C, Chang Y, Zhang Z, Hu Y, Zhang F et al. Long non-coding RNA CASC11 interacts with hnRNP-K and activates the WNT/beta-catenin pathway to promote growth and metastasis in colorectal cancer. *Cancer Lett* 2016; **376**: 62–73.
51. Wu L, Jin L, Zhang W, Zhang L. Roles of long non-coding RNA CCAT2 in cervical cancer cell growth and apoptosis. *Med Sci Monit* 2016; **22**: 875–879.
52. Liu Z, Yan HY, Xia SY, Zhang C, Xiu YC. Downregulation of long non-coding RNA TRIM52-AS1 functions as a tumor suppressor in renal cell carcinoma. *Mol Med Rep* 2016; **13**: 3206–3212.
53. Ballantyne MD, Pinel K, Dakin R, Vesey AT, Diver L, Mackenzie R et al. Smooth muscle enriched long non-coding RNA (SMILR) regulates cell proliferation. *Circulation* 2016; **133**: 2050–2065.
54. Xu S, Wang P, You Z, Meng H, Mu G, Bai X et al. The long non-coding RNA EPB41L4-AS2 inhibits tumor proliferation and is associated with favorable prognoses in breast cancer and other solid tumors. *Oncotarget* 2016; **7**: 20704–20717.
55. Wan L, Sun M, Liu GJ, Wei CC, Zhang EB, Kong R et al. Long non-coding RNA PVT1 promotes non-small cell lung cancer cell proliferation through epigenetically regulating LATS2 expression. *Mol Cancer Ther* 2016; **15**: 1082–1094.
56. Olivieri M, Ferro M, Terreri S, Durso M, Romanelli A, Avitabile C et al. Long non-coding RNA containing ultraconserved genomic region 8 promotes bladder cancer tumorigenesis. *Oncotarget* 2016; **7**: 20636–20654.
57. Li J, Zhuang C, Liu Y, Chen M, Zhou Q, Chen Z et al. shRNA targeting long non-coding RNA CCAT2 controlled by tetracycline-inducible system inhibits progression of bladder cancer cells. *Oncotarget* 2016; **7**: 28989–28997.
58. Wu X, He X, Li S, Xu X, Chen X, Zhu H. Long non-coding RNA *ucoc2kmd.1* regulates CD44-dependent cell growth by competing for miR-211-3p in colorectal cancer. *PLoS One* 2016; **11**: e0151287.
59. Esposti DD, Hernandez-Vargas H, Voegele C, Fernandez-Jimenez N, Forey N, Bancel B et al. Identification of novel long non-coding RNAs deregulated in hepatocellular carcinoma using RNA-sequencing. *Oncotarget* 2016; **7**: 31862–31877.
60. Reed DE, Shokat KM. Targeting osteosarcoma. *Proc Natl Acad Sci USA* 2014; **111**: 18100–18101.
61. McQueen P, Ghaffar S, Guo Y, Rubin EM, Zi X, Hoang BH. The Wnt signaling pathway: implications for therapy in osteosarcoma. *Expert Rev Anticancer Ther* 2011; **11**: 1223–1232.
62. Robey RW, Polgar O, Deeken J, To KW, Bates SE. ABCG2: determining its relevance in clinical drug resistance. *Cancer Metastasis Rev* 2007; **26**: 39–57.
63. Natarajan K, Xie Y, Baer MR, Ross DD. Role of breast cancer resistance protein (BCRP/ABCG2) in cancer drug resistance. *Biochem Pharmacol* 2012; **83**: 1084–1103.
64. Martins-Neves SR, Paiva-Oliveira DI, Wijers-Koster PM, Abrunhosa AJ, Fontes-Ribeiro C, Bovee JV et al. Chemotherapy induces stemness in osteosarcoma cells through activation of Wnt/beta-catenin signaling. *Cancer Lett* 2016; **370**: 286–295.
65. Kumar RM, Fuchs B. Hedgehog signaling inhibitors as anti-cancer agents in osteosarcoma. *Cancers* 2015; **7**: 784–794.
66. Lo WW, Pinnaduwage D, Gokgoz N, Wunder JS, Andruis IL. Aberrant hedgehog signaling and clinical outcome in osteosarcoma. *Sarcoma* 2014; **2014**: 261804.
67. Lo WW, Wunder JS, Dickson BC, Campbell V, McGovern K, Alman BA et al. Involvement and targeted intervention of dysregulated Hedgehog signaling in osteosarcoma. *Cancer* 2014; **120**: 537–547.
68. Li B, Yang Y, Jiang S, Ni B, Chen K, Jiang L. Adenovirus-mediated overexpression of BMP-9 inhibits human osteosarcoma cell growth and migration through downregulation of the PI3K/AKT pathway. *Int J Oncol* 2012; **41**: 1809–1819.
69. Daigang L, Jining Q, Jinlai L, Pengfei W, Chuan S, Liangu H et al. LPS-stimulated inflammation inhibits BMP-9-induced osteoblastic differentiation through crosstalk between BMP/MAPK and Smad signaling. *Exp Cell Res* 2016; **341**: 54–60.
70. Liu C, Goswami M, Talley J, Chesser-Martinez PL, Lou CH, Sater AK. TAK1 promotes BMP4/Smad1 signaling via inhibition of erk MAPK: a new link in the FGF/BMP regulatory network. *Differ Res Biol Divers* 2012; **83**: 210–219.
71. Verheyen EM. Opposing effects of Wnt and MAPK on BMP/Smad signal duration. *Dev Cell* 2007; **13**: 755–756.
72. Wang L, Park P, Zhang H, La Marca F, Claeson A, Valdivia J et al. BMP-2 inhibits the tumorigenicity of cancer stem cells in human osteosarcoma OS99-1 cell line. *Cancer Biol Ther* 2011; **11**: 457–463.
73. Wang L, Park P, La Marca F, Than K, Rahman S, Lin CY. Bone formation induced by BMP-2 in human osteosarcoma cells. *Int J Oncol* 2013; **43**: 1095–1102.
74. Weiss KR, Cooper GM, Jadlovec JA, McGough RL III, Huard J. VEGF and BMP expression in mouse osteosarcoma cells. *Clin Orthop Relat Res* 2006; **450**: 111–117.
75. Yu L, Fan Z, Fang S, Yang J, Gao T, Simoes BM et al. Cisplatin selects for stem-like cells in osteosarcoma by activating notch signaling. *Oncotarget* 2016; **7**: 33055–33068.
76. Ongaro A, Pellati A, Bagheri L, Rizzo P, Caliceti C, Massari L et al. Characterization of Notch signaling during osteogenic differentiation in human osteosarcoma cell line MG63. *J Cell Physiol* 2016; **231**: 2652–2663.
77. Mei H, Yu L, Ji P, Yang J, Fang S, Guo W et al. Doxorubicin activates the Notch signaling pathway in osteosarcoma. *Oncol Lett* 2015; **9**: 2905–2909.
78. Ji P, Yu L, Guo WC, Mei HJ, Wang XJ, Chen H et al. Doxorubicin inhibits proliferation of osteosarcoma cells through upregulation of the Notch signaling pathway. *Oncol Res* 2015; **22**: 185–191.
79. McManus MM, Weiss KR, Hughes DP. Understanding the role of Notch in osteosarcoma. *Adv Exp Med Biol* 2014; **804**: 67–92.
80. Perry JA, Kiezun A, Tonzi P, Van Allen EM, Carter SL, Baca SC et al. Complementary genomic approaches highlight the PI3K/mTOR pathway as a common vulnerability in osteosarcoma. *Proc Natl Acad Sci USA* 2014; **111**: E5564–E5573.
81. Zhang J, Yu XH, Yan YG, Wang C, Wang WJ. PI3K/Akt signaling in osteosarcoma. *Clin Chim Acta* 2015; **444**: 182–192.
82. Song R, Tian K, Wang W, Wang L. P53 suppresses cell proliferation, metastasis, and angiogenesis of osteosarcoma through inhibition of the PI3K/AKT/mTOR pathway. *Int J Surg* 2015; **20**: 80–87.
83. Berman SD, Calo E, Landman AS, Danielian PS, Miller ES, West JC et al. Metastatic osteosarcoma induced by inactivation of Rb and p53 in the osteoblast lineage. *Proc Natl Acad Sci USA* 2008; **105**: 11851–11856.
84. Zhao J, Zhang ZR, Zhao N, Ma BA, Fan QY. VEGF silencing inhibits human osteosarcoma angiogenesis and promotes cell apoptosis via PI3K/AKT signaling pathway. *Int J Clin Exp Med* 2015; **8**: 12411–12417.
85. Zhang Y, Yang CQ, Gao Y, Wang C, Zhang CL, Zhou XH. Knockdown of CXCR7 inhibits proliferation and invasion of osteosarcoma cells through inhibition of the PI3K/Akt and beta-arrestin pathways. *Oncol Rep* 2014; **32**: 965–972.
86. Zhu LB, Jiang J, Zhu XP, Wang TF, Chen XY, Luo QF et al. Knockdown of Aurora-B inhibits osteosarcoma cell invasion and migration via modulating PI3K/Akt/NF-kappaB signaling pathway. *Int J Clin Exp Pathol* 2014; **7**: 3984–3991.
87. Zhao G, Cai C, Yang T, Qiu X, Liao B, Li W et al. MicroRNA-221 induces cell survival and cisplatin resistance through PI3K/Akt pathway in human osteosarcoma. *PLoS One* 2013; **8**: e53906.
88. Liu B, Qu L, Yang Z, Tao H. Cyclooxygenase-2 inhibitors induce anoikis in osteosarcoma via PI3K/Akt pathway. *Med Hypotheses* 2012; **79**: 98–100.
89. Gobin B, Huin MB, Lamoureux F, Ory B, Charrier C, Lanel R et al. BYL719, a new alpha-specific PI3K inhibitor: single administration and in combination with conventional chemotherapy for the treatment of osteosarcoma. *Int J Cancer* 2015; **136**: 784–796.
90. Gong C, Liao H, Wang J, Lin Y, Qi J, Qin L et al. LY294002 induces G0/G1 cell cycle arrest and apoptosis of cancer stem-like cells from human osteosarcoma via down-regulation of PI3K activity. *Asian Pacific J Cancer Prev* 2012; **13**: 3103–3107.
91. Li JP, Liu LH, Li J, Chen Y, Jiang XW, Ouyang YR et al. Microarray expression profile of long noncoding RNAs in human osteosarcoma. *Biochem Biophys Res Commun* 2013; **433**: 200–206.
92. Qian M, Yang X, Li Z, Jiang C, Song D, Yan W et al. P50-associated COX-2 extragenic RNA (PACER) overexpression promotes proliferation and metastasis of osteosarcoma cells by activating COX-2 gene. *Tumour Biol* 2015; **37**: 3879–3886.
93. Bernard D, Prasanth KV, Tripathi V, Colasse S, Nakamura T, Xuan Z et al. A long nuclear-retained non-coding RNA regulates synaptogenesis by modulating gene expression. *EMBO J* 2010; **29**: 3082–3093.

94. Dong Y, Liang G, Yuan B, Yang C, Gao R, Zhou X. MALAT1 promotes the proliferation and metastasis of osteosarcoma cells by activating the PI3K/Akt pathway. *Tumour Biol* 2015; **36**: 1477–1486.
95. Cai X, Liu Y, Yang W, Xia Y, Yang C, Yang S et al. Long noncoding RNA MALAT1 as a potential therapeutic target in osteosarcoma. *J Orthop Res* 2015; **34**: 932–941.
96. Fang D, Yang H, Lin J, Teng Y, Jiang Y, Chen J et al. 17Beta-estradiol regulates cell proliferation, colony formation, migration, invasion and promotes apoptosis by upregulating miR-9 and thus degrades MALAT-1 in osteosarcoma cell MG-63 in an estrogen receptor-independent manner. *Biochem Biophys Res Commun* 2015; **457**: 500–506.
97. Taniguchi M, Fujiwara K, Nakai Y, Ozaki T, Koshikawa N, Toshio K et al. Inhibition of malignant phenotypes of human osteosarcoma cells by a gene silencer, a pyrrrole-imidazole polyamide, which targets an E-box motif. *FEBS Open Bio* 2014; **4**: 328–334.
98. Chan LH, Wang W, Yeung W, Deng Y, Yuan P, Mak KK. Hedgehog signaling induces osteosarcoma development through Yap1 and H19 overexpression. *Oncogene* 2014; **33**: 4857–4866.
99. Lamora A, Mullard M, Amiaud J, Brion R, Heymann D, Redini F et al. Anticancer activity of halofuginone in a preclinical model of osteosarcoma: inhibition of tumor growth and lung metastases. *Oncotarget* 2015; **6**: 14413–14427.
100. Li H, Zhang K, Liu LH, Ouyang Y, Bu J, Guo HB et al. A systematic review of matrix metalloproteinase 9 as a biomarker of survival in patients with osteosarcoma. *Tumour Biol* 2014; **35**: 5487–5491.
101. Wang B, Su Y, Yang Q, Lv D, Zhang W, Tang K et al. Overexpression of long non-coding RNA HOTAIR promotes tumor growth and metastasis in human osteosarcoma. *Mol Cells* 2015; **38**: 432–440.
102. Ruan W, Wang P, Feng S, Xue Y, Li Y. Long non-coding RNA small nucleolar RNA host gene 12 (SNHG12) promotes cell proliferation and migration by upregulating angiominin gene expression in human osteosarcoma cells. *Tumour Biol* 2015; **37**: 4065–4073.
103. Wang Z, Cao CJ, Huang LL, Ke ZF, Luo CJ, Lin ZW et al. EFEMP1 promotes the migration and invasion of osteosarcoma via MMP-2 with induction by AEG-1 via NF-kappaB signaling pathway. *Oncotarget* 2015; **6**: 14191–14208.
104. Sun J, Wang X, Fu C, Wang X, Zou J, Hua H et al. Long noncoding RNA FGFR3-AS1 promotes osteosarcoma growth through regulating its natural antisense transcript FGFR3. *Mol Biol Rep* 2016; **43**: 427–436.
105. Wang H, Zhang X, Teng L, Legerski RJ. DNA damage checkpoint recovery and cancer development. *Exp Cell Res* 2015; **334**: 350–358.
106. Asghar U, Witkiewicz AK, Turner NC, Knudsen ES. The history and future of targeting cyclin-dependent kinases in cancer therapy. *Nat Rev Drug Discov* 2015; **14**: 130–146.
107. Khan Z, Bisen PS. Oncoapoptotic signaling and deregulated target genes in cancers: special reference to oral cancer. *Biochim Biophys Acta* 2013; **1836**: 123–145.
108. Min L, Hong S, Duan H, Zhou Y, Zhang W, Luo Y et al. Antidifferentiation noncoding RNA regulates the proliferation of osteosarcoma cells. *Cancer Biother Radiopharmaceut* 2016; **31**: 52–57.
109. Zhang Q, Geng PL, Yin P, Wang XL, Jia JP, Yao J. Down-regulation of long non-coding RNA TUG1 inhibits osteosarcoma cell proliferation and promotes apoptosis. *Asian Pacific J Cancer Prev* 2013; **14**: 2311–2315.
110. Wang Y, Yao J, Meng H, Yu Z, Wang Z, Yuan X et al. A novel long non-coding RNA, hypoxia-inducible factor-2alpha promoter upstream transcript, functions as an inhibitor of osteosarcoma stem cells *in vitro*. *Mol Med Rep* 2015; **11**: 2534–2540.
111. Cong M, Li J, Jing R, Li Z. Long non-coding RNA tumor suppressor candidate 7 functions as a tumor suppressor and inhibits proliferation in osteosarcoma. *Tumour Biol* 2016; **37**: 9441–9450.
112. Wang Z, Jin Y, Ren H, Ma X, Wang B, Wang Y. Downregulation of the long non-coding RNA TUSC7 promotes NSCLC cell proliferation and correlates with poor prognosis. *Am J Transl Res* 2016; **8**: 680–687.
113. Zhou Q, Chen F, Fei Z, Zhao J, Liang Y, Pan W et al. Genetic variants of lncRNA HOTAIR contribute to the risk of osteosarcoma. *Oncotarget* 2016; **7**: 19928–19934.
114. Ma B, Li M, Zhang L, Huang M, Lei JB, Fu GH et al. Upregulation of long non-coding RNA TUG1 correlates with poor prognosis and disease status in osteosarcoma. *Tumour Biol* 2015; **37**: 4445–4455.
115. Li F, Cao L, Hang D, Wang F, Wang Q. Long non-coding RNA HOTTIP is up-regulated and associated with poor prognosis in patients with osteosarcoma. *Int J Clin Exp Pathol* 2015; **8**: 11414–11420.
116. Liu T, Ma Q, Zhang Y, Ke S, Yan K, Chen X et al. Interleukin-11 receptor alpha is overexpressed in human osteosarcoma, and near-infrared-labeled IL-11Ralpha imaging agent could detect osteosarcoma in mouse tumor xenografts. *Tumour Biol* 2015; **36**: 2369–2375.
117. Zhu KP, Zhang CL, Shen GQ, Zhu ZS. Long noncoding RNA expression profiles of the doxorubicin-resistant human osteosarcoma cell line MG63/DXR and its parental cell line MG63 as ascertained by microarray analysis. *Int J Clin Exp Pathol* 2015; **8**: 8754–8773.
118. Diederichs S, Bartsch L, Berkmann JC, Frose K, Heitmann J, Hoppe C et al. The dark matter of the cancer genome: aberrations in regulatory elements, untranslated regions, splice sites, non-coding RNA and synonymous mutations. *EMBO Mol Med* 2016; **8**: 442–457.
119. Li T, Mo X, Fu L, Xiao B, Guo J. Molecular mechanisms of long noncoding RNAs on gastric cancer. *Oncotarget* 2016; **7**: 8601–8612.
120. Wang J, Wang H, Zhang Y, Zhen N, Zhang L, Qiao Y et al. Mutual inhibition between YAP and SRSF1 maintains long non-coding RNA, Malat1-induced tumorigenesis in liver cancer. *Cell Signal* 2014; **26**: 1048–1059.
121. Khachane AN, Harrison PM. Mining mammalian transcript data for functional long non-coding RNAs. *PLoS One* 2010; **5**: e10316.



**Cell Death and Disease** is an open-access journal published by Nature Publishing Group. This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016