



## WWP2 promotes degradation of transcription factor OCT4 in human embryonic stem cells

Huiming Xu<sup>1,2</sup>, Weicheng Wang<sup>1,2</sup>, Chunliang Li<sup>2</sup>, Hongyao Yu<sup>2</sup>, Acong Yang<sup>1,2</sup>, Beibei Wang<sup>2</sup>, Ying Jin<sup>1,2,3</sup>

Cell Research (2009) **19**:796. doi:10.1038/cr.2009.63; published online 1 June 2009

Correction to: (2009) 19:561-573. doi: 10.1038/cr.2009.31; published online 10 March 2009

The authors want to change the affiliations of this paper. The corrected affiliations are as follows.

Huiming Xu<sup>1,2</sup>, Weicheng Wang<sup>1,2</sup>, Chunliang Li<sup>2</sup>, Hongyao Yu<sup>1</sup>, Acong Yang<sup>1,2</sup>, Beibei Wang<sup>2</sup>, Ying Jin<sup>1,2,3</sup>

<sup>1</sup>Key Laboratory of Stem Cell Biology, Institute of Health Sciences, Shanghai Jiao Tong University School of Medicine & Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, 225 South Chongqing Road, Shanghai 200025, China; <sup>2</sup>Shanghai Stem Cell Institute, Shanghai Jiao Tong University School of Medicine, 225 South Chongqing Road, Shanghai 200025, China; <sup>3</sup>Key laboratory of Cell Differentiation and Apoptosis of Chinese Ministry of Education, Shanghai Jiao Tong University School of Medicine, Shanghai, China