

The mechanisms of *IDH* mutations in tumorigenesis

Dan Ye¹, Yue Xiong^{1,2,4}, Kun-Liang Guan^{1,3,5}

¹Molecular and Cell Biology Laboratory, Institutes of Biomedical Sciences, ²College of Life Science, ³College of Medicine, Fudan University, Shanghai 200032, China; ⁴Lineberger Comprehensive Cancer Center, Department of Biochemistry and Biophysics, University of North Carolina at Chapel Hill, NC 27599, USA; ⁵Department of Pharmacology and Moores Cancer Center, University of California San Diego, La Jolla, CA 92093, USA

Cell Research (2012) 22:1102-1104. doi:10.1038/cr.2012.51; published online 27 March 2012

Tumor-associated mutations in the isocitrate dehydrogenase 1 and 2 (*IDH1* and *IDH2*) genes result in the loss of normal catalytic activity, the production of α -ketoglutarate (α -KG), and gain of a new activity, the production of an oncometabolite, *R*-2-hydroxyglutarate (*R*-2-HG). New evidence supports previous findings that *R*-2-HG acts as an antagonist of α -KG to competitively inhibit the activity of multiple α -KG-dependent dioxygenases, including both histones and DNA demethylases involved in epigenetic control of gene expression and cell differentiation, and also reveals an intriguing new facet of *R*-2-HG in tumorigenesis.

The NADP⁺-dependent isocitrate dehydrogenase *IDH1* and *IDH2* catalyze the oxidative decarboxylation of isocitrate to α -ketoglutarate (α -KG). *IDH1* and *IDH2* are localized in the cytoplasm and mitochondria, respectively, and represent by far the most frequently mutated metabolic enzymes in human cancer [1]. The tumor-derived mutants of both *IDH1* and *IDH2* lose their activity in producing α -KG [2, 3], and gain

a surprising new catalytic activity, the production of *R*-2-hydroxyglutarate (*R*-2-HG) by reduction of α -KG [4]. Previous studies have shown that *R*-2-HG acts as an antagonist of α -KG to competitively inhibit a number of α -KG-dependent dioxygenases, including the JmjC domain-containing histone demethylases (KDMs) and the TET (ten-eleven translocation) family of DNA hydroxylases that catalyze the sequential oxidation of 5-methylcytosine (5mC) to 5-hydroxymethylcytosine (5hmC), 5-formylcytosine (5fC), and 5-carboxylcytosine (5caC), leading to eventual DNA demethylation (Figure 1) [5, 6]. Three papers recently published in *Nature* provide additional evidence that α -KG-dependent dioxygenases are the pathophysiological targets of mutant *IDH1/2*, and further underscore the presumptive role of *R*-2-HG as the first oncometabolite in contributing to tumorigenesis after *IDH1/2* mutations.

A subset of glioblastoma, known as the proneural subgroup, has previously found to display hypermethylation at a large number of loci and is enriched with *IDH1* mutations [7]. In one of the three *Nature* papers, Turcan *et al.* [8] determined whether *IDH1* mutation alone is sufficient to cause the hypermethylation phenotype by ectopic expression of *IDH1*^{R132H} mutant in immortalized primary human astrocytes, a cell type from which glioblastoma is believed to

develop. The authors found that introduction of mutant *IDH1* induced extensive DNA hypermethylation, altered the methylation of specific histones, and reshaped the methylome in a fashion that mirrors the changes observed in *IDH1*-mutated low-grade gliomas. The observed hypermethylation of DNA and histones can be explained by the direct inhibition of TET methylcytosine hydroxylases and JmjC family histone demethylases by *R*-2-HG, respectively. In keeping with the notion that TET hydroxylases directly regulate genomic DNA methylation levels and can be inhibited by the *R*-2-HG accumulated in *IDH1/2*-mutated cells, Turcan *et al.* also showed that ectopic expression of TET2 in cultured astrocytes decreased 5mC and increased 5hmC, and that both changes were inhibited by the co-expression of TET2 with mutant *IDH1*. These results are consistent with the findings made in acute myeloid leukemia (AML) in which *IDH1/2* and *TET2* genes are mutated in a mutually exclusive manner [9]. Moreover, Turcan *et al.* found that expression of wild-type *IDH1* decreased the average DNA methylation level in the genome, supporting the notion that the concentration of α -KG may be a rate-limiting factor of TET-catalyzed DNA demethylation [5].

In the second paper, Lu *et al.* [10] reported that ectopic expression of tumor-derived mutant *IDH1/2* or feed-

Correspondence: Dan Ye^a, Yue Xiong^b, Kun-Liang Guan^c

^aE-mail: yedan@fudan.edu.cn

^bE-mail: yxiong@email.unc.edu

^cE-mail: kuguan@ucsd.edu

Aberrant IDH1 and IDH2 activity and R-2-HG signaling in tumor cells




Figure 1 Summarization of reported mechanisms linking *IDH* mutation to tumorigenesis. Regulation of α -KG-dependent dioxygenases by *R*-2-HG is likely to play a major role in the pathophysiology of tumors with *IDH* mutation.

ing cells with cell-permeable *R*-2-HG increases histone demethylation and results in blockade of the differentiation of 3T3-L1 adipoblasts to adipocytes. These results indicate that mutation of *IDH1/2* and accumulation of *R*-2-HG can broadly impair cell differentiation beyond the cell types in which *IDH1/2* mutations are found to associate with tumorigenesis. The authors further confirmed that *IDH1*-mutated gliomas have elevated levels of histone methylation compared with gliomas retaining the wild-type *IDH1* [5, 6]. As previously reported [5, 6], multiple KDMs that are inhibited by 2-HG, including KDM4C/JMJD2C, which causes repressive histone H3K9 di- and trimethylation and, when suppressed by RNA interference, blocks the 3T3-L1 adipogenesis. It remains to be determined whether collective inhibition of multiple KDMs or a few individual ones, such as KDM4C, is responsible for altering cell differentiation in *IDH1/2*-mutated cells. The authors also noted that expression of mutant *IDH1* increased histone methylation prior to the increase of DNA

methylation, raising an intriguing possibility that histone methylation status may affect DNA methylation.

In the third paper, Koivunen *et al.* [11] proposed an enantiomer-specific mechanism of 2-HG in tumorigenesis. The authors reported two surprising findings. They showed first that immortalized human astrocytes stably expressing tumor-derived *IDH1*^{R132H} mutant proliferate faster during late passages than those expressing either wild-type *IDH1* or *IDH1*^{R132H/3DN} mutant that lacks 2-HG-producing activity. Ectopic expression of R132H mutant *IDH1* has previously been reported to decrease the growth of D54 glioblastoma cells [12], raising an intriguing possibility that the mutation of *IDH1/2* may exhibit different effects on cell growth in a cell context-dependent manner. More surprisingly, they found that *R*-2-HG, but not its enantiomer *S*-2-HG, substitutes for α -KG as a co-substrate, as opposed to an inhibitor, of EGLN, an α -KG-dependent prolylhydroxylase responsible for promoting the degradation of hypoxia inducible factor 1 α (HIF-1 α)

(Figure 1). As the result of stimulating EGLN, accumulation of *R*-2-HG was found to associate with diminished, instead of increased, HIF-1 α levels in cells expressing mutant *IDH1/2*. At first glance, these observations appear to be at odds with the generally accepted role of both enantiomers of 2-HG as inhibitors of α -KG-dependent dioxygenases, and HIF-1 α as an oncogene in tumorigenesis, but may at least in part explain the apparent selection for *IDH* mutations to produce *R*-, but not *S*-2-HG in cancer. This data, also for the first time, reveals a qualitatively different property of two 2-HG enantiomers with respect to α -KG-dependent dioxygenases. It will be interesting to determine the structural basis of this enantiomer-specific effect of 2-HG toward different α -KG-dependent dioxygenases. The observation that ectopic increase of *R*-2-HG reduces HIF-1 α suggests that endogenous α -KG is limiting for HIF-1 α hydroxylation by EGLN. The study by Koivunen *et al.* also suggests the complexity of EGLN regulation by *R*-2-HG and subsequent downregulation

of HIF-1 α . It remains to be determined genetically whether a reduction or fluctuation of HIF-1 α levels contributes to gliomagenesis in *IDH1/2*-mutated cells, because elevated HIF-1 α generally contributes to cancer development. The only piece of genetic evidence—*IDH1/2* mutation occurs in a mutually exclusive manner with *TET2* mutation in AML—supports the notion that epigenetic alteration plays a direct and perhaps a key role in *IDH1/2* mutation-associated tumorigenesis.

IDH1/2 mutation has rapidly emerged as a favorable diagnostic and prognostic marker for certain tumors, such as low-grade gliomas and benign cartilaginous tumors. While the full mechanism linking *IDH* mutation to tumorigenesis is incompletely understood, regulation of α -KG-dependent dioxygenases by 2-HG is likely to play a major role in the pathophysiology of tumors with *IDH* mutation. These recent reports also highlight the impact of altered metabolism and metabolites on the epigenetic modification of cell differentiation and tumorigenesis.

Acknowledgments

We thank previous and current members of Fudan MCB lab for the discussion and Eric Oermann for reading the manuscript.

References

- 1 Oermann EK, Wu J, Guan KL, Xiong Y. Alterations of metabolic genes and metabolites in cancer. *Semin Cell Dev Biol* 2012; **23**:370-380.
- 2 Yan H, Parsons DW, Jin G, *et al.* *IDH1* and *IDH2* mutations in gliomas. *N Engl J Med* 2009; **360**:765-773.
- 3 Zhao S, Lin Y, Xu W, *et al.* Glioma-derived mutations in *IDH1* dominantly inhibit *IDH1* catalytic activity and induce HIF-1 α . *Science* 2009; **324**:261-265.
- 4 Dang L, White DW, Gross S, *et al.* Cancer-associated *IDH1* mutations produce 2-hydroxyglutarate. *Nature* 2009; **462**:739-744.
- 5 Xu W, Yang H, Liu Y, *et al.* Oncometabolite 2-hydroxyglutarate is a competitive inhibitor of α -ketoglutarate-dependent dioxygenases. *Cancer Cell* 2011; **19**:17-30.
- 6 Chowdhury R, Yeoh KK, Tian YM, *et al.* The oncometabolite 2-hydroxyglutarate inhibits histone lysine demethylases. *EMBO Rep* 2011; **12**:463-469.
- 7 Noushmehr H, Weisenberger DJ, Diefes K, *et al.* Identification of a CpG island methylator phenotype that defines a distinct subgroup of glioma. *Cancer Cell* 2010; **17**:510-522.
- 8 Turcan S, Rohle D, Goenka A, *et al.* *IDH1* mutation is sufficient to establish the glioma hypermethylator phenotype. *Nature* 2012; **483**:479-483.
- 9 Figueroa ME, Abdel-Wahab O, Lu C, *et al.* Leukemic *IDH1* and *IDH2* mutations result in a hypermethylation phenotype, disrupt *TET2* function, and impair hematopoietic differentiation. *Cancer Cell* 2010; **18**:553-567.
- 10 Lu C, Ward PS, Kapoor GS, *et al.* *IDH* mutation impairs histone demethylation and results in a block to cell differentiation. *Nature* 2012; **483**:474-478.
- 11 Koivunen P, Lee S, Duncan CG, *et al.* Transformation by the (R)-enantiomer of 2-hydroxyglutarate linked to EGLN activation. *Nature* 2012; **483**:484-488.
- 12 Seltzer MJ, Bennett BD, Joshi AD, *et al.* Inhibition of glutaminase preferentially slows growth of glioma cells with mutant *IDH1*. *Cancer Res* 2010; **70**:8981-8987.