

Crystal structures of two phytohormone signal-transducing α/β hydrolases: karrikin-signaling KAI2 and strigolactone-signaling DWARF14

Cell Research (2013) 23:436-439. doi:10.1038/cr.2013.19; published online 5 February 2013

Dear Editor,

Karrikins and strigolactones (SL) are two classes of butenolide compounds that control many aspects of plant physiology. Karrikins, originally found in the smoke of forest fires, have the ability to stimulate seed germination [1]. SL are endogenous plant hormones that mediate responses to low levels of soil nutrients, such as phosphate and nitrate [2]. They promote root branching to increase nutrient uptake while inhibiting shoot branching to reduce nutrient demand. SL are also secreted from roots to stimulate symbiotic associations with arbuscular mycorrhizal fungi for increased nutrient uptake, signals that are exploited to induce seed germination in parasitic weeds, including *Striga* and *Orobancha*, which are major causes of crop losses [2].

Genetic studies in *Arabidopsis* have identified KAI2 and MAX2 as two key players in karrikin signal transduction [3]. KAI2 has high sequence similarity to the bacterial signaling protein RbsQ [4] (Supplementary information, Figure S1), which encodes an α/β hydrolase. MAX2 is the F-box component of SCF E3 ubiquitin ligase. SL signaling is also mediated by MAX2 and by a paralog of KAI2, D14 [2]. Moreover, a D14 homolog from petunia, DAD2, has hydrolytic activity toward the synthetic SL, GR24, and this activity has been proposed to be essential for SL perception [5]. Similarly, hydrolysis of the butenolide moiety of karrikins by KAI2 has been proposed as a part of the karrikin signaling mechanism [6]. However, the mechanisms of catalysis as well as the molecular bases of the signaling specificity of KAI2 and D14 toward karrikins and SL remain unclear.

To further understand their functions, we crystallized KAI2 and D14 proteins from *Arabidopsis thaliana* (AtKAI2 and AtD14) and D14 from rice, *Oryza sativa* (OsD14), and determined their structures (see Supplementary information, Data S1 and Table S1). The overall structures confirm that they are members of the α/β hydrolase superfamily. All 3 structures share a common

fold of a seven-stranded β -sheet ($\beta 2$ - $\beta 8$) surrounded by five helices ($\alpha 0$, αB - αE) at one side and two (αA and αF) at the other, and a top domain displaying a double layer V-shaped helical fold containing four helices ($\alpha T1$ - $\alpha T4$) that harbor a substrate-binding pocket (Figure 1A). The two D14 structures are almost superimposable, as well as the recently solved structure of petunia DAD2 [5] (Supplementary information, Figure S3; rmsd < 0.85 Å), confirming that all three proteins are D14 orthologs.

The catalytic triad residues of the D14 proteins are S97, H247 and D218, and those of KAI2 are S95, H246 and D217, all of which are located at the bottom of the hydrophobic substrate-binding pocket and on the loops following the $\beta 4$, $\beta 7$ and $\beta 6$ strands, respectively (Figure 1A-C). The triad serine residue of hydrolases functions as highly reactive nucleophile that binds to and hydrolyzes substrates, while the histidine and aspartate residues form a charge relay network to increase the nucleophilicity of the serine and to function as acceptor of the serine hydroxyl proton.

Molecular docking of karrikins into the KAI2 pocket (Figure 1B, 1C and Supplementary information, Figure S2) and of GR24 into the AtD14 pocket (Figure 1B) indicated a snug fit of these compounds in pockets lined by bulky aromatic side chains. The lactone carboxyl groups of the docked karrikin and SL are in close proximity to the hydroxyl groups of the triad serine residues (2.9 Å and 2.5 Å, respectively) to allow a nucleophilic attack, supported by the histidine and aspartate charge relay network, that would result in the hydrolysis of the butenolide rings (Figure 1B, 1C and Supplementary information, Figure S4). Ultra performance liquid chromatography (UPLC) combined with ES-MS confirmed the hydrolysis of the GR24 butenolide ring and identified the GR24 ABC-ring and lactone D-ring as final hydrolysis products (Supplementary information, Figures S4 and S6).

When we tried to co-crystallize OsD14 with GR24, we obtained the crystal structure of D14 covalently bound




Figure 1 Structures and activities of KAI2 and D14 proteins. **(A)** Structure overview of apo KAI2, AtD14, and OsD14. Ligand-binding pockets are indicated as mesh. **(B)** KAI2 and D14 ligand-binding pockets (top) and docked ligands (bottom). **(C)** Close-up view of the KAI2 catalytic triad with docked ligand. **(D)** Structural alignment of ligands and key ligand specificity-conferring residues. **(E)** Hydrolase activity of KAI2, D14, and SABP2 toward a generic small substrate (left) and GR24 (right).

to a GR24 degradation intermediate, 2,4,4-trihydroxy-3-methyl-3-butenal, which is clearly revealed by the electron density map (Supplementary information, Figure S4A-S4C). The transition state captured in this structure shows the hydroxyl group of S97 attached to C1 of the intermediate, which is further stabilized by water-mediated hydrogen bonds with H247 and Y159 (Supplementary information, Figure S4B). This intermediate suggests that the initial nucleophilic attack causes an electron shift, followed by the addition of a water molecule, to lead to the release of the ABC ring product and the formation of a S97-stabilized open lactone. The latter then converts to the intermediate found in the crystal structure by 1,4-addition of a water molecule to the two conjugated double bonds. The C1 enol tautomer of the transition product can then undergo an intra-molecular Michael addition of the C1 hydroxyl to the carbonyl group at position 4 to form the final closed lactone ring (Supplementary information, Figure S4D) identified by MS (Supplementary information, Figure S6).

Despite that KAI2 has high sequence and structure similarity to the two D14 proteins (Supplementary information, Figure S1), KAI2 is not able to mediate the strigolactone branching signaling, and the D14 proteins do not promote seed germination in response to karrikin exposure [7]. While the overall structures of these hydrolases are very similar ($\text{rmsd} \leq 1.15 \text{ \AA}$), their substrate binding pockets differ significantly. KAI2 has a relatively small pocket of 279 \AA^3 compared to the larger pockets of AtD14 (357 \AA^3) and OsD14 (432 \AA^3). In addition, the pocket of KAI2 is constricted in the middle by the inward shift of helix α T4, whose bulky residues I193 (V194 in both D14 proteins) and F194 face into the pocket (Figure 1B, 1D). Overlay of the apo structures of KAI2 and D14 indicates that these two residues of KAI2 would clash with the position of GR24 as docked into the D14 binding pocket. Conversely, the larger binding pocket of the D14 proteins cannot accommodate karrikin as docked into the KAI2 pocket because the KAI2 ligand-binding residue F157 is replaced with a tyrosine in D14, whose hydroxyl group clashes with the oxygen atom of the 6-member pyran ring of karrikin (Figure 1D). Therefore, the positions of 3 bulky hydrophobic residues, I193/IV194, F194/F195, and F157/Y159 are likely the main determinants of ligand specificity.

To biochemically assess hydrolase activity and specificity, we first determined the activities of KAI2 and the D14 proteins towards a small generic hydrolase substrate, 4-nitrophenyl butyrate, and compared them to the activity of SABP2, a methyl salicylate-cleaving α/β hydrolase involved in systemic acquired resistance signaling [8, 9]. Our biochemical assays demonstrated that

SABP2 is able to hydrolyze 4-nitrophenyl butyrate, and its enzymatic activity can be entirely blocked by PMSF, a commonly used hydrolase inhibitor (Figure 1E). Despite the structural similarity to SABP2, KAI2 and D14 do not show significant esterase activity for 4-nitrophenyl butyrate, either in the presence or absence of PMSF, and the low activity of KAI2 and D14 are not inhibited by PMSF (Figure 1E). The crystal structure of SABP2 in complex with salicylic acid [8] revealed a small binding pocket of 167 \AA^3 , whose entrance is covered by a cap domain (Supplementary information, Figure S5), which can enhance the capture of the substrate during catalysis. In contrast, both KAI2 and D14 have a rigid open entrance to their substrate binding pockets where a bound substrate can leave before being hydrolyzed, which may explain why KAI2 and D14 have much lower activity than SABP2. In contrast to the activities toward 4-nitrophenyl butyrate, only the D14 proteins were able to hydrolyze GR24, demonstrating hormone substrate specificity for this class of signaling hydrolases. However, the GR24 hydrolase activity is extremely low, with a turnover rate of only one GR24 molecule per D14 molecule per three minutes.

D14 and KAI2 are absolutely required for SL and karrikin signaling, respectively, and are functionally linked to the same F-box protein, AtMAX2/OsD3. This pathway is reminiscent of gibberellin signal perception and transduction, in which the catalytically inactive α/β -hydrolase GID1 functions as the hormone receptor to mediate hormone-dependent complex formation between GID1, DELLA transcriptional repressors that function as coreceptors, and the GID2 F-box protein [10-12]. F-box-binding mediated degradation of the DELLA repressors leads then to induction of gibberellin-responsive genes. While SL and karrikins might be precursors of the actual signaling molecules, thought to be generated by D14- and KAI2-catalyzed hydrolysis, the extremely low hydrolase activity and the apparent lack of signaling activity of GR24 hydrolytic products [5] suggest that D14 and KAI2 may instead function as hormone receptors in analogy to GID1. Moreover, while endogenous SL occur at low concentrations, D14 alone exhibits only weak substrate-binding affinity, consistent with a possible coreceptor requirement for high-affinity binding. We speculate that D14 and KAI2, which unlike SABP2 lack moveable lids to control access to the ligand-binding pocket, likely associate with coreceptors in the ligand-bound state to retain the ligands in the pocket, thereby increasing ligand/substrate affinity. In analogy with GID1, and consistent with a GR24-mediated Y2H interaction between petunia D14 and MAX2 [5], a ternary receptor-hormone-coreceptor complex might recruit SCF^{D3/MAX2} for coreceptor

ubiquitination and degradation. Association and possible degradation of SL- and karrikin-specific corepressors would therefore provide a simple explanation for D14 and KAI2 signaling through the same F-box protein.

In summary, the structures reported in this study revealed unique pocket topologies as a basis for karrikin and SL signaling specificities, identified the pathway and mechanism of D14-catalyzed GR24 hydrolysis, and further support the likely roles of KAI2 and D14 in karrikin and SL perception.

Acknowledgments

This work was supported by the Jay and Betty Van Andel Foundation, Amway (China), the National Natural Science Foundation of China (NSFC 91217311) and US National Institute of Health (R01GM102545). The atomic coordinates have been deposited in the Protein Data Bank with accession codes listed in Supplementary information, Table S1. We thank staff members at Shanghai Synchrotron Radiation Facility (SSRF) beamline BL17U and the Life Science Collaborative Access Team of the Advanced Photon Source (APS) for assistance in data collection at the beam lines of sector 21, which is in part funded by the Michigan Economic Development Corporation and the Michigan Technology Tri-Corridor (Grant 085P1000817). Use of APS was supported by the Office of Science of the US Department of Energy, under Contract No. DE-AC02-06CH11357.

Li-Hua Zhao^{1,*}, X Edward Zhou^{2,*},
Zhong-Shan Wu^{1,3,*}, Wei Yi¹, Yong Xu^{1,4},
Suling Li¹, Ting-Hai Xu¹, Yue Liu¹, Run-Ze Chen¹,
Amanda Kovach², Yangyong Kang², Li Hou^{1,2},
Yuanzheng He², Cen Xie⁵, Wanling Song⁵,
Dafang Zhong⁵, Yechun Xu⁵, Yonghong Wang⁶,
Jiayang Li⁶, Chenghai Zhang¹, Karsten Melcher²,
H Eric Xu^{1,2}

¹VARI-SIMM Center, Center for Structure and Function of Drug Targets, CAS-Key Laboratory of Receptor Research, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China; ²Labora-

tory of Structural Sciences, Center for Structural Biology and Drug Discovery, Van Andel Research Institute, MI 49503, USA; ³Wuhan National Laboratory for Optoelectronics, Britton Chance Center for Biomedical Photonics, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China; ⁴Institute of Chemical Biology, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, Guangdong 510530, China; ⁵Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China; ⁶State Key Laboratory of Plant Genomics and National Center for Plant Gene Research (Beijing), Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China

*These three authors contributed equally to this work.

Correspondence: H Eric Xu^a, Karsten Melcher^b, Chenghai Zhang^c

^aE-mail: eric.xu@vai.org

^bE-mail: karsten.melcher@vai.org

^cE-mail: chenghai@mail.shnc.ac.cn

References

- 1 Flematti GR, Ghisalberti EL, Dixon KW, *et al.* *Science* 2004; **305**:977.
- 2 Brewer PB, Koltai H, Beveridge CA. *Mol Plant* 2012 Nov 15. doi: 10.1093/mp/sss130
- 3 Nelson DC, Scaffidi A, Dun EA, *et al.* *Proc Natl Acad Sci USA* 2011; **108**:8897-8902.
- 4 Kaneko T, Tanaka N, Kumasaka T. *Protein Sci* 2005; **14**:558-565.
- 5 Hamiaux C, Drummond RS, Janssen BJ, *et al.* *Curr Biol* 2012; **22**:2032-2036.
- 6 Scaffidi A, Waters MT, Bond CS, *et al.* *Bioorg Med Chem Lett* 2012; **22**:3743-3746.
- 7 Waters MT, Nelson DC, Scaffidi A, *et al.* *Development*; 2012; **139**:1285-1295.
- 8 Forouhar F, Yang Y, Kumar D, *et al.* *Proc Natl Acad Sci USA* 2005; **102**:1773-1778.
- 9 Kumar D, Klessig DF. *Proc Natl Acad Sci USA* 2003; **100**:16101-16106.
- 10 Shimada A, Ueguchi-Tanaka M, Nakatsu T, *et al.* *Nature* 2008; **456**:520-523.
- 11 Murase K, Hirano Y, Sun TP, *et al.* *Nature* 2008; **456**:459-463.
- 12 Sun TP. *Plant Physiol* 2010; **154**:567-570.

(Supplementary information is linked to the online version of the paper on the *Cell Research* website.)



This work is licensed under the Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0>