

RESEARCH HIGHLIGHT

Aberrant ceRNA activity drives lung cancer

Cell Research (2014) 24:259-260. doi:10.1038/cr.2014.21; published online 14 February 2014

In a recent issue of *Nature*, Kumar *et al.* demonstrate that the oncogenic potential of the *Hmga2* gene is largely due to the ability of its transcript to operate as a competing endogenous RNA in a protein coding-independent manner. The *Hmga2* mRNA decoys the let-7 microRNA family to regulate *Tgfb3* expression and enhance TGF- β signaling, thereby promoting lung cancer progression.

MicroRNAs are a large class of small non-coding RNAs which regulate gene expression by binding to microRNA response elements (MREs) on target transcripts. It has become increasingly clear in recent years that transcripts which contain MREs for shared microRNAs can co-regulate each other by titrating microRNA availability, thus acting as natural microRNA sponges or competing endogenous RNAs (ceRNAs) [1, 2]. Although ceRNA activity has been attributed to both protein-coding and non-coding RNAs such as small non-coding RNAs, long non-coding RNAs, pseudogenes and circular RNAs in diverse species, little is known about the precise molecular conditions needed for optimal ceRNA crosstalk.

A recent study by Kumar *et al.* has led to important insights into this recently described post-transcriptional regulatory dimension [3]. The authors focused on *Hmga2*, a non-histone chromosomal high-mobility group protein which contributes to lung cancer progression and metastasis. The *Hmga2* 3' UTR contains seven conserved MREs for the let-7 family of microRNAs [4], which has previously been demonstrated to constrain lung cancer development [5]. Kumar *et al.* generated a series of

Hmga2 expression constructs (wild-type *Hmga2*, WT; *Hmga2* with all seven let-7 binding sites mutated, m7; *Hmga2* with the single in-frame start codon mutated, ATG WT; and *Hmga2* with both the start codon and let-7 binding sites mutated, ATG m7). This allelic series enabled them to disentangle the contribution of *Hmga2*'s function as a let-7 ceRNA from its protein-coding function in lung cancer development.

Kumar *et al.* found that both WT and ATG WT constructs dramatically promoted anchorage-independent growth of lung cancer cell lines *in vitro*, rescued tumor formation in an autochthonous lung cancer model *in vivo*, and reduced survival of transplanted mice. A more modest effect was observed with the m7 construct and no growth was observed with the double mutant ATG m7 construct. Critically, the *in vitro* effects were rescued by exogenous expression of let-7. These results indicate that the interaction of the *Hmga2* transcript with let-7 is critical for the oncogenic function while the *Hmga2* protein is largely dispensable for lung cancer progression, suggesting a ceRNA function for the *Hmga2* transcript.

To examine potential targets of *Hmga2*'s ceRNA activity, the authors predicted let-7 targets among mRNAs that are differentially expressed in *Hmga2*-expressing vs non-expressing lung cancer cells. This approach identified six ceRNA targets that are regulated by *Hmga2* in a let-7-dependent manner. Subsequent analyses validate the TGF- β co-receptor *Tgfb3* as a *bona fide* target of *Hmga2* ceRNA (Figure 1A), and establish *Tgfb3* and TGF- β signaling to be critical for the oncogenic effect

of *Hmga2*. Importantly, *HMGA2* and *TGFBR3* expression is reciprocally positively correlated in human lung cancer, indicating that the oncogenic ceRNA function of *HMGA2* is conserved across species.

Kumar *et al.* performed several important experiments that further validated ceRNA crosstalk between *Hmga2* and *Tgfb3* and addressed a number of key open questions about such crosstalk. First, the authors demonstrated that *Tgfb3* recruitment to the Ago2-based microRNA repression complex was reduced in both *Hmga2* WT and ATG WT cells by RNA immunoprecipitation. Intriguingly, the authors reported that *Hmga2*'s ceRNA activity does not result in changes in the levels of let-7 isoforms or their association with Ago2, suggesting that *Hmga2* was able to block the association of *Tgfb3* with Ago2 via its function as a let-7 ceRNA. As previous studies have shown that microRNA decoying can direct the degradation and thus decrease the abundance of decoyed microRNAs [6], this observation suggests that ceRNA regulation may have varying effects on targeted microRNAs which may be sequence-specific and context-dependent.

Moreover, RNA-seq analyses demonstrated that *Hmga2* was one of the most highly expressed predicted let-7 target transcripts, *Hmga2* and *Tgfb3* transcripts were expressed at similar levels, and total expression levels of let-7 family members was within an order of magnitude of *Hmga2* and *Tgfb3*. These observations are consistent with previous reports suggesting that optimal ceRNA crosstalk would occur when the abundance of microRNA and ceRNA




Figure 1 (A) *Hmga2* acts as a natural microRNA sponge for the let-7 family, which also targets *Tgfr3*. An increase in *Hmga2* transcript levels and hence ceRNA activity results in a concomitant increase in *Tgfr3* levels. (B) *Hmga2* also regulates the expression of five other target genes via ceRNA crosstalk as well as through *Tgfr3*/TGF- β -mediated transcriptional control.

transcripts were near equimolarity [7]. *Hmga2* shares let-7 sites with numerous other transcripts besides *Tgfr3* and, considering simple stoichiometry, the effect of reduced let-7 availability in response to *Hmga2* overexpression should be “diluted” by all let-7 targets. Indeed, using Sylamer analysis, Kumar *et al.* report an enrichment of let-7 site-containing transcripts upon *Hmga2* overexpression, suggesting a more global effect on let-7 targets. Intriguingly, however, only six differentially expressed let-7 targets were identified as *Hmga2* ceRNA targets. These data could indicate a marginal effect of *Hmga2*-mediated sequestration of let-7 on most let-7 targets, while a few let-7 targets are affected significantly. Stoichiometry dictates that the overexpression of an abundant transcript such as *Hmga2* has profound ceRNA effects on lowly expressed let-7 targets. However, five out of six identified *Hmga2* ceRNA targets (*Tgfr3*, *Angptl2*, *Fndc3*, *Hmga1*, and *Skil*) are among the highly expressed let-7 targets, suggesting that additional factors such as subcellular localization may determine the effect of let-7 sponging. Indeed, another prominent target of the let-7 family, the potent oncogene K-Ras, was unaffected by *Hmga2* overexpression. Future work will provide further insight into the intricacies of regulation of ceRNA crosstalk. Additionally, recent reports have identified other *bona fide*

Hmga2-targeting microRNAs, including miR-33a and miR-154 [8, 9]. The ability of *Hmga2* to sequester these and other additional microRNAs may confer further complexity to its function as a ceRNA and should be investigated in future studies.

This study also provided evidence for the co-evolution of ceRNA and transcriptional networks. Kumar *et al.* found that the five *Hmga2* ceRNA targets besides *Tgfr3* are transcriptionally regulated by TGF- β signaling. This suggests that *Hmga2* regulates the expression of these transcripts in a feed-forward manner, first through ceRNA crosstalk, and second through transcriptional control via the *Tgfr3*-TGF- β axis (Figure 1B). Many more of such integrated ceRNA and transcriptional networks likely remain to be found, and their identification would indicate that ceRNA crosstalk is not a mere coincidence but a sophisticated means to fine-tune biological networks.

In summary, this report by Kumar *et al.* provides convincing evidence for the provocative hypothesis that in specific contexts, the primary function of a protein-coding gene may act as a microRNA decoy. It is also the first study which provides mutational analyses of the relevant microRNA-binding sites to demonstrate the functional relevance of ceRNA activity in tumorigenesis *in vivo*. Furthermore, Kumar *et al.* demonstrate that even highly abundant microRNAs

like let-7 can be effectively regulated by highly expressed ceRNAs with multiple MREs. As both *Hmga2* and let-7 have been implicated in multiple cancers as well as developmental processes, further insights into the *Hmga2*-let7 ceRNA network will have important implications for human health and disease.

Yvonne Tay¹, Florian A Karreth¹,
Pier Paolo Pandolfi¹

¹Cancer Research Institute, Beth Israel Deaconess Cancer Center, Department of Medicine and Pathology, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA 02215, USA

Correspondence: Pier Paolo Pandolfi
Tel: +1-617-735-2121; Fax: +1-617-735-2120
E-mail: ppandolfi@bidmc.harvard.edu

References

- Salmena L, Poliseno L, Tay Y, *et al.* *Cell* 2011; **146**:353-358.
- Tay Y, Rinn J, Pandolfi PP. *Nature* 2014; **505**:344-352.
- Kumar MS, Armenteros-Montero E, East P, *et al.* *Nature* 2014; **505**:212-217.
- Mayr C, Hemann MT, Bartel DP. *Science* 2007; **315**:1576-1579.
- Kumar MS, Erkeland SJ, Pester RE, *et al.* *Proc Natl Acad Sci USA* 2008; **105**:3903-3908.
- Cazalla D, Yario T, Steitz JA. *Science* 2010; **328**:1563-1566.
- Ala U, Karreth FA, Bosia C, *et al.* *Proc Natl Acad Sci USA* 2013; **110**:7154-7159.
- Rice SJ, Lai SC, Wood LW, *et al.* *J Biol Chem* 2013; **288**:16348-16360.
- Zhu C, Li J, Cheng G, *et al.* *Mol Cell Biochem* 2013; **379**:69-75.