

ORIGINAL ARTICLE

Genome wide linkage disequilibrium in Chinese asparagus bean (*Vigna. unguiculata* ssp. *sesquipedialis*) germplasm: implications for domestication history and genome wide association studies

P Xu¹, X Wu¹, B Wang¹, J Luo², Y Liu¹, JD Ehlers³, TJ Close³, PA Roberts³, Z Lu¹, S Wang¹ and G Li¹

Association mapping of important traits of crop plants relies on first understanding the extent and patterns of linkage disequilibrium (LD) in the particular germplasm being investigated. We characterize here the genetic diversity, population structure and genome wide LD patterns in a set of asparagus bean (*Vigna. unguiculata* ssp. *sesquipedialis*) germplasm from China. A diverse collection of 99 asparagus bean and normal cowpea accessions were genotyped with 1127 expressed sequence tag-derived single nucleotide polymorphism markers (SNPs). The proportion of polymorphic SNPs across the collection was relatively low (39%), with an average number of SNPs per locus of 1.33. Bayesian population structure analysis indicated two subdivisions within the collection sampled that generally represented the 'standard vegetable' type (subgroup SV) and the 'non-standard vegetable' type (subgroup NSV), respectively. Level of LD (r^2) was higher and extent of LD persisted longer in subgroup SV than in subgroup NSV, whereas LD decayed rapidly (0–2 cM) in both subgroups. LD decay distance varied among chromosomes, with the longest (≈ 5 cM) five times longer than the shortest (≈ 1 cM). Partitioning of LD variance into within- and between-subgroup components coupled with comparative LD decay analysis suggested that linkage group 5, 7 and 11 may have undergone the most intensive epistatic selection toward traits favorable for vegetable use. This work provides a first population genetic insight into domestication history of asparagus bean and demonstrates the feasibility of mapping complex traits by genome wide association study in asparagus bean using a currently available cowpea SNPs marker platform. *Heredity* (2012) **109**, 34–40; doi:10.1038/hdy.2012.8; published online 29 February 2012

Keywords: asparagus bean; association mapping; cowpea; domestication history; linkage disequilibrium; population structure

INTRODUCTION

Linkage disequilibrium (LD) refers to association of particular allelic configurations at distinct loci in the genome of a sampled population (Weir, 1979). LD mapping or association mapping can be applied to natural populations, sets of germplasm accessions or cultivars developed recently (Rostoks *et al.*, 2006; Robbins *et al.*, 2011) and assumes that only markers in strong LD with a functionally important gene will be significantly associated to the variation in quantitative traits (Ardlie *et al.*, 2002; Garris *et al.*, 2003). LD mapping thus takes advantage of the much larger number of historical recombination events that have occurred over time compared with the relatively restricted levels of recombination inherent in biparental mapping populations where quantitative trait loci (QTL) mapping studies have frequently been used to unravel the genetic basis of traits (Cardon and Bell, 2001). In order to appropriately apply LD mapping in crop plants, it is a prerequisite to understand the extent and patterns of LD in the particular populations being investigated (Lander and Schork, 1994). It is also crucial to be able to distinguish between physical LD and the other different forces that can create LD in natural populations, to avoid the detection of spurious associations (Flint-Garcia *et al.*, 2003).

LD can be caused by unknown population structure and several forces, including mutation, drift, genetic bottlenecks, founder effects, selection, and specifically for plants, level of inbreeding caused by their mating systems (Hartl and Clark, 1997). In contrast, physical LD tends to be continuously decreasing because of accumulated recombinations, which means that loci located far apart along chromosomes will generally remain in LD for shorter periods (Hartl and Clark, 1997; Mather *et al.*, 2007). Outcrossing species generally exhibit low levels of LD because of many opportunities for effective recombination events in large highly heterozygous populations (Yan *et al.*, 2009). In contrast, many crop species including asparagus bean, are inbreeders and would thus be expected to exhibit relatively higher levels of LD (Flint-Garcia *et al.*, 2003; Morrell *et al.*, 2005).

The extent and patterns of LD can be investigated on a haplotype, chromosome or whole genome level. Tenailon *et al.* (2001) sequenced 21 loci located on the maize chromosome 1 and estimated that in landraces the LD decay distance was < 1000 bp. In contrast, a study of 18 maize genes in 36 commercial inbred lines revealed extensive LD blocks as long as 100 Kb (Ching *et al.*, 2002). Thus, as pointed out by Yan *et al.* (2009), LD decay estimation based on a single chromosome or a limited number of loci may be biased. Uncovering LD patterns of

¹Institute of Vegetables, Zhejiang Academy of Agricultural Sciences, Hangzhou, PR China; ²Institute of Digital Agricultural Research, Zhejiang Academy of Agricultural Sciences, Hangzhou, PR China and ³Department of Botany and Plant Sciences, University of California, Riverside, CA, USA
Correspondence: Professor G Li, Institute of Vegetables, Zhejiang Academy of Agricultural Sciences, 198 shiqiao Road, Hangzhou 310021, PR China.
E-mail: Guojing_li@yahoo.com.cn

Received 18 July 2011; revised 31 December 2011; accepted 23 January 2012; published online 29 February 2012

plant species of interest at a genome level is therefore essential for the optimal design of genotyping efforts before conducting LD mapping and for assessing the statistical power and resolution in whole-genome association studies (McNally *et al.*, 2009).

Cowpea (*Vigna unguiculata* L. Walp., $2n=2x=22$) is an important self-pollinating grain legume, fodder and vegetable crop in many tropical/subtropical regions of the world. Two main divisions of cultivated cowpea are the dominant subspecies *unguiculata* used primarily for dry grain and fodder and *sesquipedialis*, which is harvested at the immature green pod stage and used as a vegetable (Timko *et al.*, 2007). The latter is also known as asparagus bean or 'yard-long' bean, for it is characterized by its very long (0.5–1 m) pods with less fiber, narrow kidney-shaped seeds and climbing growth habit. Genetic similarity between the two subspecies is high as evidenced by fully fertile hybrids and synteny for most of the shared molecular markers between genetic maps (Muchero *et al.*, 2009; Xu *et al.*, 2011). Cowpea is thought to have undergone a severe bottleneck during domestication (Panella and Gepts, 1992). Although genetic diversity has not been comprehensively assessed in the asparagus bean germplasm, it is likely low, keeping the assumption that the crop may have derived from a limited sample of the wider cowpea gene pool that moved from Africa to east Asia, followed by a further genetic narrowing as strong selection for long pods gave rise to the present day ssp. *sesquipedialis* (Fang *et al.*, 2007; Xu *et al.*, 2010). Assessing both genetic variation levels and genome-wide LD patterns in asparagus bean thus would not only be important for LD mapping but also shed some light on the past effects of intensive human selection for vegetable use, compared with materials that would be closer to the ancestral species.

Recently, a high-throughput Illumina GoldenGate assay platform with 1536 expressed sequence tag-derived single nucleotide polymorphism markers (SNPs) became available for cowpea. Using this system, high-density consensus genetic linkage maps for the ssp. *unguiculata* were constructed (Muchero *et al.*, 2009; Lucas *et al.*, 2011). A comparable genetic linkage map of asparagus bean also was constructed in the authors' lab using the same system and additional SSR markers (Xu *et al.*, 2011). The main objective of the present study was to analyze the genetic variation and LD patterns in asparagus bean based on a panel of 95 Chinese accessions genotyped using the above SNP markers.

MATERIALS AND METHODS

Plant materials

A sample set consisting of 95 asparagus bean accessions from a wide geographic origin across China and four ssp. *unguiculata* accessions (as a control) were used in the current study. Details of the plant materials including names, origins, subspecies assignment and morphological characteristics can be found in Supplementary Table 1.

DNA extraction

Genomic DNA was extracted from leaves of 2-week-old plants using a DNeasy Plant DNA miniprep kit (Qiagen, Hilden, Germany) according to the procedures described by the manufacturer.

SNP genotyping

Each of the 99 accessions was genotyped for SNPs using the KASPar (KBiosciences, Hoddeston, UK) 1127-SNP genotyping platform converted from the Illumina cowpea GoldenGate 1536-SNP assay system (Muchero *et al.*, 2009). Names of the 1127 SNPs are listed in Supplementary Table 2 and their details including map positions information can be accessed at HarvEST (<http://www.harvest-web.org/hweb/bin/wc.dll?hwebProcess-hmain~&versid=68>). After fluorescence scanning of the reactions, the results were

interpreted by the software KlusterCaller 1.1 (KBiosciences), which can be accessed at the Dryad repository (doi:10.5061/dryad.6tv35cc2).

Inference of structure and kinship

Population structure was calculated using a Bayesian model-based clustering method implemented in the software STRUCTURE 2.3.3 (Pritchard *et al.*, 2000) using data from 422 informative SNPs (see results below). We ran STRUCTURE under the 'admixture model' with a burn-in period of 100 000 followed by 100 000 replications of Markov Chain Monte Carlo. Three independent runs each were performed with the number of clusters (K) varying from 1 to 10. A statistic ΔK based on the relative rate of change in the likelihood of the data between successive K values were used to determine the optimal number of clusters (Evanno *et al.*, 2005). Lines with probability of membership >70% were assigned to a subgroup. No *a priori* population information was used. Pairwise genetic distances were calculated using the software Powermarker 3.25 under the Nei 1983 model (Liu and Muse, 2005). Relative kinship matrix was constructed using the software SPAGeDi and negative values between two individuals were changed to 0 (Hardy and Vekemans, 2002).

Analysis of LD

LD was measured by calculating the square value of correlation coefficient (r^2) between each SNP pair with the software package TASSEL 2.1 (Bradbury *et al.*, 2007). Only SNP loci with minor allele frequency values above 0.1 and having at least 80% successful calls among the sample set were included further for LD analyses. *P*-values for each r^2 estimate were obtained with a two-sided Fisher's exact test as implemented in TASSEL. The critical r^2 for LD decay was determined by taking the parametric 95th percentile of the distribution of r^2 square root for all unlinked loci (Brescaghello and Sorrells, 2006). LD plots against map distance for each linkage group (LG) were generated in Microsoft Excel (Redmond, WA, USA), where only r^2 values with $P < 0.01$ were included. Map position of each marker was obtained from the improved cowpea consensus map, allowing including the maximum number of markers (Lucas *et al.*, 2011). The LD decay trend lines were generated using a window size based method (Yan *et al.*, 2009).

Analysis of LD variance components

The computer program LinkDos (Garnier-gere and Dillmann, 1992) was employed to partition the variance of LD into within- (D_{IS}^2 and D'_{ST}^2) and between- (D_{ST}^2 and D'_{IS}^2) subpopulation components, following Ohta's *D*-statistics estimation method (1982). The greater between-subpopulation than within-populations variance components would suggest that genetic drift has an important role in shaping observed patterns of LD. For each chromosome, all the marker pairs with an $R^2 \geq 0.2$ and $P < 0.01$ were included in Ohta's analysis and the results were averaged to give a comprehensive insight. Significance of the differences between within- and between-subpopulation variance components was tested by ANOVA.

RESULTS

Data quality and genome wide SNP diversity

The KASPar SNP assay harboring 1127 SNPs gave a technical success rate of 97.7%, which means that 1102 SNPs were successfully called in the sample set. Of these, eight SNPs showed >20% missing data, 247 SNP loci were found to be monomorphic in all 99 lines and 420 SNPs had a minor allele frequency value below 0.1, which were removed from further analyses. Only five SNP loci were heterozygous in >10% of the accessions, consistent with the self-pollinating nature of asparagus bean and confirming the high level of homozygosity in the sample set. A final total of 422 SNPs were used for further data analysis. Of these, 415 SNP markers having known map positions and representing 370 unique loci were included for creating the LD plots (Supplementary Table 3). All the 11 LGs were covered by these SNPs, with the highest (76 SNPs) and lowest (23 SNPs) numbers observed in LG 3 and LG 10, respectively. The average marker distance was 1.58 cM, ranging from 1.01 to 2.69 cM across the 11 LGs. The allele




Figure 1 Allele frequency of the SNPs. Only SNPs with a minor allele frequency (MAF) >0.1 are shown. SNP frequencies >0.5 were shown as 0.5.

frequency distribution of the SNPs, which are bi-allelic, showed a continuous pattern, with a peak position falling into the region of 0.1–0.18 (Figure 1).

Identification of clusters and relative kinship in the sample set

Clustering inference performed with K from 1 to 10 showed that the most significant change of likelihood occurred when K increased from 1 to 2, and the highest ΔK value was observed at $K=2$ followed by a drastic decline of ΔK from $K=3$ (Table 1, Supplementary Figure 1). This suggested that the 99 genotypes could be assigned into two subgroups. Using a probability of membership threshold of 70%, 43 and 33 lines were assigned into the two subgroups, respectively. The remaining 23 lines were considered as intermediates (Supplementary Table 1, Supplementary Figure 2).

In general there was no association between subgroups inferred from structure and the geographic origin of the materials, reflecting the probable extensive exchange of parental lines by breeders nationwide; however, we found strong associations of subgrouping with plant morphology and usage. For example, subgroup I was consisted entirely of those ‘standard vegetable’ type lines, which means they all have very long tender pods (mean 50.5 cm, median 50.2 cm) typical of commercial varieties for vegetable use and strong climbing growth habit. In contrast, subgroup II ($n=29$ excluding the four *ssp. unguiculata* lines) mainly comprised the ‘non-standard vegetable’ type lines, which in general have shorter pods (mean 30.7 cm, median 30.6 cm) with higher fiber content, oval-shaped seeds suitable for grain use and dwarf or bush-type plant architecture (Supplementary Table 1 and 4). The two subgroups were hereafter renamed as subgroup SV (standard vegetable) and NSV (non-standard vegetable), respectively. All the four *ssp. unguiculata* lines were grouped into the NSV subgroup, suggesting a closer genetic relationship between this subgroup and the subspecies *unguiculata*.

The pairwise genetic distance among the 99 genotypes ranged from 0.01 to 0.58, with an average of 0.32. The greatest genetic distance was observed between ‘Charleston Greenpack’, an improved cowpea (*ssp. unguiculata*) cultivar from the southern United States (Fery, 1998) and ‘Prince Charming’, an improved Chinese asparagus bean (*ssp. sesquipedialis*) cultivar. The average genetic distance within subgroup SV (0.22) was lower than that in subgroup NSV (0.37). Analysis of relative kinship showed that 3124 (69%) of all the pairwise kinship values were between 0 and 0.05, whereas 18.9 % of the values were above 0.25, indicating that most individual pairs are not or only weakly related whereas members of subsets have a considerable level of relatedness.

LD in the asparagus bean genome

As the 95 asparagus bean accessions could be divided into two distinct subgroups, pairwise LD estimates were performed within each sub-

Table 1 Average $\ln P(D)$, s.d. and ΔK values at K from 1 to 10 showing the population structure of the sample set

K	$\ln P(D)$	s.d.	ΔK
1	-39 598	1.0	-
2	-33 986	2.4	1527.1
3	-32 009	20.8	2.9
4	-30 116	203.9	1.9
5	-28 620	162.1	3.9
6	-27 756	116.1	1.8
7	-26 680	456.8	0.3
8	-25 746	429.4	0.3
9	-24 682	56.0	3.5
10	-23 815	197.3	-

gene pool. Globally, 3937 (4.6%) and 2909 (3.4%) of the total possible SNP locus pairs were in significant LD ($P<0.01$) in subgroup SV and NSV, respectively (Table 2). Of these, 756 and 628 of the significant associations were intrachromosomal, accounting for 8.6% and 7.1 % of the total possible intrachromosomal correlations. The proportion of significant LD among unlinked loci to the total possible interchromosomal correlations was lower, being 4.3% and 3% in the two subgroups, respectively. In both subgroups, an uneven distribution of LD among the 11 chromosomes was observed.

Intrachromosomal LD extended to a relatively long distance in asparagus bean, with the means of 15 (median 9.5) and 11.2 (median 3.5) cM in the two subgene pools, respectively. Significant LD was even observed at a distance over 60 cM, despite the very small proportions (2.4% and 1.7% of all the significant intrachromosomal associations in subgroup SV and NSV, respectively). The strength of LD was high in asparagus bean, as reflected by the mean r^2 of 0.39 and 0.34 in subgroup SV and NSV, respectively. Albeit variable among LGs, the level (r^2) and extent of LD were generally higher in subgroup SV than in subgroup NSV (Table 2, Figure 2).

Based on the parametric 95th percentile of the distribution of square-rooted r^2 values for unlinked markers, r^2 thresholds of 0.5 and 0.4 were adopted for estimating LD decay in subgroup SV and NSV, respectively. It was showed that, though high in r^2 , LD decayed rapidly with increasing genetic distance in both subgene pools. On the whole genome scale, LD decayed within 2 cM (Figure 2). At the chromosome level, the LD for LG 3, 4 and 6 all decayed at 0–2 cM in subgroup SV, and for LG 1, 2, 7, 9, 10 and 11 within 2.5–5 cM. LD appeared more complicated for LG 5 and LG 8, as indicated by the uneven decay along the chromosomes. In subgroup NSV, LD decayed more rapidly. Six of the eleven chromosomes (LG 1, 2, 4, 6, 7 and 8) had their LD decay distances near or shorter than 1 cM, whereas LD for the other five chromosomes all decayed within 2–3 cM.

LD partitioning

To allow for discrimination between genetic drift and epistatic selection, the different possible causes that create and maintain LD in natural populations, Ohta’s (1982) analysis was performed by comparing the variance components within- (D_{IS}^2 and D_{ST}^2) and between- (D_{ST}^2 and D'_{IS}^2) subpopulations. As shown in Table 3, D_{ST}^2 was significantly greater than D_{IS}^2 on the whole genome scale, whereas D'_{IS}^2 was significantly smaller than D'_{ST}^2 , suggesting that globally the effect of epistatic selection and genetic drift cannot be simply compared. Similar results also were observed at the chromosome level; however, LD in at least two of the 11 LGs, that is, LG 5 and 7

Table 2 Locus pairs in significant ($P < 0.01$) LD, r^2 values and extent of LD in the two subgene pools

LG	Gene pool	No. Loci	No. (%) of locus pair in LD	Range (mean, median) of r^2	Extent (mean, median) of LD (cM)
1	SV	38	77 (12.2)	0.05–1 (0.40, 0.37)	30 (8, 6.6)
	NSV		40 (6.3)	0.03–1 (0.33, 0.24)	57 (13.7, 2.5)
2	SV	56	57 (3.7)	0.02–1 (0.33, 0.21)	61 (24, 27)
	NSV		151 (9.8)	0.04–1 (0.33, 0.24)	72 (13.5, 6.5)
3	SV	76	204 (7.2)	0.01–1 (0.28, 0.2)	78 (23, 19)
	NSV		136 (4.8)	0.02–0.9 (0.25–0.2)	73 (14.4, 6.4)
4	SV	30	28 (6.4)	0.05–0.9 (0.29, 0.21)	35 (9.9, 8)
	NSV		27 (6.2)	0.06–0.87 (0.29, 0.24)	27 (6.2, 3.3)
5	SV	28	62 (16.4)	0.12–1 (0.57, 0.54)	50 (11, 9.5)
	NSV		21 (5.6)	0.09–1 (0.4, 0.39)	38 (6.4, 1.2)
6	SV	41	38 (4.6)	0.12–1 (0.48, 0.44)	42 (9.8, 7.3)
	NSV		67 (8.2)	0.04–0.9 (0.3, 0.25)	64 (17.3, 11.8)
7	SV	32	71 (14.3)	0.06–0.9 (0.4, 0.31)	26 (10, 7.5)
	NSV		23 (4.6)	0.04–0.6 (0.3, 0.26)	32 (7.3, 0.6)
8	SV	27	35 (10)	0.21–1 (0.53, 0.48)	37 (10.6, 6.9)
	NSV		32 (9.1)	0.06–0.9 (0.39, 0.34)	42 (5.1, 1.1)
9	SV	36	39 (6.2)	0.15–1 (0.46, 0.38)	40 (12.7, 10)
	NSV		36 (5.7)	0.02–1 (0.26, 0.17)	29 (9.2, 6)
10	SV	23	17 (6.7)	0.05–0.95 (0.4, 0.36)	42 (17.8, 8.8)
	NSV		13 (5.1)	0.06–0.87 (0.4, 0.37)	24 (4.4, 1.1)
11	SV	28	71 (18.8)	0.06–0.9 (0.46, 0.33)	31 (7, 10)
	NSV		48 (12.7)	0.18–0.87 (0.7, 0.86)	13 (1.2, 0.5)
Genome-wide	SV	415	3937 (4.6) ^a	0.01–1 (0.39, 0.3)	78 (15, 9.5)
	NSV		2909 (3.4) ^a	0.02–1 (0.34, 0.27)	66 (11.2, 3.5)

Abbreviations: NSV, non-standard vegetable; SV, standard vegetable.

^aThe numbers and proportions of locus pairs in linkage disequilibrium (LD) on the whole genome scale includes both intrachromosomal and interchromosomal associations.

may primarily be because of epistatic selection, because the within-subpopulations variances components were consistently greater than those between subpopulations (see also discussion below).

DISCUSSION

Genetic diversity and domestication history of Chinese asparagus bean germplasm

Unlike staple food crops such as rice and maize where abundant germplasm resources are available for LD analysis, the genetic resources of asparagus bean is limited. In the current study, 99 genotypes were selected, based on morphology, geographic origin and pedigree (if known), from the available Chinese asparagus bean germplasm to represent a putative core collection. Despite their diverse morphological characteristics and origins, the genetic diversity within the sample set is low, as reflected by the low level of SNP polymorphism and genetic distance. This result is consistent with previous phylogenetic studies based on AFLP or SSR markers (Fang *et al.*, 2007; Xu *et al.*, 2010). However, these results should be viewed with some caution because of potential ascertainment bias in the original selection of SNPs put on the platform used in the assays (Muchero *et al.*, 2009). Although highly variable in morphology, several authors provide evidence that cowpea went through a severe genetic bottleneck during domestication and therefore has lower inherent genetic diversity (Panella and Gepts 1992; Pasquet 1999). Asparagus bean (*ssp. unguiculata*), as a subspecies of cowpea originating from only a small portion of the genetic variation of domesticated *ssp. unguiculata* would have gone through an additional bottleneck due to founder effects and intense selection for pod characteristics favorable for vegetable use (Fang *et al.* 2007). Therefore, the low level of genetic diversity observed in this study across the Chinese asparagus bean germplasm fits well with its likely domestication history.

The classification of the 99 accessions into two subgroups that differ mainly in pod length and growth habit indicates the impact of long-term human selection toward traits for vegetable use on the differentiation of *ssp. unguiculata*/asparagus bean, as lines with pods shorter than 36 cm or dwarf/semi-dwarf architecture were grouped together with the four *ssp. unguiculata* accessions. The analysis of LD variance components also suggested that epistatic selection might be the main force shaping the current pattern of LD in at least part of the genome. Our results therefore provide solid evidence proving the long-existing hypothesis of the domestication history of cultivated asparagus bean as mentioned above from a population genetic perspective. We also note that three landrace accessions (No.7, 18 and 80) morphologically more fitting the 'standard vegetable' type were classified into the NSV subgroup. This observation, though common, indicates that genetic factors controlling morphological variations of pod length and growth habit are not the only forces dominating the stratification of Chinese asparagus bean germplasm. An interesting task in the near future is to more thoroughly address the mechanisms behind the domestication of cultivated asparagus bean.

Comparison of LD patterns between asparagus bean and other plant species

The LD level appears high in asparagus bean; however, decay of LD is still rapid (0–2 cM genome wide). Given the genome size of 630 Mb and the genetic linkage map of 680 cM (Lucas *et al.*, 2011; Xu *et al.*, 2011), this approximately equals 1.84 Mb of physical distance. Such level and extent of LD are comparable to those observed in the European barley collection (Rostoks *et al.*, 2006), and the LD decay distance is even shorter than that in a small population of common bean (*Phaseolus vulgaris* L.) from America, a close relative of asparagus




Figure 2 Decay of LD (r^2) as a function of genetic distance (cM) between pairs of loci on individual LGs and the whole genome in asparagus bean. Only r^2 values with $P < 0.01$ are shown.

bean, where LD decays to 0.1–0.2 within 6–12 cM (McConnell *et al.*, 2010). Collectively, these observations indicate that in inbreeding species including asparagus bean, the LD decay distances are fairly

long. In contrast, LD declines very rapidly in outcrossing species where physical recombination is more effective. For instance, LD decays within only a few kilobases in maize (Remington *et al.*, 2001;

Table 3 Overall and linkage group (LG)-based Ohta's variance components of LD

LG	D_{IS}^2	D_{ST}^2	D_{IS}^2/D_{ST}^2	D'_{ST}^2	D'_{IS}^2	D'_{ST}^2/D'_{IS}^2
1	0.060	0.067	0.89	0.241	0.052	4.63*
2	0.031	0.089	0.34*	0.316	0.033	9.57*
3	0.032	0.085	0.37*	0.319	0.036	8.86*
4	0.035	0.072	0.48*	0.263	0.037	7.1*
5	0.083	0.047	1.76*	0.214	0.058	3.68*
6	0.024	0.131	0.18*	0.486	0.040	12.15*
7	0.092	0.039	2.35*	0.160	0.065	2.46*
8	0.035	0.143	0.24*	0.493	0.074	6.66*
9	0.018	0.068	0.26*	0.261	0.019	13.73*
10	0.018	0.131	0.12*	0.456	0.040	11.4*
11	0.177	0.036	4.91*	0.157	0.160	0.98
Overall	0.041	0.087	0.47*	0.322	0.044	7.31*

*Significant at $P=0.01$.

Yan *et al.*, 2009) and only 200 bp in a wild sunflower population (Liu and Burke, 2006).

Variation in LD magnitudes across different germplasm and chromosomes

The extent of LD is usually affected by population structure. For example, Morrell *et al.* (2005) reported a rapid decay of LD in wild barley compared with domesticated barley, even though both are highly self-pollinated. Similar results were reported between the *indica* and *japonica* groups of rice (Mather *et al.*, 2007), the wild and domesticated subspecies of grapevine (Barnaud *et al.*, 2010) and the landrace and modern varieties groups of wheat (Hao *et al.*, 2011). Our results here showed that LD patterns in the two asparagus bean subdivisions are also different and that the rate of interchromosomal LD was reduced to, on average, 3.6% compared with 8% when using the combined gene pool (data not shown). The subgroup SV globally had a higher level and longer decay distance of LD, suggesting relatively limited historical recombination events and less genetic diversity in this more typical vegetable-type subdivision perhaps because of a more severe historical population bottleneck. The distances and patterns of LD decay are most different on LG 5, 7 and 11 between the two subgroups, suggesting that these chromosomes may carry more genes/QTLs related to agronomic traits favorable for vegetable use that have been strongly selected by cultivators and breeders of this crop. This is in good agreement with the Ohta's LD variance component analysis. An important future task is to fine-dissect the LD patterns between the two subgroups of asparagus bean germplasm using broader populations that may also include more *ssp. unguiculata* accessions to get better understanding of the genetic architecture of the cowpea natural resources globally.

Implications on genome wide association study in asparagus bean

For genome wide association mapping, long-range LD will reduce the number of markers needed to cover the genome, although the resolution for mapping is expected to be low (Yu and Buckler, 2006). The level of LD decay at a $-cM$ level in the Chinese asparagus bean germplasm we observed suggests that a genome wide detection of QTLs is feasible using the currently available SNP maps and marker system. Assuming that one marker is needed per 1.5 cM along the 680 cM asparagus bean genome, ~ 450 informative markers evenly distributed across the genome would be sufficient to uncover any significant associations using genome wide association studies. Given

the expressed sequence tag-derived nature of the SNP markers employed here, the power for association mapping is expected to be higher than using other marker systems such as genomic SSRs, of which the majority are located in the non-coding regions of the genome (Yan *et al.*, 2009). In addition, the presence of population stratification and kinship among a subset of the plant materials, as well as the 3.6% interchromosomal associations, all should be taken into account in association analyses (Yan *et al.*, 2009; Hao *et al.*, 2011).

To further complement the possible low mapping resolution caused by high-level LD in asparagus bean, a two-step procedure as proposed by Barnaud *et al.* (2010) would be considered for future work. Based on the discovery here that landrace asparagus bean lines (and especially the *ssp. unguiculata*), are genetically more diverse and thus might should exhibit shorter LD decay distance, QTL locations determined from 'standard' asparagus bean populations could be refined by studying populations of more 'rustic' asparagus bean or *ssp. unguiculata* accessions with much more limited LD. In addition, bi-parental QTL mapping using large RIL populations or NIL sets would also provide higher resolution for QTLs of interest. In our lab, a project that aims to uncover the genetic factors controlling the pod length and climbing ability of asparagus bean using the combined two-step association mapping and bi-parental QTL mapping approach is now underway.

DATA ARCHIVING

Data have been deposited at Dryad: doi:10.5061/dryad.6tv35cc2.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This study was partially supported by the Science Foundation of Zhejiang Province (No. Y3090146), the National Science Foundation of China (No. 30900985) and the Sino-US joint project funded by Zhejiang Academy of Agricultural Sciences, China. We are grateful to Pauline Garnier-Géré, for advising us on Ohta's LD variance components analysis and for valuable comments on earlier versions of the manuscript.

- Ardlie KG, Kruglyak L, Seielstad M (2002). Patterns of linkage disequilibrium in the human genome. *Nat Rev Genet* **3**: 299–309.
- Barnaud A, Laucou V, This P, Lacombe T, Doligez A (2010). Linkage disequilibrium in wild French grapevine, *Vitis vinifera* L. *subsp. Silvestris*. *Heredity* **104**: 431–437.
- Bradbury PJ, Zhang ZW, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES (2007). TASSEL: software for association mapping of complex traits in diverse samples. *Bioinformatics* **23**: 2633–2635.
- Breseghele F, Sorrells ME (2006). Association mapping of kernel size and milling quality in wheat (*Triticum aestivum* L. cultivars). *Genetics* **172**: 1165–1177.
- Cardon LR, Bell JI (2001). Association study designs for complex diseases. *Nat Rev Genet* **2**: 91–99.
- Ching A, Caldwell KS, Jung M, Dolan M, Smith OS, Tingey S *et al.* (2002). SNP frequency, haplotype structure and linkage disequilibrium in elite maize inbred lines. *BMC Genetics* **3**: 19.
- Evanno G, Regnaut S, Goudet J (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol* **14**: 2611–2620.
- Fang JG, Chao CCT, Roberts PA, Ehlers JD (2007). Genetic diversity of cowpea [*Vigna unguiculata* L. Walp.] in four West African and USA breeding programs as determined by AFLP analysis. *Genet Resour Crop Ev* **54**: 1197–1209.
- Fery RL (1998). Charleston Greenpack' a pinkeye-type southernpea with a green cotyledon phenotype. *HortSci* **33**: 907–908.
- Flint-Garcia SA, Thornsberry JM, Buckler ES (2003). Structure of linkage disequilibrium in plants. *Ann Rev Plant Biol* **54**: 357–374.
- Garnier-Gere P, Dillmann C (1992). A computer program for testing pairwise linkage disequilibria in subdivided populations. *J Hered* **83**: 239.
- Garris AJ, McCouch SR, Kresovich S (2003). Population structure and its effect on haplotype diversity and linkage disequilibrium surrounding the *xa5* locus of rice (*Oryza sativa* L.). *Genetics* **165**: 759–769.

- Hao C, Wang L, Ge H, Dong Y, Zhang X (2011). Genetic diversity and linkage disequilibrium in Chinese bread wheat (*Triticum aestivum* L.) revealed by SSR markers. *PLoS One* **6**: e17279.
- Hardy OJ, Vekemans X (2002). Spagedi: a versatile computer program to analyze spatial genetic structure at the individual or population levels. *Mol Ecol Notes* **2**: 618–620.
- Hartl DL, Clark AG (1997). *Principles of population genetics*. Sinauer Associates: Sunderland, MA.
- Lander ES, Schork NJ (1994). The genetic dissection of complex traits. *Science* **265**: 2037–2048.
- Liu A, Burke JM (2006). Patterns of nucleotide diversity in wild and cultivated sunflower. *Genetics* **173**: 321–330.
- Liu KJ, Muse SV (2005). PowerMarker: an integrated analysis environment for genetic marker analysis. *Bioinformatics* **21**: 2128–2129.
- Lucas MR, Diop N-N, Wanamaker S, Ehlers JD, Roberts PA, Close TJ (2011). Cowpea-soybean synteny clarified through an improved genetic map. *Plant Genome* **4**: 218–225.
- Mather KA, Caicedo AL, Polato NR, Olsen KM, McCouch S, Purugganan MD (2007). The extent of linkage disequilibrium in rice (*Oryza sativa* L.). *Genetics* **177**: 2223–2232.
- McConnell M, Mamidi S, Lee R, Chikara S, Rossi M, Papa R, McClean P (2010). Syntenic relationships among legumes revealed using a gene-based genetic linkage map of common bean (*Phaseolus vulgaris* L.). *Theor Appl Genet* **121**: 1103–1116.
- McNally KL, Childs KL, Bohnert R, Davidson RM, Zhao K, Ulat VJ *et al.* (2009). Genomewide SNP variation reveals relationships among landraces and modern varieties of rice. *PNAS* **106**: 12273–12278.
- Morrell PL, Toleno DM, Lundy KE, Clegg MT (2005). Low levels of linkage disequilibrium in wild barley (*Hordeum vulgare* ssp. *spontaneum*) despite high rates of self-fertilization. *PNAS* **102**: 2442–2447.
- Muchero W, Diop NN, Bhat PR, Fenton RD, Wanamaker S, Pottorff M *et al.* (2009). A consensus genetic map of cowpea (*Vigna unguiculata* L. Walp) and synteny based on EST-derived SNPs. *PNAS* **106**: 18159–18164.
- Ohta T (1982). Linkage disequilibrium due to random genetic drift in finite subdivided populations. *PNAS* **79**: 1940–1944.
- Panella L, Gepts P (1992). Genetic relationship within *Vigna unguiculata* L. Walp. based on isozyme analyses. *Genet Resour Crop Ev* **39**: 71–88.
- Pasquet RS (1999). Genetic relationships among subspecies of *Vigna unguiculata* L. Walp. based on allozyme variation. *Theor Appl Genet* **98**: 1104–1119.
- Pritchard JK, Stephens M, Donnelly P (2000). Inference of population structure from multilocus genotype data. *Genetics* **155**: 945–959.
- Remington DL, Thornsberry JM, Matsuoka Y, Wilson LM, Whitt SR, Doebley J *et al.* (2001). Structure of linkage disequilibrium and phenotypic associations in the maize genome. *PNAS* **98**: 11479–11484.
- Robbins MD, Sim SC, Yang W, Deynze AV, Knaap E, Joobeur T *et al.* (2011). Mapping and linkage disequilibrium analysis with a genome-wide collection of SNPs that detect polymorphism in cultivated tomato. *J Exp Bot* **62**: 1831–1845.
- Rostoks N, Ramsay L, MacKenzie K, Cardle L, Bhat PR, Roose ML *et al.* (2006). Recent history of artificial outcrossing facilitates whole-genome association mapping in elite inbred crop varieties. *PNAS* **103**: 18656–18661.
- Tenaillon MI, Sawkins MC, Long AD, Gaut RL, Doebley JF, Gaut BS (2001). Patterns of DNA sequence polymorphism along chromosome 1 of maize (*Zea mays* ssp. *mays* L.). *PNAS* **98**: 9161–9166.
- Timko MP, Ehlers JD, Roberts PA (2007). Cowpea. In: Kole C (ed) *Pulses, sugar and tuber crops, Genome mapping and molecular breeding in plants*, vol 3. Springer-Verlag: Berlin Heidelberg, pp 49–67.
- Weir B (1979). Inferences about linkage disequilibrium. *Biometrics* **35**: 235–254.
- Xu P, Wu X, Wang B, Liu Y, Ehlers JD, Close TJ *et al.* (2011). A SNP and SSR based genetic map of asparagus bean (*Vigna unguiculata* ssp. *sesquipedialis*) and comparison with the broader species. *PLoS One* **6**: e15952.
- Xu P, Wu XH, Wang BG, Liu YH, Qin DH, Ehlers JD *et al.* (2010). Development and polymorphism of *Vigna unguiculata* ssp. *unguiculata* microsatellite markers used for phylogenetic analysis in asparagus bean (*Vigna unguiculata* ssp. *sesquipedialis* L. Verdc.). *Mol Breeding* **25**: 675–684.
- Yan J, Shah T, Warburton ML, Buckler ES, McMullen MD, Crouch J (2009). Genetic characterization and linkage disequilibrium estimation of a global maize collection using SNP markers. *PLoS One* **4**: e8451.
- Yu J, Buckler ES (2006). Genetic association mapping and genome organization of maize. *Curr Opin Biotech* **17**: 155–160.

Supplementary Information accompanies the paper on Heredity website (<http://www.nature.com/hdy>)