

CORRIGENDUM

Expression of thyrotropin and thyroid hormone receptors in the adipose tissue of patients with morbid obesity and/or type 2 diabetes: effects of weight loss

M Nannipieri, F Cecchetti, M Anselmino, S Camastra, P Nicolini, M Lamacchia, M Rossi, G Iervasi and E Ferrannini

International Journal of Obesity (2010) 34, 215; doi:10.1038/ijo.2009.246

Correction to: *International Journal of Obesity* (2009) 33, 1001–1006; doi:10.1038/ijo.2009.140; published online 28 July 2009

After the publication of the above article, the authors noticed a mistake in Table 1. The correct table is reproduced below.

Table 1 Baseline clinical and metabolic characteristics

	<i>Obese-NGT</i>	<i>Obese-IGT</i>	<i>Obese-T2DM</i>	<i>Controls</i>
Subjects (<i>n</i>)	55	20	32	12
Sex (F/M) ^a	49/6	14/6	23/9	8/4
Age (years) ^a	40 ± 11	39 ± 10	48 ± 8	45 ± 10
BMI (kg m ⁻²) ^a	47.4 ± 9.7	46.9 ± 7.1	45.4 ± 6.4	23.6 ± 3.0
Waist girth (cm) ^a	124.2 ± 15.0	132.1 ± 9.9	130.0 ± 15.1	84.6 ± 8.2
SBP (mm Hg) ^a	126 ± 15	126 ± 17	139 ± 14	123 ± 10
DBP (mm Hg) ^a	77 ± 11	80 ± 11	86 ± 9	77 ± 7
TG (mg per 100 ml) ^a	112 [44]	135 [90]	186 [153]	121 [11]
T-Chol. (mg per 100 ml)	195 ± 33	207 ± 56	215 ± 52	192 ± 6
HDL-Chol. (mg per 100 ml)	49 ± 10	43 ± 11	45 ± 12	47 ± 2
FPG (mg per 100 ml) ^a	88 [11]	97 [16]	153 [40]	88 [14]
2-h PG (mg per 100 ml) ^a	112 [27]	166 [18]	243 [23]	113 [11]
FPI (μU ml ⁻¹) ^a	16.8 [12]	15.4 [16]	20.7 [25]	7.7 [4]
2-h PI (μU ml ⁻¹)	64 [79]	100 [63]	94 [115]	73 [42]
TSH (μU ml ⁻¹) ^a	1.49 [1.46]	1.43 [1.78]	2.00 [1.54]	0.80 [0.32]
FT ₃ (pg ml ⁻¹) ^a	3.7 [1.5]	3.8 [1.0]	3.5 [0.9]	2.7 [0.5]
FT ₄ (pg ml ⁻¹) ^a	10.2 [2.6]	9.6 [4.7]	10.7 [4.3]	12.5 [4.2]
FT ₄ /FT ₃	2.76	2.52	3.10	4.63

^a*P* ≤ 0.05 by Kruskal–Wallis test. Data are expressed as mean ± s.d. or median [interquartile range].

Also one of the author's name is not correct: Nicolini P should be Nicolini G, as the full name is Giuseppina Nicolini.

The authors would like to apologize for this mistake.