

www.nature.com/ismei

## **CORRIGENDUM**

## An active atmospheric methane sink in high Arctic mineral cryosols

MCY Lau, BT Stackhouse, AC Layton, A Chauhan, TA Vishnivetskaya, K Chourey, J Ronholm, NCS Mykytczuk, PC Bennett, G Lamarche-Gagnon, N Burton, WH Pollard, CR Omelon, DM Medvigy, RL Hettich, SM Pfiffner, LG Whyte and TC Onstott

The ISME Journal (2015) 9, 1904; doi:10.1038/ismej.2015.112

**Correction to**: *The ISME Journal* (2015) **9**, 1880–1891; doi:10.1038/ismej.2015.13; published online 14 April 2015

Since the publication of this article, the authors have identified an error in the reported values for  $CH_4$  oxidation rates calculated from the microcosm experiment (Table 1), which also affected the reported  $Q_{10}$  values (on page 7 and Figure 3):

At 66% and 100% saturations, the  $Q_{10}$  values fell within the range of 1.3–2.9 values previously reported from incubation experiments carried out

at atmospheric or elevated  $CH_4$  concentrations (Figure 3). In contrast, the  $Q_{10}$  value was 8 at 33% saturation (Figure 3).

The revised text is:

At 100% saturations, the  $Q_{10}$  values fell within the range of 1.3–2.9 values previously reported from incubation experiments carried out at atmospheric or elevated CH<sub>4</sub> concentrations (Figure 3). In contrast, the  $Q_{10}$  value was 6 at 33% and 66% saturations (Figure 3).

Table 1 and Figure 3 have been amended to reflect these changes.

The authors would like to apologize for any inconvenience this may have caused.