

## Xylactam, a New Nitrogen-containing Compound from the Fruiting Bodies of Ascomycete *Xylaria euglossa*

Xing-Na Wang, Ren-Xiang Tan, Ji-Kai Liu

Received: November 17, 2004 / Accepted: March 22, 2005

© Japan Antibiotics Research Association

**Abstract** A novel nitrogen-containing compound, named xylactam (**1**), was isolated from the fruiting bodies of ascomycete *Xylaria euglossa* together with two known compounds penochalasin B2 and neoechinulin A. Their structures were elucidated on the basis of spectral data.

**Keywords** *Xylaria euglossa*, xylactam, ascomycete

*Xylaria euglossa* Fr. is a rot-wood-inhabiting ascomycete, mainly occurring on stumps and fallen branches of forested areas in the Southwest of China [1]. Many unique secondary metabolites have been found in the fungi of this genus. During the study of *Xylaria* sp., various new metabolites had been discovered, including cytochalasins [2, 3], globoscin [4], lactones [5], maldoxin [6], sesquiterpenoids [7, 8], xylaramide [9], xylarin [10], and xyloketal [11]. As a part of our search for naturally occurring bioactive metabolites of the higher fungi in Yunnan province [12], we have carried out a detailed chemical investigation on the fungus *Xylaria euglossa* Fr. and isolated a new nitrogen-containing compound, xylactam (**1**), along with two known alkaloids, penochalasin B2 and neoechinulin A from extracts of the fruiting bodies.

Fruiting bodies of *Xylaria euglossa* were collected at Ailao mountain of Yunnan province, P. R. China, in July, 2002 and identified by Prof. Mu Zang, Kunming Institute of Botany, the Chinese Academy of Sciences. The voucher


specimen was deposited in the Herbarium of the Kunming Institute of Botany, the Chinese Academy of Sciences. The air-dried fruiting bodies of *X. euglossa* (0.5 kg) were crushed and extracted with chloroform/methanol (1/1, v/v) four times at room temperature. The combined extracts were concentrated *in vacuo* to give a syrup (25 g), which was partitioned between chloroform and water. The chloroform soluble part (17 g) was subjected to silica gel column chromatography employing a gradient elution with chloroform/methanol (from 100:0 to 85:15, v/v) to give six fractions. Fraction II (100:1, v/v) was chromatographed on silica gel (petroleum ether/acetone 85:15), yielding penochalasin B2 (30 mg). Fraction III (98:2, v/v) was rechromatographed through silica gel, eluting with petroleum ether/acetone 80:20 (v/v) to give neoechinulin A (5 mg). Fraction VI (85:15, v/v) was passed through Sephadex LH-20, eluting with chloroform/methanol (1:1, v/v). A fraction exhibited strong yellow and white-blue fluorescence at 254 nm and was further purified on silica gel in petroleum ether/acetone (7:3, v/v) to afford compound **1** (8 mg).

Xylactam (**1**) was obtained as a white powder. Its molecular formula was determined as C<sub>23</sub>H<sub>31</sub>NO<sub>6</sub> from the quasi-molecular ion peak at *m/z* 418.2211 ([M+H]<sup>+</sup>, calcd. 418.2229) in the positive-ion HR-ESI-MS. Twenty-three signals in the <sup>13</sup>C-NMR (DEPT) spectra of **1** were recognized (9×C, 2×CH, 11×CH<sub>2</sub>, 1×CH<sub>3</sub>), including three carbonyl C-atoms ( $\delta$  172.4, 173.5 and 204.0) and six aromatic quaternary C-atoms ( $\delta$  102.5, 120.8, 125.9, 128.2, 155.0, 164.8). To fulfill the molecular formula of

**J.-K. Liu** (Corresponding author), **X.-N. Wang**: Kunming Institute of Botany, the Chinese Academy of Sciences, Kunming 650204, P. R. China, E-mail: jkliu@mail.kib.ac.cn

**X.-N. Wang, R.-X. Tan**: Institute of Functional Biomolecule, School of Life Sciences, Nanjing University, Nanjing 210093, P. R. China


compound **1**, the presence of one benzene ring, three carbonyl carbons ( $-\text{COOH}$ ,  $-\text{CO}-$ ,  $-\text{CONH}-$ ) and two phenolic hydroxyl groups (the exchangeable proton signals at  $\delta_{\text{H}}$  10.02 and 13.78) was suggested. In the mass spectrum of **1** a significant fragment ion at  $m/z=181$  ( $\text{C}_{12}\text{H}_{21}\text{O}$ ) was recognized. The signals in the  $^1\text{H-NMR}$  spectrum for one  $\text{A}_2\text{B}_2$ -system at  $\delta$  2.97 and 3.12 were assigned to one isolated  $-\text{CH}_2\text{CH}_2-$  group, and signals at  $\delta$  6.13 (H-12, d,  $J=16$ ) and 7.02 (H-13) were consistent with the presence of one *trans* double bond in the structure. The correlations from H-9 ( $\delta_{\text{H}}$  2.97) and H-12 to C-11 ( $\delta_{\text{C}}$  204.0) were observed in the HMBC spectrum of **1**. All these data suggested the existence of a long chain  $\text{CH}_3(\text{CH}_2)_8\text{CH}=\text{CHCO}-\text{CH}_2\text{CH}_2-$  in the structure of **1**. The  $^1\text{H-NMR}$  spectrum also presented a characteristic amide NH at  $\delta$  6.81 (1H, exchangeable with  $\text{D}_2\text{O}$ ) and  $^{13}\text{C-NMR}$  (DEPT) spectrum of **1** showed the presence of an amide functionality at  $\delta$  173.5 (C-1) and  $\delta$  44.6 (C-3). In addition, the correlations between NH ( $\delta_{\text{H}}$  6.81) and C-3a ( $\delta_{\text{C}}$  125.9), C-7a ( $\delta_{\text{C}}$  128.2) was observed in the HMBC spectrum of **1**. Combined all these evidence and comparison with the data of xylaral (**2**) from *Xylaria polymorpha* previously reported [13], compound **1** can be considered as one substructure A with four substituent



**Fig. 1** Structure of xylactam (**1**) and xylaral (**2**).

groups.

The HMBC spectrum revealed that the exchangeable proton at  $\delta_{\text{H}}$  13.78 belonged to a phenolic OH group coupled to carbons at  $\delta_{\text{C}}$  102.5 (C-7) and 164.8 (C-6). The latter signal was attributed to a carbon bearing a phenolic hydroxyl, and the strong coupling of the OH proton to the carbon at  $\delta_{\text{C}}$  102.5 and 120.8 suggested the direct neighborhood of the OH group to the aromatic



**Table 1**  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR data and HMBC correlations of **1**

Position			HMBC	$^1\text{H}$ , $^1\text{H-COSY}$
1		173.5 (C)		
2	6.81 (s)		C-3a, C-7a	
3	4.48 (s)	44.6 ( $\text{CH}_2$ )	C-1, C-4, C-7a	
3a		125.9 (C)		
4		155.0 (C)		
5		120.8 (C)		
6		164.8 (C)		
7		102.5 (C)		
7a		128.2 (C)		
8		172.4 (C)		
9	2.97 (t, $J=5.2$ )	16.8 ( $\text{CH}_2$ )	C-4, C-6, C-11	H-10
10	3.12 (t, $J=5.2$ )	39.3 ( $\text{CH}_2$ )	C-5, C-11	H-9
11		204.0 (C)		
12	6.13 (d, $J=16$ )	129.0 (CH)	C-11, C-14	H-13
13	7.02 (m)	151.4 (CH)	C-11, C-14, C-15	H-12, H-14
14	2.20 (m)	32.6 ( $\text{CH}_2$ )	C-12, C-13	H-13, H-15
15	1.42 (m)	27.8 ( $\text{CH}_2$ )	C-13	H-14
16~19	1.24 (m)	29.1~29.4 ( $\text{CH}_2$ )		
20	1.24 (m)	31.8 ( $\text{CH}_2$ )		
21	1.24 (m)	22.6 ( $\text{CH}_2$ )		
22	0.84 (t, $J=6$ )	14.1 ( $\text{CH}_3$ )	C-20	H-21
4-OH	10.02 (s)		C-5, C-3a, C-4	
6-OH	13.78 (s)		C-5, C-7, C-6	

carbons. The exchangeable proton at  $\delta_{\text{H}}$  10.02 exhibited strong HMBC correlations to carbons at  $\delta_{\text{C}}$  120.8 (C-5) and 125.9 (C-3a) as well as 155.0 (C-4). The HMBC spectrum also displayed correlations between H-3 ( $\delta_{\text{H}}$  4.48) and C-1 ( $\delta_{\text{C}}$  173.5), C-4 ( $\delta_{\text{C}}$  155.0), C-7a ( $\delta_{\text{C}}$  128.2), and also between H-9 ( $\delta_{\text{H}}$  2.97) and C-4 ( $\delta_{\text{C}}$  155.0), C-6 ( $\delta_{\text{C}}$  164.8). A combination of the substructure A and substituents leads to structure **1** for xylactam.

The spectral data and physical properties of penochalasin B2 and neoechinulin A were identical with the data previously reported [14, 15]. Penochalasin B was reported to exhibit potent cytotoxicity against cultured P 388 cells [14].

Xylactam (**1**). White powder. EI-MS: 417 (9,  $[\text{M}]^+$ ), 399 (5,  $[\text{M}-\text{H}_2\text{O}]^+$ ), 246 (19), 236 (34), 218 (100), 204 (20), 181 (45). FAB-MS (posit.): 418 (100,  $[\text{M}+1]^+$ ), 400 (40,  $[\text{M}+1-\text{H}_2\text{O}]^+$ ). HR-ESI-MS (posit.): 418.2211 ( $[\text{M}+1]^+$ , calc. 418.2229). UV ( $\text{CHCl}_3$ ): 242, 277, 325. IR (KBr): 3432, 3190, 2953, 2926, 2854, 1655, 1601, 1490, 1450, 1391, 1338, 1295, 1227, 1105, 766.  $^1\text{H}$ -,  $^{13}\text{C}$ -NMR, HMBC, and  $^1\text{H}$ - $^1\text{H}$  COSY: Table 1.

**Acknowledgements** This project was supported by the National Natural Science Foundation of China (30470027 and 30225048).

## References

- Mao XL. The Macrofungi in China, pp. 571, Henan Sciences and Technology Press, Henan, 2000
- Espada A, Rivera-Sagredo A, De La Fuente JM, Hueso-Rodriguez JA, Elson SW. New cytochalasins from the fungus *Xylaria hypoxylon*. *Tetrahedron* 53: 6485–6492 (1997)
- Dagne E, Gunatilaka AAL, Asmellash S, Abate D, Kingston DGI, Hofmann GA, Johnson RK. Two new cytotoxic cytochalasins from *Xylaria obovata*. *Tetrahedron* 50: 5615–5620 (1994)
- Adeboya M, Edwards RL, Lassoe T, Maitland DJ, Whalley A. Metabolites of the higher fungi. Part 28. Globoscinic acid and globoscin, a labile acid-lactone system from *Xylaria globosa* and *Xylaria obovata*. *J Chem Soc Perkin Trans 1* 1995: 2067–2072 (1995)
- Boonphong S, Kittakoop P, Isaka M, Pittayakhajonwut D, Tanticharoen M, Thebtaranonth Y. Multiplolides A and B, new antifungal 10-membered lactones from *Xylaria multiplex*. *J Nat Prod* 64: 965–967 (2001)
- Adeboya M, Edwards RL, Lassoe T, Maitland DJ, Shields L, Whalley A. Metabolites of the higher fungi. Part 29. Maldoxin, maldoxone, dihydromaldoxin, isodihydromaldoxin and dechlorodihydromaldoxin. A spirocyclohexadienone, a depsidone and three diphenylethers: keys in the depsidone biosynthetic pathway from a member of the fungus genus *Xylaria*. *J Chem Soc Perkin Trans 1* 1996: 1419–1425 (1996)
- Smith CJ, Morin NR, Bills GF, Dombrowski AW, Salituro GM, Smith SK, Zhao A, Macneil DJ. Novel sesquiterpenoids from the fermentation of *Xylaria persicaria* and selective ligands for the NPY Y5 receptor. *J Org Chem* 67: 5001–5004 (2002)
- Singh SB, Zink D, Polishhook J, Valentino D, Shafiee A, Silverman K, Felock P, Teran A, Vilella D, Hazuda DJ, Lingham RB. Structure and absolute stereochemistry of HIV-1 integrase inhibitor integric acid. A novel eremophilane sesquiterpenoid produced by a *Xylaria* sp. *Tetrahedron Lett* 40: 8775–8779 (1999)
- Schneider G, Anke H, Sterner O. Xylaramide, a new antifungal compound, and other secondary metabolites from *Xylaria longipes*. *Z Naturforsch* 51c: 802–806 (1996)
- Schneider G, Anke H, Sterner O. Xylarin, an antifungal *Xylaria* metabolite with an unusual tricyclic uronic acid moiety. *Nat Prod Lett* 7: 309–316 (1995)
- Lin Y, Wu X, Feng S, Jiang G, Luo J, Zhou S, Vrijmoed LLP, Jones EBG, Steingrover K, Zasila F. Five unique compounds: xyloketal from mangrove fungus *Xylaria* sp. from the South China Sea coast. *J Org Chem* 66: 6252–6256 (2001)
- Liu JK. Biologically active substances from mushrooms in Yunnan, China. *Heterocycles* 57: 157–167 (2002)
- Gunawan S, Steffan B, Steglich W. Xylaral, ein Hydroxyphthalid-Derivat aus Fruchtkoerpern von *Xylaria polymorpha* (Ascomycetes). *Liebigs Ann Chem* 1990: 825–827 (1990)
- Hiromichi N, Akira I, Akinori S, Saburo T.  $^{13}\text{C}$ -NMR spectra and stereochemistry of isoechinulins A, B and C. *Agric Biol Chem* 43: 1759–1763 (1979)
- Numata A, Takahashi C, Ito Y, Minoura K, Yamada T, Matsuda C, Nomoto K. Penochalasin, a novel class of cytotoxic cytochalasins from a *Penicillium* species separated from a marine alga: structure determination and solution confirmation. *J Chem Soc Perkin Trans 1* 1996: 239–246 (1996)