

Table 3 (Revised). *Continued.*

Analyte	Surface (floor) wipe (ng/cm ²)				Toy wipe (ng/cm ²)				Cotton socks ^d (ng/sample)			
	p25	p50	P75	Range	p25	p50	p75	Range	p25	p50	p75	Range
Bifenthrin	—	—	—	ND–0.035	—	—	—	ND–0.14	—	—	19	ND–92
λ -Cyhalothrin	—	—	—	ND–0.026	—	—	—	—	—	—	—	—
Cyfluthrin	—	—	—	ND–0.40	—	—	—	—	—	—	—	ND–1,400
Cypermethrin	—	—	0.34	ND–2.8	—	—	—	—	—	—	—	ND–2,400
Deltamethrin	—	—	—	—	—	—	—	—	—	—	—	—
Esfenvalerate	—	—	—	—	—	—	—	—	—	—	—	ND–230
<i>cis</i> -Permethrin	0.053	0.10	0.21	ND–1.7	—	—	—	ND–0.053	50	120	380	17–5,600
<i>trans</i> -Permethrin	0.14	0.23	0.39	ND–3.6	—	—	—	ND–0.072	140	220	490	10–9,200
Resmethrin	—	—	—	—	—	—	—	—	—	—	—	ND–130
Sumithrin	—	—	—	ND–3.5	—	—	—	—	—	—	—	ND–2,100
<i>Other</i>												
Chlorthal-dimethyl (herbicide)	0.022	0.044	0.079	0.0018–0.28	—	0.0059	0.012	ND–0.035	9.8	16	31	ND–200
<i>p,p'</i> -DDE (OC)	—	—	—	ND–0.15	—	—	—	—	—	28	52	ND–86
<i>p,p'</i> -DDT (OC)	—	—	—	—	—	—	—	—	—	—	—	ND–13,000
Iprodione (fungicide)	—	—	—	ND–0.53	—	—	—	—	—	—	—	ND–11,000

Abbreviations: “—” and ND = nondetectable; OC = organochlorine

^aAnalyte concentrations presented only when detectable levels were measured in one or more matrices.

^bHouse dust samples are collected from 1 m² surface.

^cOne missing *trans*-permethrin house dust sample measurement.

^dOne participant's sock sample was missing.

Table 6 (Revised). Spearman's rank sum correlation matrix for the five most frequently detected analytes across sampling media.^a

	Dust	Socks	Union suit	Surface wipe	Indoor air	Outdoor air	Toy	Overnight urine	Spot urine
<i>Chlorpyrifos</i>									
Dust	1								
Socks	–0.25	1							
Union suit	–0.24	0.54*	1						
Surface wipe	0.14	–0.056	0.16	1					
Indoor air	0.42	–0.12	–0.044	–0.066	1				
Outdoor air	–0.020	0.077	–0.093	–0.31	0.31	1			
Toy	–0.087	–0.19	–0.34	0.17	0.43	–0.14	1		
Overnight urine	–0.24	0.057	–0.17	0.021	0.084	0.045	0.50*	1	
Spot urine	–0.39	0.12	0.19	–0.091	–0.18	0.21	0.056	0.48*	1
<i>Chlorthal-dimethyl</i>									
Dust	1								
Socks	0.39	1							
Union suit	0.50*	0.76*	1						
Surface wipe	0.053	0.37	0.080	1					
Indoor air	0.66*	0.49*	0.63*	0.00	1				
Outdoor air	0.56*	0.31	0.57*	0.21	0.92*	1			
Toy	0.075	–0.043	0.13	–0.22	0.37	0.25	1		
<i>Diazinon</i>									
Dust	1								
Socks	0.52*	1							
Union suit	0.47*	0.76*	1						
Surface wipe	0.29	0.25	0.16	1					
Indoor air	0.47*	0.60*	0.62*	0.36	1				
Outdoor air	0.14	0.27	0.51*	0.30	0.80*	1			
Toy	0.33	0.76*	0.52*	0.28	0.62*	0.33	1		
Overnight urine	0.49*	0.25	0.069	0.081	–0.0033	–0.24	0.33	1	
Spot urine	0.40	0.11	0.27	–0.14	–0.0053	–0.069	0.20	0.48*	1
<i>cis-Permethrin</i>									
Dust	1								
Socks	0.43	1							
Union suit	0.40	0.18	1						
Surface wipe	0.39	0.44	0.32	1					
Indoor air	0.062	0.11	0.18	0.16	1				
Outdoor air	–0.16	0.12	–0.41	–0.27	–0.026	1			
Toy	0.22	0.30	0.036	0.061	0.032	–0.29	1		
<i>trans-Permethrin</i>									
Dust	1								
Socks	0.58*	1							
Union suit	0.52*	0.45	1						
Surface wipe	0.40	0.34	0.26	1					
Indoor air	0.39	0.30	0.13	0.20	1				
Outdoor air	–0.34	–0.42	–0.17	–0.19	–0.26	1			
Toy	0.33	0.44	0.13	–0.098	0.23	–0.16	1		

*Statistically significant Spearman's ρ ($P < 0.05$).

^aSpot and overnight urine samples are total diethyl phosphate metabolite levels (nmol/l).