

## CORRIGENDUM

### Preferential sensitivity of hematopoietic (HPs) and mesenchymal (MPs) progenitors to fludarabine suggests impaired bone marrow niche and HP mobilization

MG Berger<sup>1,2,3</sup>, J Berger<sup>1,2,3</sup>, C Richard<sup>1,2,3</sup>, S Jeanpierre<sup>4,5,6,7</sup>, FE Nicolini<sup>8</sup>, O Tournilhac<sup>1,3</sup>, M Michallet<sup>8</sup> and V Maguer-Satta<sup>4,5,6,7</sup>

<sup>1</sup>Hematology (Biology), CHU Hôtel-Dieu, Clermont-Ferrand, France; <sup>2</sup>UPRES EA 3846, Faculté Médecine, Université de Clermont 1, Clermont-Ferrand, France; <sup>3</sup>Department of Hematology and Cell Therapy, CHU Hôtel-Dieu, Clermont-Ferrand, France; <sup>4</sup>Université de Lyon, Lyon, France; <sup>5</sup>ISPB, Université de Lyon 1, Lyon, France; <sup>6</sup>INSERM, U590, Lyon, France; <sup>7</sup>IFR62, Lyon, France and <sup>8</sup>Department of Hematology, E Herriot Hospital, Lyon, France

*Leukemia* (2008) **22**, 2135; doi:10.1038/leu.2008.202

**Correction to:** *Leukemia* advance online publication, 3 July 2008; doi:10.1038/leu.2008.167

Since the publication of the above paper, the authors have

noticed errors in Dr V Maguer-Satta's name and also in the affiliations. The correct name and affiliations are shown above.

The authors apologize for any inconvenience caused.